

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
23 August 2001 (23.08.2001)

PCT

(10) International Publication Number
WO 01/60416 A2

- (51) International Patent Classification⁷: **A61K 49/00** (74) Agent: **DOLAN, Peter, L.**; DuPont Pharmaceuticals Company, Legal Patent Records Center, 1007 Market Street, Wilmington, DE 19898 (US).
- (21) International Application Number: **PCT/US01/04870**
- (22) International Filing Date: 15 February 2001 (15.02.2001) (81) Designated States (*national*): AU, BR, CA, CN, CZ, EE, HU, IL, IN, JP, KR, LT, LV, MX, NO, NZ, PL, RO, SG, SI, SK, UA, VN, ZA.
- (25) Filing Language: English
- (26) Publication Language: English (84) Designated States (*regional*): Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR).
- (30) Priority Data:
60/182,712 15 February 2000 (15.02.2000) US
- (71) Applicant: **DUPONT PHARMACEUTICALS COMPANY** [US/US]; 974 Centre Road, Wilmington, DE 19805 (US). Published:
— *without international search report and to be republished upon receipt of that report*
- (72) Inventors: **CARPENTER, Alan, P., Jr.**; 73 Cranberry Hill Lane, Carlisle, MA 01741 (US). **RAJOPADHYE, Milind**; 21 Honeysuckle Road, Westford, MA 01886 (US). *For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

WO 01/60416 A2

(54) Title: **MATRIX METALLOPROTEINASE INHIBITORS**

(57) Abstract: Thus the present invention describes diagnostic agents comprising a diagnostic metal and a compound, wherein the compound comprises: 1-10 targeting moieties; a chelator; and 0-1 linking groups between the targeting moiety and chelator; wherein the targeting moiety is a matrix metalloproteinase inhibitor; and wherein the chelator is capable of conjugating to the diagnostic metal. The present invention also provides novel compositions of the compounds of the invention, kits, and their uses in diagnosis of diseases associated with MMPs.

BEST AVAILABLE COPY

TITLE

MATRIX METALLOPROTEINASE INHIBITORS

5

FIELD OF THE INVENTION

The present invention provides novel compounds useful for the diagnosis of cardiovascular pathologies associated with extracellular matrix degradation, such as atherosclerosis, heart failure, and restenosis, methods of imaging these pathologies in
10 a patient, and pharmaceutical compositions comprising the compounds. The pharmaceuticals are comprised of a targeting moiety that inhibits a matrix metalloproteinase that is expressed in these pathologies, an optional linking group, and a diagnostically effective imageable moiety. The imageable moiety
15 is a gamma ray or positron emitting radioisotope, a magnetic resonance imaging contrast agent, an X-ray contrast agent, or an ultrasound contrast agent.

20

BACKGROUND OF THE INVENTION

The ability to detect increased levels of matrix metalloproteinases (MMPs) in the heart would be extremely useful for the detection of tissue degradation which occurs in many heart conditions. The composition and vulnerability of
25 atheromatous plaque in the coronary arteries has recently been recognized as a key determinant in thrombus-mediated acute coronary events, such as unstable angina, myocardial infarction and death (Falk E, Shah PK and Fuster V; Circulation 1995; 92: 657-671). Among the many components involved in the
30 inflammatory atheromatous plaque are macrophages which secrete the matrix metalloproteinases (Davies MJ; Circulation 1996; 94: 2013-2020). The MMPs are a family of enzymes which specialize in the cleavage of the usually protease-resistant fibrillar extracellular matrix components of the heart, such as collagen.
35 These extracellular matrix proteins confer strength to the

fibrous cap of atheroma (Libby P, Circulation 1995; 91: 2844-2850).

Macrophages which accumulate in areas of inflammation such as atherosclerotic plaques release these MMPs which degrade
5 connective tissue matrix proteins (Falk, 1995). In fact, studies have demonstrated that both the metalloproteinases and their mRNA are present in atherosclerotic plaques (Coker ML, Thomas CV, Clair MJ, et al.; Am. J. Physiol. 1998; 274:H1516-1523; Dollery CM, McEwan JR, Henney A., et al.; Circ. Res. 1995;
10 77:863-868; Henney A., Wakeley P., Davies M., et al.; Proc Natl Acad Sci 1991; 88:8154-8158), particularly in the vulnerable regions of human atherosclerotic plaques (Galis Z, Sukhova G, Lark M. and Libby P.; J Clin Invest. 1994; 94: 2493-2503). Amongst the metalloproteinases that may be released by
15 macrophages present at the site of human atheroma are interstitial collagenase (MMP-1), gelatinases A and B (MMP-2 and MMP-9, respectively) and stromelysin (MMP-3 Moreno PR, Falk E., Palacios IF et al.; Circulation 1994; 90: 775-778). Although all MMPs may be elevated at the site of human atheroma, it has
20 been suggested that gelatinase B may be one of the most prevalent MMPs in the plaque because it can be expressed by virtually all activated macrophages (Brown D., Hibbs M, Kerney M., et al.; Circulation 1995; 91: 2125-2131). The MMP-9 has also been shown to be more prevalent in atherectomy material
25 from unstable angina relative to stable angina patients (Brown, 1995).

The left ventricular extracellular matrix, containing a variety of collagens and elastin, are also proposed to participate in the maintenance of left ventricle (LV) geometry.
30 Therefore, alterations in these extracellular components of the myocardium may influence LV function and be a marker of progressive changes associated with LV degeneration and ultimately heart failure (Coker, 1998).

In the situation of congestive heart failure (CHF), the
35 relationship of CHF state to MMP activity in the LV remains somewhat unclear, at least in the clinical setting. In pre-

clinical models of CHF, however, the functional changes in the LV have been correlated with increased MMP activity. For example, in a pig model of CHF, the decrease in LV function was observed to coincide with a marked increase in MMP-1 (~300%),
5 MMP-2 (~200%), and MMP-3 (500%) (Coker, 1998). Moderate ischemia and reperfusion in a pig model has been demonstrated to selectively activate MMP-9 (Lu L, et al., Circulation, 1999, 100 Suppl. 1, I-12). Similarly in a dog model of CHF the levels of gelatinases (e.g. MMP-2 and MMP-9) were found to be elevated in
10 severe heart failure (Armstrong PW, Moe GW, et al., Can J Cardiol 1994; 10: 214-220). The levels of MMP-2 and MT1-MMP (membrane type MMP, MMP-14) were found to be increased in biopsy samples of human myocytes from patients suffering from dilated cardiomyopathy (Bond BR, et al., Circulation, 1999, 100 Suppl.
15 1, I-12).

Ahrens, et al. United States Patent No. 5,674,754 discloses methods for the detection of Matrix Metallo-Proteinase No. 9, using antibodies which selectively recognize pro-MMP-9 and complexes of pro-MMP-9 with tissue inhibitor of matrix
20 metallo proteinase-1 (TIMP-1), with no substantial binding to active MMP-9. Venkatesan, et al. United States Patent No. 6,172,057 discloses non-peptide inhibitors of matrix metalloproteinases (MMPs) and TNF-.alpha. converting enzyme (TACE) for the treatment of arthritis, tumor metastasis, tissue
25 ulceration, abnormal wound healing, periodontal disease, bone disease, diabetes (insulinresistance) and HIV infection.

Pathologically, MMPs have been identified as associated with several disease states. For example, anomalous MMP-2 levels have been detected in lung cancer patients, where it was
30 observed that serum MMP-2 levels were significantly elevated in stage IV disease and in those patients with distant metastases as compared to normal sera values (Garbisa et al., 1992, Cancer Res., 53: 4548, incorporated herein by reference.). Also, it was observed that plasma levels of MMP-9 were elevated in patients
35 with colon and breast cancer (Zucker et al., 1993, Cancer Res. 53: 140 incorporated herein by reference).

Elevated levels of stromelysin (MMP-3) and interstitial collagenase (MMP-1) have been noted in synovial fluid derived from rheumatoid arthritis patients as compared to post-traumatic knee injury (Walakovits et al., 1992, Arth. Rheum., 35: 35) incorporated herein by reference.. Increased levels of mRNA expression for collagenase type I (MMP-1) and collagenase type IV (MMP-2) have been shown to be increased in ulcerative colitis as compared to Crohn's disease and controls (Matthes et al., 1992, Gastroenterology, Abstract 661, incorporated herein by reference). Furthermore, Anthony et al., 1992, Gastroenterology, Abstract 591, demonstrated increased immuno-histochemical expression of the gelatinase antigen in arabbit model of chronic inflammatory colitis.

It has been shown that the gelatinase MMPs are most intimately involved with the growth and spread of tumors. It is known that the level of expression of gelatinase is elevated in malignancies, and that gelatinase can degrade the basement membrane which leads to tumor metastasis. Angiogenesis, required for the growth of solid tumors, has also recently been shown to have a gelatinase component to its pathology. Furthermore, there is evidence to suggest that gelatinase is involved in plaque rupture associated with atherosclerosis. Other conditions mediated by MMPs are restenosis, MMP-mediated osteopenias, inflammatory diseases of the central nervous system, skin aging, tumor growth, osteoarthritis, rheumatoid arthritis, septic arthritis, corneal ulceration, abnormal wound healing, bone disease, proteinuria, aneurysmal aortic disease, degenerative cartilage loss following traumatic joint injury, demyelinating diseases of the nervous system, cirrhosis of the liver, glomerular disease of the kidney, premature rupture of fetal membranes, inflammatory bowel disease, periodontal disease, age related macular degeneration, diabetic retinopathy, proliferative vitreoretinopathy, retinopathy of prematurity, ocular inflammation, keratoconus, Sjogren's syndrome, myopia, ocular tumors, ocular angiogenesis/neo-vascularization and corneal graft rejection. For recent reviews, see: (1) Recent Advances in

Matrix Metalloproteinase Inhibitor Research, R. P. Beckett, A. H. Davidson, A. H. Drummond, P. Huxley and M. Whittaker, Research Focus, Vol. 1, 16-26, (1996), (2) Curr. Opin. Ther. Patents (1994) 4(1): 7-16, (3) Curr. Medicinal Chem. (1995) 2: 743-762, (4) Exp. Opin. Ther. Patents (1995) 5(2): 1087-110, (5) Exp. Opin. Ther. Patents (1995) 5(12): 1287-1196, all of which are incorporated herein by reference.

Therefore, the present imaging agents targeted to one or more MMP's would be very useful for detecting and monitoring the degree of extracellular matrix degradation in CHF, atherosclerosis and other degradative disease processes. These imaging agents, containing a ligand directed at one or more MMP's (e.g. MMP-1, MMP-2, MMP-3, MMP-9), will localize a diagnostic imaging probe to the site of pathology for the purpose of non-invasive imaging of these diseases. The imaging agent may be a MMP inhibitor linked to radioisotopes which are known to be useful for imaging by gamma scintigraphy or positron emission tomography (PET). Alternatively, the MMP targeting ligand could be bound to a single or multiple chelator moieties for attachment of one or more paramagnetic metal atoms, which would cause a local change in magnetic properties, such as relaxivity or susceptibility, at the site of tissue damage, which could then be imaged with magnetic resonance imaging systems. Alternatively, the MMP inhibitor can be bound to a phospholipid or polymer material which would be used to encapsulate/stabilize microspheres of gas which would be detectable by ultrasound imaging following localization at the site of tissue injury.

Therefore, imaging agents based on MMP inhibitors would be extremely useful in the detection, staging and monitoring of cardiovascular diseases such as atherosclerosis (especially unstable arterial plaque) and various cardiomyopathies including congestive heart failure. Compounds of the present invention, which localize in areas of MMP activity in the heart, will allow detection and localization of these cardiac diseases which are

associated with altered MMP levels relative to normal myocardial tissue.

These imaging agents, whether for gamma scintigraphy, positron emission tomography, MRI, ultrasound or x-ray image enhancement, have utility to detect and monitor changes in cardiovascular diseases over time. Since the degree of overexpression of MMPs is related to the degradation of cardiac or vascular tissue (Rohde LE, Aikawa M, Cheng GC, et al., JACC 1999; 33: 835-842) it is possible to assess the severity and current activity of cardiovascular disease lesions (i.e. plaques) by quantitating the degree of localization of these imaging agents at the diseased sites of interest. Moreover, with these imaging agents it is possible to monitor changes in MMP activity associated with the institution of pharmaceutical therapies which slow the progression or cause a reversal of atherosclerotic changes in the vascular system or a reversal of myocardial degradation associated with congestive heart failure.

Therefore, it can be appreciated that the imaging of MMPs in the heart would be generally useful for the detection, localization and monitoring the progression/regression of a variety of cardiac diseases which are associated with alterations in the MMP content of cardiac tissues.

SUMMARY OF THE INVENTION

It is one object of the present invention to provide imaging agents for cardiovascular pathologies associated with extracellular matrix degradation, such as atherosclerosis, heart failure, and restenosis, comprised of matrix metalloproteinase inhibiting compounds conjugated to an imageable moiety, such as a gamma ray or positron emitting radioisotope, a magnetic resonance imaging contrast agent, an X-ray contrast agent, or an ultrasound contrast agent.

Another aspect of the present invention are diagnostic kits for the preparation of radiopharmaceuticals useful as imaging agents for cardiovascular pathologies associated with extracellular matrix degradation, such as atherosclerosis, heart

failure, and restenosis. Diagnostic kits of the present invention comprise one or more vials containing the sterile, non-pyrogenic, formulation comprised of a predetermined amount of a reagent of the present invention, and optionally other
5 components such as one or two ancillary ligands, reducing agents, transfer ligands, buffers, lyophilization aids, stabilization aids, solubilization aids and bacteriostats. The inclusion of one or more optional components in the formulation will frequently improve the ease of synthesis of the
10 radiopharmaceutical by the practicing end user, the ease of manufacturing the kit, the shelf-life of the kit, or the stability and shelf-life of the radiopharmaceutical. The inclusion of one or two ancillary ligands is required for diagnostic kits comprising reagent comprising a hydrazine or
15 hydrazone bonding moiety. The one or more vials that contain all or part of the formulation can independently be in the form of a sterile solution or a lyophilized solid.

Another aspect of the present invention contemplates a method of imaging cardiovascular pathologies associated with
20 extracellular matrix degradation, such as atherosclerosis, heart failure, and restenosis in a patient involving: (1) synthesizing a diagnostic radiopharmaceutical of the present invention, using a reagent of the present invention, capable of localizing at the loci of the cardiovascular pathology; (2) administering said
25 radiopharmaceutical to a patient by injection or infusion; (3) imaging the patient using planar or SPECT gamma scintigraphy, or positron emission tomography.

Another aspect of the present invention contemplates a method of imaging cardiovascular pathologies associated with
30 extracellular matrix degradation, such as atherosclerosis, heart failure, and restenosis in a patient involving: (1) administering a paramagnetic metallopharmaceutical of the present invention capable of localizing the loci of the cardiovascular pathology to a patient by injection or infusion;
35 and (2) imaging the patient using magnetic resonance imaging.

Another aspect of the present invention contemplates a method of imaging cardiovascular pathologies associated with extracellular matrix degradation, such as atherosclerosis, heart failure, and restenosis in a patient involving: (1)

- 5 administering a X-ray contrast agent of the present invention capable of localizing in tumors to a patient by injection or infusion; and (2) imaging the patient using X-ray computed tomography.

- Another aspect of the present invention contemplates a
10 method of imaging cardiovascular pathologies associated with extracellular matrix degradation, such as atherosclerosis, heart failure, and restenosis in a patient involving: (1)
administering a ultrasound contrast agent of the present invention capable of localizing in tumors to a patient by
15 injection or infusion; and (2) imaging the patient using sonography.

DETAILED DESCRIPTION

- 20 Thus the present invention includes the following embodiments:

(1) A diagnostic agent comprising a diagnostic metal and a compound, wherein the compound comprises:

- i) 1-10 targeting moieties;
25 ii) a chelator; and
iii) 0-1 linking groups between the targeting moiety and chelator;

wherein the targeting moiety is a matrix metalloproteinase inhibitor; and

- 30 wherein the chelator is capable of conjugating to the diagnostic metal.

- (2) A diagnostic agent according to embodiment 1, wherein the targeting moiety is a matrix metalloproteinase inhibitor having
35 an inhibitory constant K_i of <1000 nM.

(3) A diagnostic agent according to any of embodiments 1-2, wherein the targeting moiety is a matrix metalloproteinase inhibitor having an inhibitory constant K_i of <100 nM.

5 (4) A diagnostic agent according to any of embodiments 1-3, comprising 1-5 targeting moieties.

(5) A diagnostic agent according to any of embodiments 1-4, comprising one targeting moiety.

10


(6) A diagnostic agent any of embodiments 1-5, wherein the targeting moiety is an inhibitor of one or more matrix metalloproteinases selected from the group consisting of MMP-1, MMP-2, MMP-3, MMP-9, and MMP-14.

15

(7) A diagnostic agent of any one of embodiments 1-6, wherein the targeting moiety is an inhibitor of one or more matrix metalloproteinases selected from the group consisting of MMP-2, MMP-9, and MMP-14.

20

(8) A diagnostic agent according to any one of embodiments 1-7 wherein the targeting moiety is a matrix metalloproteinase inhibitor of the formulae (Ia) or (Ib):




R is independently OH or $-\text{CH}_2\text{SH}$;

30 R^1 is independently selected at each occurrence from the group:
H, OH, C₁₋₃ alkyl, C₂₋₃ alkenyl, C₂₋₃ alkynyl, and heterocycle-S-CH₂-;

R^2 is independently C₁₋₂₀ alkyl;

X is independently C=O or SO₂, provided when X is C=O, R³ is



, and when X is SO₂, R³ is independently selected from the group: aryl substituted with 0-2 R⁶, and
5 heterocycle substituted with 0-2 R⁶;

R⁴ is independently selected at each occurrence from the group:
C₁₋₆ alkyl, phenyl, and benzyl;

10 R⁵ is independently at each occurrence from the group: NH(C₁₋₆ alkyl), NH-phenyl, and NH-heterocycle; wherein said alkyl, phenyl and heterocycle groups are optionally substituted with a bond to the linking group or a bond to the chelator;

15 R⁶ is independently aryloxy substituted with 0-3 R⁷;

R⁷ is independently halogen or methoxy;

or alternatively,

20

R¹ and R⁴ may be taken together to form a bridging group of the formula -(CH₂)₃-O-phenyl-CH₂-, optionally substituted with a bond to the linking group or a bond to the chelator;

25

or alternatively,

R¹ and R² may be taken together to form a bridging group of the formula -(CH₂)₃-NH-, optionally substituted with a bond to
30 the linking group or a bond to the chelator; or

R¹ and R² taken together with the nitrogen and carbon atom through which they are attached form a C₅₋₇ atom saturated

ring system substituted with one or more substituents selected from the group consisting of: a bond to Ln, a bond to Ch, and $-C(=O)-NR^{29}R^{30}$;

- 5 R^8 is independently at each occurrence OH or phenyl, optionally substituted with a bond to the linking group or a bond to the chelator, provided that when R^8 is phenyl, R^{10} is $-C(=O)-CR^{12}-NH-CH(CH_3)-COOH$;
- 10 R^9 and $R^{9'}$ are independently H, C₁₋₆ alkyl optionally substituted with a bond to the linking group or a bond to the chelator, or are taken together with the carbon atom to which R^9 and $R^{9'}$ are attached to form a 5-7 atom saturated, partially unsaturated or aromatic ring system containing 0-3 heteroatoms selected from O, N, SO₂ and S, said ring system substituted with R^6 and optionally substituted with a bond to the linking group or a bond to the chelator;
- 15
- R^{10} and R^{11} are independently H, or C₁₋₆ alkyl optionally substituted with a bond to the linking group or a bond to the chelator, or are taken together with the nitrogen atom to which they are attached to form a 5-7 atom saturated, partially unsaturated or aromatic ring system containing 0-3 heteroatoms selected from O, N, SO₂ and S, said ring system optionally substituted with 0-3 R^{27} , a bond to the linking group or a bond to the chelator;
- 20
- 25

or alternatively,

- 30 R^9 and R^{10} are taken together with the carbon atom to which they are attached to form a 5-7 atom saturated, partially unsaturated or aromatic ring system containing 0-3 heteroatoms selected from O, N, SO₂ and S, said ring system optionally substituted with a bond to the linking group or a bond to the chelator; and
- 35

R¹² is independently C₁-20 alkyl;

R²⁷ is =O, C₁-4 alkyl, or phenyl substituted with R²⁸;


R²⁸ is a phenoxy group substituted with 0-2 OCH₃ groups;

5 R²⁹ and R³⁰ taken together with the nitrogen atom through which they are attached form a C₅-7 atom saturated ring system substituted with R³¹; and

R³¹ is a benzyloxy group substituted with C₁-4 alkyl.

10

(9). A diagnostic agent according to any one of embodiments 1-8 wherein the targeting moiety is a matrix metalloproteinase inhibitor of the formulae (Ia) or (Ib):



R is OH;

20 R¹ is independently selected at each occurrence from the group:
H, OH, C₁-3 alkyl, C₂-3 alkenyl, C₂-3 alkynyl, and
heterocycle-S-CH₂-;

R² is independently C₁-6 alkyl;

25 X is C=O;

R⁴ is independently selected at each occurrence from the group:
C₁-6 alkyl, phenyl, and benzyl;

30 R⁵ is independently at each occurrence from the group: NH(C₁-6 alkyl), NH-phenyl, and NH-heterocycle; wherein said alkyl,

phenyl and heterocycle groups are optionally substituted with a bond to the linking group or a bond to the chelator;

R⁶ is independently aryloxy substituted with 0-3 R⁷;

5

R⁷ is independently halogen or methoxy;

or alternatively,

10 R¹ and R⁴ may be taken together to form a bridging group of the formula $-(CH_2)_3-O-phenyl-CH_2-$, optionally substituted with a bond to the linking group or a bond to the chelator;

or alternatively,

15

R¹ and R² may be taken together to form a bridging group of the formula $-(CH_2)_3-NH-$, optionally substituted with a bond to the linking group or a bond to the chelator; or

20 R¹ and R² taken together with the nitrogen and carbon atom through which they are attached form a C₅₋₇ atom saturated ring system substituted with one or more substituents selected from the group consisting of: a bond to Ln, a bond to Ch, and $-C(=O)-NR^{29}R^{30}$;

25

R⁸ is OH;

R⁹ and R^{9'} are independently H, C₁₋₆ alkyl optionally substituted with a bond to the linking group or a bond to the chelator, or are taken together with the carbon atom to which R⁹ and R^{9'} are attached to form a 5-7 atom saturated, partially unsaturated or aromatic ring system containing 0-1 heteroatoms selected from O, N, , said ring system

30

optionally substituted with a bond to the linking group or a bond to the chelator;

R¹⁰ and R¹¹ are independently H, or C₁₋₆ alkyl optionally substituted with a bond to the linking group or a bond to the chelator, or are taken together with the nitrogen atom to which they are attached to form a 5-7 atom saturated, partially unsaturated or aromatic ring system containing 0-1 heteroatoms selected from O, N, , said ring system optionally substituted with 0-3 R²⁷, a bond to the linking group or a bond to the chelator;

or alternatively,

R⁹ and R¹⁰ are taken together with the carbon atom to which they are attached to form a 5-7 atom saturated, partially unsaturated or aromatic ring system containing 0-1 heteroatoms selected from O, N, , said ring system optionally substituted with a bond to the linking group or a bond to the chelator; and

R¹² is independently C₁₋₆ alkyl;

R²⁷ is =O, C₁₋₄ alkyl, or phenyl substituted with R²⁸;

R²⁸ is a phenoxy group substituted with 0-2 OCH₃ groups;

R²⁹ and R³⁰ taken together with the nitrogen atom through which they are attached form a C₅₋₇ atom saturated ring system substituted with R³¹; and

R³¹ is a benzyloxy group substituted with C₁₋₄ alkyl.


(10). A diagnostic agent according to any one of embodiments 1-9 wherein the targeting moiety is a matrix metalloproteinase inhibitor of the formulae (Ia) or (Ib):

wherein:

R is -OH;

R² is C₁₋₆ alkyl;

X is C=O;



R¹ and R⁴ are taken together to form a bridging group of formula
5 -(CH₂)₃-O-phenyl-CH₂-;

R⁵ is NH(C₁₋₆alkyl), substituted with a bond to the linking
group or a bond to the chelator.

(11) A diagnostic agent according to any one of embodiments 1-
10 10, wherein:

R is -OH;

R⁹ is C₁ alkyl substituted with a bond to Ln;

R¹⁰ and R¹¹ taken together with the nitrogen atom to which they
are attached form a 5 atom saturated ring system, said right
15 system is substituted with 0-3 R²⁷;

R²⁷ is =O, C₁₋₄ alkyl, or phenyl substituted with R²⁸; and

R²⁸ is a phenoxy group substituted with 0-2 OCH₃ groups.

(12) A diagnostic agent according to any one of embodiments 1-11
20 wherein the


R is -OH;

R¹ and R² taken together with the nitrogen and carbon atom
through which they are attached form a C₅₋₇ atom saturated ring
system substituted with one or more substituents selected from
25 the group consisting of: a bond to Ln, a bond to Ch, and -C(=O)-
NR²⁹R³⁰;

R²⁹ and R³⁰ taken together with the nitrogen atom through which
they are attached form a C₅₋₇ atom saturated ring system
substituted with R³¹; and

30 R³¹ is a benzyloxy group substituted with C₁₋₄ alkyl.

(13) A diagnostic agent according to any one of embodiments 1-12 wherein the linking group is of the formula:



5

W^1 and W^2 are independently selected at each occurrence from the group: O, S, NH, NHC(=O), C(=O)NH, $NR^{15}C(=O)$, $C(=O)NR^{15}$, C(=O), C(=O)O, OC(=O), NHC(=S)NH, NHC(=O)NH, SO₂, SO₂NH, - (OCH₂CH₂)₇₆₋₈₄, (OCH₂CH₂)_s, (CH₂CH₂O)_{s'}, (OCH₂CH₂CH₂)_{s''},
 10 (CH₂CH₂CH₂O)_t, and (aa)_{t'};

aa is independently at each occurrence an amino acid;

Z is selected from the group: aryl substituted with 0-3 R^{16} ,
 15 C₃-10 cycloalkyl substituted with 0-3 R^{16} , and a 5-10 membered heterocyclic ring system containing 1-4 heteroatoms independently selected from N, S, and O and substituted with 0-3 R^{16} ;

20 R^{13} , R^{13a} , R^{14} , R^{14a} , and R^{15} are independently selected at each occurrence from the group: H, =O, COOH, SO₃H, PO₃H, C₁-C₅ alkyl substituted with 0-3 R^{16} , aryl substituted with 0-3 R^{16} , benzyl substituted with 0-3 R^{16} , and C₁-C₅ alkoxy substituted with 0-3 R^{16} , NHC(=O) R^{17} , C(=O)NHR¹⁷,
 25 NHC(=O)NHR¹⁷, NHR¹⁷, R^{17} , and a bond to the chelator;

R^{16} is independently selected at each occurrence from the group: a bond to the chelator, COOR¹⁷, C(=O)NHR¹⁷, NHC(=O) R^{17} , OH, NHR¹⁷, SO₃H, PO₃H, -OPO₃H₂, -OSO₃H, aryl substituted with
 30 0-3 R^{17} , C₁-5 alkyl substituted with 0-1 R^{18} , C₁-5 alkoxy substituted with 0-1 R^{18} , and a 5-10 membered heterocyclic

ring system containing 1-4 heteroatoms independently
selected from N, S, and O and substituted with 0-3 R¹⁷;

R¹⁷ is independently selected at each occurrence from the group:

5 H, alkyl substituted with 0-1 R¹⁸, aryl substituted with
0-1 R¹⁸, a 5-10 membered heterocyclic ring system
containing 1-4 heteroatoms independently selected from N,
S, and O and substituted with 0-1 R¹⁸, C₃-10 cycloalkyl
substituted with 0-1 R¹⁸, polyalkylene glycol substituted
10 with 0-1 R¹⁸, carbohydrate substituted with 0-1 R¹⁸,
cyclodextrin substituted with 0-1 R¹⁸, amino acid
substituted with 0-1 R¹⁸, polycarboxyalkyl substituted with
0-1 R¹⁸, polyazaalkyl substituted with 0-1 R¹⁸, peptide
substituted with 0-1 R¹⁸, wherein the peptide is comprised
15 of 2-10 amino acids, 3,6-O-disulfo-B-D-galactopyranosyl,
bis(phosphonomethyl)glycine, and a bond to the chelator;

R¹⁸ is a bond to the chelator;

20 k is selected from 0, 1, and 2;
h is selected from 0, 1, and 2;
h' is selected from 0, 1, and 2;
g is selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;
g' is selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;
25 s is selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;
s' is selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;
s" is selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;
t is selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;
t' is selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;
30 x is selected from 0, 1, 2, 3, 4, and 5; and
x' is selected from 0, 1, 2, 3, 4, and 5.

(14) A diagnostic agent according to any one of embodiments 1-13 wherein

W^1 and W^2 are independently selected at each occurrence from the group: O, NH, $NHC(=O)$, $C(=O)NH$, $NR^{15}C(=O)$, $C(=O)NR^{15}$,
 5 $C(=O)$, $C(=O)O$, $OC(=O)$, $NHC(=S)NH$, $NHC(=O)NH$, SO_2 , -
 $(CH_2CH_2O)_{76-84-}$, $(OCH_2CH_2)_s$, $(CH_2CH_2O)_s'$, $(OCH_2CH_2CH_2)_s''$,
 $(CH_2CH_2CH_2O)_t$, and $(aa)_t$;

aa is independently at each occurrence an amino acid;

10

Z is selected from the group: aryl substituted with 0-1 R^{16} ,
 C_3 -10 cycloalkyl substituted with 0-1 R^{16} , and a 5-10
 membered heterocyclic ring system containing 1-4
 heteroatoms independently selected from N, S, and O and
 15 substituted with 0-1 R^{16} ;

R^{13} , R^{13a} , R^{14} , R^{14a} , and R^{15} are independently selected at each
 occurrence from the group: H, =O, COOH, SO_3H , C_1 - C_5 alkyl
 substituted with 0-1 R^{16} , aryl substituted with 0-1 R^{16} ,
 20 benzyl substituted with 0-1 R^{16} , and C_1 - C_5 alkoxy
 substituted with 0-1 R^{16} , $NHC(=O)R^{17}$, $C(=O)NHR^{17}$,
 $NHC(=O)NHR^{17}$, NHR^{17} , R^{17} , and a bond to the chelator;

k is 0 or 1;

25 s is selected from 0, 1, 2, 3, 4, and 5;

s' is selected from 0, 1, 2, 3, 4, and 5;

s'' is selected from 0, 1, 2, 3, 4, and 5; and

t is selected from 0, 1, 2, 3, 4, and 5.

30 (15) A diagnostic agent according to embodiment 13 wherein
 wherein:

W^1 is $C(=O)NR^{15}$;

h is 1;

g is 3;

R¹³ and R¹⁴ are independently H;

x is 1;

k is 0;

g' is 0;

5 h' is 1;

W² is NH; and

x' is 1.

(16) A diagnostic agent according to embodiment 13 wherein

10 x is 0;

k is 1;

Z is aryl substituted with 0-3 R¹⁶;

g' is 1;

W² is NH;

15 R^{13a} and R^{14a} are independently H;

h' is 1; and

x' is 1.

(17) A diagnostic agent according to embodiment 13 wherein

20 W¹ is C(=O)NR¹⁵;

h is 1;

g is 2;

R¹³ and R¹⁴ are independently H;

x is 1;

25 k is 0;

g' is 1;

R^{13a} and R^{14a} are independently H; or C1-5 alkyl substituted
with 0-3 R¹⁶;

R¹⁶ is SO₃H;

30 W² is NHC(=O) or NH;

h' is 1; and

x' is 2.

(18). A diagnostic agent according to embodiment 13 wherein

- W¹ is C(=O)NH;
h is 1;
g is 3;
R¹³ and R¹⁴ are independently H;
5 k is 0;
g' is 0;
x is 1;
W² is -NH(C=O)- or -(OCH₂CH₂)₇₆₋₈₄-;
h' is 2; and
10 x' is 1.
- (19) A diagnostic agent according to embodiment 13 wherein
x is 0;
k is 0;
15 g' is 3;
h' is 1;
W² is NH; and
x' is 1.
- 20 (20) A diagnostic agent according to embodiment 13 wherein
x is 0;
Z is aryl substituted with 0-3 R¹⁶;
k is 1;
g' is 1;
25 R^{13a}R^{14a} are independently H;
W² is NHC(=O) or -(OCH₂CH₂)₇₆₋₈₄-; and
x' is 1.
- (21) A diagnostic agent according to embodiment 13 wherein
30 W¹ is C=O;
g is 2;
R¹³ and R¹⁴ are independently H;
k is 0;
g' is 0;


h' is 1;



w^2 is NH; and

x' is 1.

5 (22) A compound according to embodiment 1 wherein the linking group is absent.

(23) A diagnostic agent according to any one of embodiments 1-22
 10 wherein the chelator is a metal bonding unit having a formula selected from the group:





A¹, A², A³, A⁴, A⁵, A⁶, A⁷, and A⁸ are independently selected at
 5 each occurrence from the group: N, NR²⁶, NR¹⁹, NR¹⁹R²⁰, S,
 SH, -S(Pg), O, OH, PR¹⁹, PR¹⁹R²⁰, -O-P(O)(R²¹)-O-,
 P(O)R²¹R²², a bond to the targeting moiety and a bond to
 the linking group;

10 Pg is a thiol protecting group;

E¹, E², E³, E⁴, E⁵, E⁶, E⁷, and E⁸ are independently a bond, CH,
 or a spacer group independently selected at each occurrence
 from the group: C₁-C₁₆ alkyl substituted with 0-3 R²³,
 15 aryl substituted with 0-3 R²³, C₃-10 cycloalkyl substituted
 with 0-3 R²³, heterocyclo-C₁-10 alkyl substituted with 0-3
 R²³, wherein the heterocyclo group is a 5-10 membered
 heterocyclic ring system containing 1-4 heteroatoms
 independently selected from N, S, and O, C₆-10 aryl-C₁-10
 20 alkyl substituted with 0-3 R²³, C₁-10 alkyl-C₆-10 aryl-
 substituted with 0-3 R²³, and a 5-10 membered heterocyclic

ring system containing 1-4 heteroatoms independently selected from N, S, and O and substituted with 0-3 R²³;

R¹⁹ and R²⁰ are each independently selected from the group: a
5 bond to the linking group, a bond to the targeting moiety, hydrogen, C₁-C₁₀ alkyl substituted with 0-3 R²³, aryl substituted with 0-3 R²³, C₁-10 cycloalkyl substituted with 0-3 R²³, heterocyclo-C₁-10 alkyl substituted with 0-3 R²³, wherein the heterocyclo group is a 5-10 membered
10 heterocyclic ring system containing 1-4 heteroatoms independently selected from N, S, and O, C₆-10 aryl-C₁-10 alkyl substituted with 0-3 R²³, C₁-10 alkyl-C₆-10 aryl-substituted with 0-3 R²³, a 5-10 membered heterocyclic ring system containing 1-4 heteroatoms independently selected
15 from N, S, and O and substituted with 0-3 R²³, and an electron, provided that when one of R¹⁹ or R²⁰ is an electron, then the other is also an electron;

R²¹ and R²² are each independently selected from the group: a
20 bond to the linking group, a bond to the targeting moiety, -OH, C₁-C₁₀ alkyl substituted with 0-3 R²³, C₁-C₁₀ alkyl substituted with 0-3 R²³, aryl substituted with 0-3 R²³, C₃-10 cycloalkyl substituted with 0-3 R²³, heterocyclo-C₁-10 alkyl substituted with 0-3 R²³, wherein
25 the heterocyclo group is a 5-10 membered heterocyclic ring system containing 1-4 heteroatoms independently selected from N, S, and O, C₆-10 aryl-C₁-10 alkyl substituted with 0-3 R²³, C₁-10 alkyl-C₆-10 aryl-substituted with 0-3 R²³, and a 5-10 membered heterocyclic ring system containing 1-4
30 heteroatoms independently selected from N, S, and O and substituted with 0-3 R²³;

R^{23} is independently selected at each occurrence from the group:
 a bond to the linking group, a bond to the targeting
 moiety, =O, F, Cl, Br, I, $-CF_3$, $-CN$, $-CO_2R^{24}$, $-C(=O)R^{24}$,
 $-C(=O)N(R^{24})_2$, $-CHO$, $-CH_2OR^{24}$, $-OC(=O)R^{24}$, $-OC(=O)OR^{24a}$,
 5 $-OR^{24}$, $-OC(=O)N(R^{24})_2$, $-NR^{25}C(=O)R^{24}$, $-NR^{25}C(=O)OR^{24a}$,
 $-NR^{25}C(=O)N(R^{24})_2$, $-NR^{25}SO_2N(R^{24})_2$, $-NR^{25}SO_2R^{24a}$, $-SO_3H$,
 $-SO_2R^{24a}$, $-SR^{24}$, $-S(=O)R^{24a}$, $-SO_2N(R^{24})_2$, $-N(R^{24})_2$,
 $-NHC(=S)NHR^{24}$, $=NOR^{24}$, NO_2 , $-C(=O)NHOR^{24}$, $-C(=O)NHN(R^{24})_2R^{24a}$,
 $-OCH_2CO_2H$, 2-(1-morpholino)ethoxy, C₁-C₅ alkyl, C₂-C₄
 10 alkenyl, C₃-C₆ cycloalkyl, C₃-C₆ cycloalkylmethyl, C₂-C₆
 alkoxyalkyl, aryl substituted with 0-2 R^{24} , and a 5-10
 membered heterocyclic ring system containing 1-4
 heteroatoms independently selected from N, S, and O; and
 wherein at least one of A^1 , A^2 , A^3 , A^4 , A^5 , A^6 , A^7 , A^8 or R^{23} is
 15 a bond to the linking group or targeting moiety;
 R^{24} , R^{24a} , and R^{25} are independently selected at each occurrence
 from the group: a bond to the linking group, a bond to the
 targeting moiety, H, C₁-C₆ alkyl, phenyl, benzyl, C₁-C₆ alkoxy,
 halide, nitro, cyano, and trifluoromethyl; and
 20 R^{26} is a co-ordinate bond to a metal or a hydrazine protecting
 group; or a pharmaceutically acceptable salt thereof.

(24) A diagnostic agent according to any one of embodiments 1-23
 25 wherein:

A^1 , A^2 , A^3 , A^4 , A^5 , A^6 , A^7 , and A^8 are independently selected at
 each occurrence from the group: NR^{19} , $NR^{19}R^{20}$, S, SH, OH,
 a bond to the targeting moiety and a bond to the linking
 group;

30 E^1 , E^2 , E^3 , E^4 , E^5 , E^6 , E^7 , and E^8 are independently a bond,
 CH, or a spacer group independently selected at each
 occurrence from the group: C₁-C₁₀ alkyl substituted with

0-3 R²³, aryl substituted with 0-3 R²³, C₃-10 cycloalkyl substituted with 0-3 R²³, and a 5-10 membered heterocyclic ring system containing 1-4 heteroatoms independently selected from N, S, and O and substituted with 0-3 R²³;

5


wherein at least one of A¹, A², A³, A⁴, A⁵, A⁶, A⁷, A⁸ and R²³ is a bond to the linking group or the targeting moiety;

R¹⁹, and R²⁰ are each independently selected from the group: a bond to the targeting moiety, a bond to the linking group, hydrogen, C₁-C₁₀ alkyl substituted with 0-3 R²³, aryl substituted with 0-3 R²³, a 5-10 membered heterocyclic ring system containing 1-4 heteroatoms independently selected from N, S, and O and substituted with 0-3 R²³, and an electron, provided that when one of R¹⁹ or R²⁰ is an electron, then the other is also an electron;

R²³ is independently selected at each occurrence from the group: a bond to the targeting moiety, a bond to the linking group, =O, F, Cl, Br, I, -CF₃, -CN, -CO₂R²⁴, -C(=O)R²⁴, -C(=O)N(R²⁴)₂, -CH₂OR²⁴, -OC(=O)R²⁴, -OC(=O)OR^{24a}, -OR²⁴, -OC(=O)N(R²⁴)₂, -NR²⁵C(=O)R²⁴, -NR²⁵C(=O)OR^{24a}, -NR²⁵C(=O)N(R²⁴)₂, -NR²⁵SO₂N(R²⁴)₂, -NR²⁵SO₂R^{24a}, -SO₃H, -SO₂R^{24a}, -S(=O)R^{24a}, -SO₂N(R²⁴)₂, -N(R²⁴)₂, -NHC(=S)NHR²⁴, =NOR¹⁸, -C(=O)NHN(R¹⁸)R^{18a}, -OCH₂CO₂H, and 2-(1-morpholino)ethoxy; and

R²⁴, R^{24a}, and R²⁵ are independently selected at each occurrence from the group: a bond to the linking group, H, and C₁-C₆ alkyl.

(25) A diagnostic agent according to any one of embodiments 1-24 wherein the chelator is of the formula:



5

A¹ is a bond to the linking group;

A², A⁴, and A⁶ are each N;

10

A³, A⁵, A⁷ and A⁸ are each OH;

E¹, E², and E⁴ are C2 alkyl;

E³, E⁵, E⁷, and E⁸ are C2 alkyl substituted with 0-1 R²³;


15

R²³ is =O.

(26) A diagnostic agent according to any one of embodiments 1-25 wherein the chelator is of the formula:

20

Ch is



wherein:

A⁵ is a bond to Ln;

25

A¹, A³, A⁷ and A⁸ are each OH;


A², A⁴ and A⁶ are each NH;

E¹, E³, E⁵, E⁷, and E⁸ are C2 alkyl substituted with 0-1 R²³;

E², and E⁴, are C₂ alkyl;

R²³ is =O.

(27) A diagnostic agent according to any one of embodiments 1-26
5 wherein the chelator is of the formula:



10 A¹, A², A³ and A⁴ are each N;

A⁵, A⁶ and A⁸ are each OH;

A⁷ is a bond to L_n;

E¹, E², E³, E⁴ are each independently C₂ alkyl; and

15 E⁵, E⁶, E⁷, E⁸ are each independently C₂ alkyl substituted with
0-1 R²³;

R²³ is =O.

20 (28) A diagnostic agent according to any one of embodiments 1-27

wherein the chelator is of the formula:
$$\begin{array}{c} \text{E}^1\text{---A}^2 \\ \diagup \\ \text{A}^1 \end{array} ;$$

A¹ is NR²⁶;

R^{26} is a co-ordinate bond to a metal or a hydrazine protecting group;;

E^1 is a bond;

5

A^2 is NHR^{19} ;

R^{19} is a heterocycle substituted with R^{23} , the heterocycle being selected from pyridine and pyrimidine;


10

R^{23} is selected from a bond to the linking group, $C(=O)NHR^{24}$ and $C(=O)R^{24}$; and

R^{24} is a bond to the linking group.

15

(29) A diagnostic agent according to any one of embodiments 1-28 wherein the chelator is of the formula:



wherein:

20 A^1 and A^5 are each $-S(Pg)$;

Pg is a thiol protecting group;

E^1 and E^4 are C_2 alkyl substituted with 0-1 R^{23} ;

R^{23} is $=O$;

A^2 and A^4 are each $-NH$;


25 E^2 is CH_2 ;

E^3 is C_{1-3} alkyl substituted with 0-1 R^{23} ;

A^3 is a bond to Ln .

(30) A diagnostic agent according to any one of embodiments 1-29 wherein the chelator is of the formula:

30



wherein:

A¹ is a bond to Ln;

E¹ is C₁ alkyl substituted by R²³;

5 A² is NH;

E² is C₂ alkyl substituted with 0-1R²³;

A³ is -O-P(O)(R²¹)-O;

E³ is C₁ alkyl;

A⁴ and A⁵ are each -O-;

10 E⁴ and E⁶ are each independently C₁₋₁₆ alkyl substituted with 0-1R²³;

E⁵ is C₁ alkyl;


R²¹ is -OH; and

R²³ is =O.


15

(31) A diagnostic agent according to embodiment 1 having the formula:

20



wherein, Q is a compound of Formulae (Ia) or (Ib):




25

R is independently OH or -CH₂SH;

R¹ is independently selected at each occurrence from the group:
H, OH, C₁-3 alkyl, C₂-3 alkenyl, C₂-3 alkynyl, and
heterocycle-S-CH₂-;

5 R² is independently C₁-20 alkyl;

X is independently C=O or SO₂, provided when X is C=O, R³ is



, and when X is SO₂, R³ is independently selected
from the group: aryl substituted with 0-2 R⁶, and
10 heterocycle substituted with 0-2 R⁶;

R⁴ is independently selected at each occurrence from the group:
C₁-6 alkyl, phenyl, and benzyl;

15 R⁵ is independently at each occurrence from the group: NH(C₁-6
alkyl), NH-phenyl, and NH-heterocycle; wherein said alkyl,
phenyl and heterocycle groups are optionally substituted
with a bond to L_n;

20 R⁶ is independently aryloxy substituted with 0-3 R⁷;

R⁷ is independently halogen or methoxy;

or alternatively,

25

R¹ and R⁴ may be taken together to form a bridging group of the
formula -(CH₂)₃-O-phenyl-CH₂-, optionally substituted with a
bond to L_n;

30 or alternatively,

R¹ and R² may be taken together to form a bridging group of the formula -(CH₂)₃-NH-, optionally substituted with a bond to L_n; or

5 R¹ and R² taken together with the nitrogen and carbon atom through which they are attached form a C₅₋₇ atom saturated ring system substituted with one or more substituents selected from the group consisting of: a bond to L_n, a bond to Ch, and -C(=O)-NR²⁹R³⁰;

10

R⁸ is independently at each occurrence OH or phenyl, optionally substituted with a bond to L_n, provided that when R⁸ is phenyl, R¹⁰ is -C(=O)-CR¹²-NH-CH(CH₃)-COOH;

15 R⁹ and R^{9'} are independently H, C₁₋₆ alkyl optionally substituted with a bond to L_n, or are taken together with the carbon atom to which they are attached to form a 5-7 atom saturated, partially unsaturated or aromatic ring system containing 0-3 heteroatoms selected from O, N, SO₂ and S, said ring system substituted with R⁶ and optionally substituted with a bond to L_n;

20

R¹⁰ and R¹¹ are independently H, or C₁₋₆ alkyl optionally substituted with a bond to L_n, or are taken together with the nitrogen atom to which they are attached to form a 5-7 atom saturated, partially unsaturated or aromatic ring system containing 0-3 heteroatoms selected from O, N, SO₂ and S, said ring system optionally substituted with 0-3 R²⁷ or a bond to L_n;

25

30

or alternatively,

R⁹ and R¹⁰ are taken together with the carbon atom to which they are attached to form a 5-7 atom saturated, partially

unsaturated or aromatic ring system containing 0-3 heteroatoms selected from O, N, SO₂ and S, said ring system optionally substituted with a bond to L_n;

5 R¹² is independently C₁₋₂₀ alkyl;

d is selected from 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;

L_n is a linking group having the formula:

10

$$((W^1)_h - (CR^{13}R^{14})_g)_x - (Z)_k - ((CR^{13a}R^{14a})_{g'} - (W^2)_{h'})_{x'};$$

W¹ and W² are independently selected at each occurrence from the group: O, S, NH, NHC(=O), C(=O)NH, NR¹⁵C(=O), C(=O)NR¹⁵,
 15 C(=O), C(=O)O, OC(=O), NHC(=S)NH, NHC(=O)NH, SO₂, SO₂NH, -
 (OCH₂CH₂)₇₆₋₈₄, (OCH₂CH₂)_s, (CH₂CH₂O)_{s'}, (OCH₂CH₂CH₂)_{s''},
 (CH₂CH₂CH₂O)_t, and (aa)_{t'};

aa is independently at each occurrence an amino acid;
 20

Z is selected from the group: aryl substituted with 0-3 R¹⁶,
 C₃₋₁₀ cycloalkyl substituted with 0-3 R¹⁶, and a 5-10
 membered heterocyclic ring system containing 1-4
 heteroatoms independently selected from N, S, and O and
 25 substituted with 0-3 R¹⁶;

R¹³, R^{13a}, R¹⁴, R^{14a}, and R¹⁵ are independently selected at each occurrence from the group: H, =O, COOH, SO₃H, PO₃H, C₁₋₅
 alkyl substituted with 0-3 R¹⁶, aryl substituted with 0-3
 30 R¹⁶, benzyl substituted with 0-3 R¹⁶, and C₁₋₅ alkoxy
 substituted with 0-3 R¹⁶, NHC(=O)R¹⁷, C(=O)NHR¹⁷,
 NHC(=O)NHR¹⁷, NHR¹⁷, R¹⁷, and a bond to C_H;

R¹⁶ is independently selected at each occurrence from the group:
a bond to C_H, COOR¹⁷, C(=O)NHR¹⁷, NHC(=O)R¹⁷, OH, NHR¹⁷,
SO₃H, PO₃H, -OPO₃H₂, -OSO₃H, aryl substituted with 0-3 R¹⁷,
C₁₋₅ alkyl substituted with 0-1 R¹⁸, C₁₋₅ alkoxy
5 substituted with 0-1 R¹⁸, and a 5-10 membered heterocyclic
ring system containing 1-4 heteroatoms independently
selected from N, S, and O and substituted with 0-3 R¹⁷;

R¹⁷ is independently selected at each occurrence from the group:
10 H, alkyl substituted with 0-1 R¹⁸, aryl substituted with
0-1 R¹⁸, a 5-10 membered heterocyclic ring system
containing 1-4 heteroatoms independently selected from N,
S, and O and substituted with 0-1 R¹⁸, C₃₋₁₀ cycloalkyl
substituted with 0-1 R¹⁸, polyalkylene glycol substituted
15 with 0-1 R¹⁸, carbohydrate substituted with 0-1 R¹⁸,
cyclodextrin substituted with 0-1 R¹⁸, amino acid
substituted with 0-1 R¹⁸, polycarboxyalkyl substituted with
0-1 R¹⁸, polyazaalkyl substituted with 0-1 R¹⁸, peptide
substituted with 0-1 R¹⁸, wherein the peptide is comprised
20 of 2-10 amino acids, 3,6-O-disulfo-B-D-galactopyranosyl,
bis(phosphonomethyl)glycine, and a bond to C_H;

R¹⁸ is a bond to C_H;


25 k is selected from 0, 1, and 2;
h is selected from 0, 1, and 2;
h' is selected from 0, 1, and 2;
g is selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;
g' is selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;
30 s is selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;
s' is selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;
s" is selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;
t is selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;

t' is selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;



x is selected from 0, 1, 2, 3, 4, and 5;

x' is selected from 0, 1, 2, 3, 4, and 5;

- 5 Ch is a metal bonding unit having a formula selected from the group:



10



A¹, A², A³, A⁴, A⁵, A⁶, A⁷, and A⁸ are independently selected at each occurrence from the group: N, NR²⁶, NR¹⁹, NR¹⁹R²⁰, S, SH, -S(Pg), O, OH, PR¹⁹, PR¹⁹R²⁰, -O-P(O)(R²¹)-O-, P(O)R²¹R²²; a bond to the targeting moiety and a bond to
5 the linking group;

Pg is a thiol protecting group;

E¹, E², E³, E⁴, E⁵, E⁶, E⁷, and E⁸ are independently a bond, CH,
10 or a spacer group independently selected at each occurrence from the group: C₁-C₁₆ alkyl substituted with 0-3 R²³, aryl substituted with 0-3 R²³, C₃-₁₀ cycloalkyl substituted with 0-3 R²³, heterocyclo-C₁-₁₀ alkyl substituted with 0-3 R²³, wherein the heterocyclo group is a 5-10 membered
15 heterocyclic ring system containing 1-4 heteroatoms independently selected from N, S, and O, C₆-₁₀ aryl-C₁-₁₀ alkyl substituted with 0-3 R²³, C₁-₁₀ alkyl-C₆-₁₀ aryl-substituted with 0-3 R²³, and a 5-10 membered heterocyclic ring system containing 1-4 heteroatoms independently
20 selected from N, S, and O and substituted with 0-3 R²³;

R¹⁹ and R²⁰ are each independently selected from the group: a bond to the linking group, a bond to the targeting moiety, hydrogen, C₁-C₁₀ alkyl substituted with 0-3 R²³, aryl
25 substituted with 0-3 R²³, C₁-₁₀ cycloalkyl substituted with 0-3 R²³, heterocyclo-C₁-₁₀ alkyl substituted with 0-3 R²³, wherein the heterocyclo group is a 5-10 membered heterocyclic ring system containing 1-4 heteroatoms independently selected from N, S, and O, C₆-₁₀ aryl-C₁-₁₀
30 alkyl substituted with 0-3 R²³, C₁-₁₀ alkyl-C₆-₁₀ aryl-substituted with 0-3 R²³, a 5-10 membered heterocyclic ring system containing 1-4 heteroatoms independently selected

from N, S, and O and substituted with 0-3 R²³, and an electron, provided that when one of R¹⁹ or R²⁰ is an electron, then the other is also an electron;

5 R²¹ and R²² are each independently selected from the group: a bond to the linking group, a bond to the targeting moiety, -OH, C₁-C₁₀ alkyl substituted with 0-3 R²³, C₁-C₁₀ alkyl substituted with 0-3 R²³, aryl substituted with 0-3 R²³, C₃-10 cycloalkyl substituted with 0-3 R²³,
 10 heterocyclo-C₁-10 alkyl substituted with 0-3 R²³, wherein the heterocyclo group is a 5-10 membered heterocyclic ring system containing 1-4 heteroatoms independently selected from N, S, and O, C₆-10 aryl-C₁-10 alkyl substituted with 0-3 R²³, C₁-10 alkyl-C₆-10 aryl substituted with 0-3 R²³,
 15 and a 5-10 membered heterocyclic ring system containing 1-4 heteroatoms independently selected from N, S, and O and substituted with 0-3 R²³;

R²³ is independently selected at each occurrence from the group:
 20 a bond to the linking group, a bond to the targeting moiety, =O, F, Cl, Br, I, -CF₃, -CN, -CO₂R²⁴, -C(=O)R²⁴, -C(=O)N(R²⁴)₂, -CHO, -CH₂OR²⁴, -OC(=O)R²⁴, -OC(=O)OR^{24a}, -OR²⁴, -OC(=O)N(R²⁴)₂, -NR²⁵C(=O)R²⁴, -NR²⁵C(=O)OR^{24a}, -NR²⁵C(=O)N(R²⁴)₂, -NR²⁵SO₂N(R²⁴)₂, -NR²⁵SO₂R^{24a}, -SO₃H,
 25 -SO₂R^{24a}, -SR²⁴, -S(=O)R^{24a}, -SO₂N(R²⁴)₂, -N(R²⁴)₂, -NHC(=S)NHR²⁴, =NOR²⁴, NO₂, -C(=O)NHR²⁴, -C(=O)NHN(R²⁴)₂, -OCH₂CO₂H, 2-(1-morpholino)ethoxy, C₁-C₅ alkyl, C₂-C₄ alkenyl, C₃-C₆ cycloalkyl, C₃-C₆ cycloalkylmethyl, C₂-C₆ alkoxyalkyl, aryl substituted with 0-2 R²⁴, and a 5-10
 30 membered heterocyclic ring system containing 1-4 heteroatoms independently selected from N, S, and O; and

wherein at least one of A¹, A², A³, A⁴, A⁵, A⁶, A⁷, A⁸ or R²³ is
a bond to the linking group or targeting moiety;

R²⁴, R^{24a}, and R²⁵ are independently selected at each occurrence
from the group: a bond to the linking group, a bond to the

5 targeting moiety, H, C₁-C₆ alkyl, phenyl, benzyl, C₁-C₆ alkoxy,
halide, nitro, cyano, and trifluoromethyl; and

R²⁶ is a co-ordinate bond to a metal or a hydrazine protecting
group; or

a pharmaceutically acceptable salt thereof.

10

(32) A diagnostic agent according to Embodiment 31, wherein:

h' is 1;

w² is NH; and

15 x' is 1.

(33) A diagnostic agent according to any one of embodiments 1-,
wherein:

x is 0;

20 Z is aryl substituted with 0-3 R¹⁶;

k is 1;

g' is 1;

R^{13a}R^{14a} are independently H;

w² is NHC(=O) or -(OCH₂CH₂)₇₆₋₈₄-; and

25 x' is 1.

(34) A diagnostic agent according to any one of embodiments 31-
33, wherein:

w¹ is C=O;

30 g is 2;

R¹³ and R¹⁴ are independently H;

k is 0;

g' is 0;

h' is 1;

w² is NH; and

x' is 1.

(35) A diagnostic agent according to to any one of embodiments
5 31-34, wherein:

2-{{[5-(3-{2-[(6-Hydroxycarbamoyl-7-isobutyl-8-oxo-2-oxa-9-aza-
bicyclo[10.2.2]hexadeca-1(15),12(16),13-triene-10-carbonyl)-
amino]-acetyl-amino)-propylcarbamoyl]-pyridin-2-yl]-
hydrazonomethyl}-benzenesulfonic acid;

10

2-{{[5-(4-{{[(6-Hydroxycarbamoyl-7-isobutyl-8-oxo-2-oxa-9-aza-
bicyclo[10.2.2]hexadeca-1(15),12(16),13-triene-10-carbonyl)-
amino]-methyl}-benzylcarbamoyl]-pyridin-2-yl]-hydrazonomethyl}-
benzenesulfonic acid;

15

2-[7-({N-[3-(2-{[7-(N-hydroxycarbamoyl)(3S,6R,7S)-4-aza-6-(2-
methylpropyl)-11-oxa-5-oxobicyclo[10.2.2]hexadeca-
1(15),12(16),13-trien-3-
yl]carbonylamino}acetyl-amino)propyl]carbamoyl}methyl)-1,4,7,10-
20 tetraaza-4,10-bis(carboxymethyl)cyclododecyl}acetic acid;

20

2-{7-({N-{{[4-({[7-(N-hydroxycarbamoyl)(3S,6R,7S)-4-aza-6-(2-
methylpropyl)-11-oxa-5-oxobicyclo[10.2.2]hexadeca-
1(15),12(16),13-trien-3-yl]-
25 carbonylamino}methyl)phenyl]methyl}carbamoyl}methyl)-1,4,7,10-
tetraaza-4,10-bis(carboxymethyl)cyclododecyl}acetic acid;

25

2-(7-{{N-(1-{N-[3-(2-{[7-(N-hydroxycarbamoyl)(3S,6R,7S)-4-aza-6-
(2-methylpropyl)-11-oxa-5-oxobicyclo[10.2.2]hexadeca-
30 1(15),12(16),13-trien-3-
yl]carbonylamino}acetyl-amino)propyl]carbamoyl}-2-
sulfoethyl)carbamoyl}methyl)-1,4,7,10-tetraaza-4,10-
bis(carboxymethyl)cyclododecyl}acetic acid;

30


35 2-[7-({N-[1-(N-{{[4-({[7-(N-hydroxycarbamoyl)(3S,6R,7S)-4-aza-6-
(2-methylpropyl)-11-oxa-5-oxobicyclo[10.2.2]hexadeca-

- 1 (15), 12 (16), 13-trien-3-yl]-
 carbonylamino)methyl]phenyl)methyl}carbamoyle)-2-
 sulfoethyl}carbamoyle)methyl)-1,4,7,10-tetraaza-4,10-
 bis(carboxymethyl)cyclododecyl}acetic acid;
- 5 2-({2-[(N-[3-(2-([7-(N-hydroxycarbamoyle)(3S,6R,7S)-4-aza-6-(2-
 methylpropyl)-11-oxa-5-oxobicyclo[10.2.2]hexadeca-
 1 (15), 12 (16), 13-trien-3-
 yl]carbonylamino)acetylamino)propyl}carbamoyle)methyl}(carboxymet
 10 hyl)amino)ethyl){2-[bis(carboxymethyl)amino]ethyl}amino}acetic
 acid;
- 2-[(2-[(N-[4-([7-(N-hydroxycarbamoyle)(3S,6R,7S)-4-aza-6-(2-
 methylpropyl)-11-oxa-5-oxobicyclo[10.2.2]hexadeca-
 15 1 (15), 12 (16), 13-trien-3-yl]-
 carbonylamino)methyl]phenyl)methyl}carbamoyle)methyl}(carboxymeth
 yl)amino)ethyl){2-[bis(carboxymethyl)amino]ethyl}amino}acetic
 acid;
- 20 N-[3-(2-([7-(N-hydroxycarbamoyle)(3S,6R,7S)-4-aza-6-(2-
 methylpropyl)-11-oxa-5-oxobicyclo[10.2.2]hexadeca-
 1 (15), 12 (16), 13-trien-3-yl]carbonylamino)acetylamino)propyl]-
 4,5-bis[2-(ethoxyethylthio)acetylamino]pentanamide;
- 25 N-[4-([7-(N-hydroxycarbamoyle)(3S,6R,7S)-4-aza-6-(2-
 methylpropyl)-11-oxa-5-oxobicyclo[10.2.2]hexadeca-
 1 (15), 12 (16), 13-trien-3-yl]carbonylamino)methyl]-phenyl)methyl}-
 4,5-bis[2-(ethoxyethylthio)acetylamino]-pentanamide;
- 30 1-(1,2-Dipalmitoyl-sn-glycero-3-phosphoethanolamino)- α , ω -
 dicarbonylPEG3400-2-([7-(N-hydroxycarbamoyle)(3S,6R,7S)-4-aza-6-
 (2-methylpropyl)-11-oxa-5-oxobicyclo[10.2.2]hexadeca-
 1 (15), 12 (16), 13-trien-3-yl]carbonylamino)-N-(3-
 aminopropyl)acetamide;
- 35

1-(1,2-Dipalmitoyl-sn-glycero-3-phosphoethanolamino)- α,ω -
dicarbonylPEG₃₄₀₀-[7-(N-hydroxycarbamoyl) (3S,6R,7S)-4-aza-6-(2-
methylpropyl)-11-oxa-5-oxobicyclo[10.2.2]hexadeca-
1(15),12(16),13-trien-3-yl]-N-{[4-
5 (aminomethyl)phenyl]methyl}carboxamide conjugate;

2-[2-({5-[N-(5-(N-hydroxycarbamoyl) (5R)-5-(3-[4-(3,4-
dimethoxyphenoxy)phenyl]-3-methyl-2-
oxopyrrolidinyl)pentyl]carbamoyl} (2-pyridyl)amino) (1Z)-2-
10 azavinyl]benzenesulfonic acid;

2-(2-({5-[N-(3-[3-(N-hydroxycarbamoyl) (4S)-4-({4-[(4-
methylphenyl)methoxy]piperidyl)carbonyl]piperidyl)-3-
oxopropyl]carbamoyl} (2-pyridyl)amino) (1Z)-2-
15 azavinyl]benzenesulfonic acid; and



(36) A diagnostic agent according to to any one of embodiments 31-
20 35 wherein the diagnostic metal is selected from the group
consisting of: a paramagnetic metal, a ferromagnetic metal, a
gamma-emitting radioisotope, or an x-ray absorber.

(37) A diagnostic agent according to to any one of embodiments
25 31-36 wherein the diagnostic metal is radioisotope selected from
the group consisting of ^{99m}Tc, ⁹⁵Tc, ¹¹¹In, ⁶²Cu, ⁶⁴Cu, ⁶⁷Ga,
and ⁶⁸Ga.

- (38). A diagnostic agent according to to any one of
embodiments 31-37 further comprising a first ancillary
ligand and a second ancillary ligand capable of stabilizing
the radioisotope.
- 5
- (39) A diagnostic agent according to Embodiment 37, wherein the
radioisotope is ^{99m}Tc .
- (40) A diagnostic agent according to Embodiment 37, wherein the
10 radioisotope is ^{111}In .
- (41) A diagnostic agent according to embodiment 36 wherein the
paramagnetic metal ion is selected from the group
consisting of Gd(III), Dy(III), Fe(III), and Mn(II).
- 15
- (42). A diagnostic agent according to embodiment 36 wherein
the x-ray absorber is a metal is selected from the group
consisting of: Re, Sm, Ho, Lu, Pm, Y, Bi, Pd, Gd, La, Au,
Au, Yb, Dy, Cu, Rh, Ag, and Ir.
- 20
- (43) A diagnostic composition comprising a compound according to
any one of embodiments 1-42 or a pharmaceutically
acceptable salt thereof, and a pharmaceutically acceptable
carrier.
- 25
- (44) A kit comprising a compound of to any one of embodiments 1-
42, or a pharmaceutically acceptable salt form thereof and a
pharmaceutically acceptable carrier.
- 30
- (45) A kit according to Embodiment 44, wherein the kit further
comprises one or more ancillary ligands and a reducing agent.
- (46) A kit according to Embodiment 45, wherein the ancillary
ligands are tricine and TPPTS.
- 35

(47) A kit according to Embodiment 45, wherein the reducing agent is tin(II).

5 (48) A diagnostic agent comprising an echogenic gas and a compound, wherein the compound comprises:

- i) 1-10 targeting moieties;
- ii) a surfactant (Sf); and
- iii) 0-1 linking groups between the targeting moiety and surfactant;

10 wherein the targeting moiety is a matrix metalloproteinase inhibitor; and

wherein the surfactant is capable of forming an echogenic gas filled lipid sphere or microbubble.

15 (49) A diagnostic agent according to embodiment 48, wherein the targeting moiety is a matrix metalloproteinase inhibitor having an inhibitory constant K_i of <1000 nM.

20 (50) A diagnostic agent according to any one of embodiments 48-49, wherein the targeting moiety is a matrix metalloproteinase inhibitor having an inhibitory constant K_i of <100 nM.

(51) A diagnostic agent according to embodiment 48, comprising 1-5 targeting moieties.

25


(52). A diagnostic agent according to embodiment 48, comprising one targeting moiety.

30 (53) A diagnostic agent according to any one of embodiments 48-52, wherein the targeting moiety is an inhibitor of one or more matrix metalloproteinases selected from the group consisting of MMP-1, MMP-2, MMP-3, MMP-9, and MMP-14.

35 (54) A diagnostic agent according to any one of embodiments 48-53, wherein the targeting moiety is an inhibitor of one or more

matrix metalloproteinases selected from the group consisting of MMP-2, MMP-9, and MMP-14.

(55) A diagnostic agent according to embodiment 48, wherein the
5 targeting moiety is of the formulae (Ia) or (Ib):




R is independently OH or $-\text{CH}_2\text{SH}$;

10

R^1 is independently selected at each occurrence from the group:
H, OH, C₁-3 alkyl, C₂-3 alkenyl, C₂-3 alkynyl, and
heterocycle-S-CH₂-;

15 R^2 is independently C₁-20 alkyl;

X is independently C=O or SO₂, provided when X is C=O, R^3 is



, and when X is SO₂, R^3 is independently selected
from the group: aryl substituted with 0-2 R^6 , and
20 heterocycle substituted with 0-2 R^6 ;

R^4 is independently selected at each occurrence from the group:
C₁-6 alkyl, phenyl, and benzyl;

25 R^5 is independently at each occurrence from the group: NH(C₁-6
alkyl), NH-phenyl, and NH-heterocycle; wherein said alkyl,
phenyl and heterocycle groups are optionally substituted
with a bond to the linking group or a bond to the
surfactant;

30

R⁶ is independently aryloxy substituted with 0-3 R⁷;

R⁷ is independently halogen or methoxy;

5 or alternatively,

R¹ and R⁴ may be taken together to form a bridging group of the formula $-(CH_2)_3-O-phenyl-CH_2-$, optionally substituted with a bond to the linking group or a bond to the surfactant;

10

or alternatively,

R¹ and R² may be taken together to form a bridging group of the formula $-(CH_2)_3-NH-$, optionally substituted with a bond to the linking group or a bond to the surfactant; or

R¹ and R² taken together with the nitrogen and carbon atom through which they are attached form a C₅₋₇ atom saturated ring system substituted with one or more substituents selected from the group consisting of: a bond to Ln, a bond to Sf, and $-C(=O)-NR^{29}R^{30}$;

R⁸ is independently at each occurrence OH or phenyl, optionally substituted with a bond to the linking group or a bond to the surfactant, provided that when R⁸ is phenyl, R¹⁰ is $-C(=O)-CR^{12}-NH-CH(CH_3)-COOH$;

R⁹ and R^{9'} are independently H, C₁₋₆ alkyl optionally substituted with a bond to the linking group or a bond to the surfactant, or are taken together with the carbon atom to which R⁹ and R^{9'} are attached to form a 5-7 atom saturated, partially unsaturated or aromatic ring system containing 0-3 heteroatoms selected from O, N, SO₂ and S, said ring system substituted with R⁶ and optionally

substituted with a bond to the linking group or a bond to the surfactant;

R¹⁰ and R¹¹ are independently H, or C₁₋₆ alkyl optionally substituted with a bond to the linking group or a bond to the surfactant, or are taken together with the nitrogen atom to which they are attached to form a 5-7 atom saturated, partially unsaturated or aromatic ring system containing 0-3 heteroatoms selected from O, N, SO₂ and S, said ring system optionally substituted with 0-3 R²⁷, a bond to the linking group or a bond to the surfactant;

or alternatively,

R⁹ and R¹⁰ are taken together with the carbon atom to which they are attached to form a 5-7 atom saturated, partially unsaturated or aromatic ring system containing 0-3 heteroatoms selected from O, N, SO₂ and S, said ring system optionally substituted with a bond to the linking group or a bond to the surfactant; and

R¹² is independently C₁₋₂₀ alkyl;

R²⁷ is =O, C₁₋₄ alkyl, or phenyl substituted with R²⁸;


R²⁸ is a phenoxy group substituted with 0-2 OCH₃ groups;

R²⁹ and R³⁰ taken together with the nitrogen atom through which they are attached form a C₅₋₇ atom saturated ring system substituted with R³¹; and

R³¹ is a benzyloxy group substituted with C₁₋₄ alkyl.

30

(56) A diagnostic agent according to embodiment 55 wherein wherein the targeting moiety is a matrix metalloproteinase inhibitor of the formulae (Ia) or (Ib):



R is OH;

- 5 R^1 is independently selected at each occurrence from the group:
H, OH, C₁₋₃ alkyl, C₂₋₃ alkenyl, C₂₋₃ alkynyl, and
heterocycle-S-CH₂-;

R^2 is independently C₁₋₆ alkyl;

10

X is C=O;

R^4 is independently selected at each occurrence from the group:
C₁₋₆ alkyl, phenyl, and benzyl;

15

R^5 is independently at each occurrence from the group: NH(C₁₋₆
alkyl), NH-phenyl, and NH-heterocycle; wherein said alkyl,
phenyl and heterocycle groups are optionally substituted
with a bond to the linking group or a bond to the
20 surfactant;

R^6 is independently aryloxy substituted with 0-3 R^7 ;

R^7 is independently halogen or methoxy;

25

or alternatively,

R^1 and R^4 may be taken together to form a bridging group of the
formula -(CH₂)₃-O-phenyl-CH₂-, optionally substituted with a
30 bond to the linking group or a bond to the surfactant;

or alternatively,

R¹ and R² may be taken together to form a bridging group of the formula $-(CH_2)_3-NH-$, optionally substituted with a bond to the linking group or a bond to the surfactant; or

5

R¹ and R² taken together with the nitrogen and carbon atom through which they are attached form a C₅₋₇ atom saturated ring system substituted with one or more substituents selected from the group consisting of: a bond to Ln, a bond to Sf, and $-C(=O)-NR^{29}R^{30}$;

10

R⁸ is OH;

R⁹ and R^{9'} are independently H, C₁₋₆ alkyl optionally substituted with a bond to the linking group or a bond to the surfactant, or are taken together with the carbon atom to which R⁹ and R^{9'} are attached to form a 5-7 atom saturated, partially unsaturated or aromatic ring system containing 0-1 heteroatoms selected from O, N, , said ring system optionally substituted with a bond to the linking group or a bond to the surfactant;

15

20

R¹⁰ and R¹¹ are independently H, or C₁₋₆ alkyl optionally substituted with a bond to the linking group or a bond to the surfactant, or are taken together with the nitrogen atom to which they are attached to form a 5-7 atom saturated, partially unsaturated or aromatic ring system containing 0-1 heteroatoms selected from O, N, , said ring system optionally substituted with 0-3 R²⁷, a bond to the linking group or a bond to the surfactant;

25

30

or alternatively,

R⁹ and R¹⁰ are taken together with the carbon atom to which they are attached to form a 5-7 atom saturated, partially unsaturated or aromatic ring system containing 0-1 heteroatoms selected from O, N, , said ring system optionally substituted with a bond to the linking group or a bond to the surfactant; and

R¹² is independently C₁₋₆ alkyl;
 R²⁷ is =O, C₁₋₄ alkyl, or phenyl substituted with R²⁸;
 10 R²⁸ is a phenoxy group substituted with 0-2 OCH₃ groups;
 R²⁹ and R³⁰ taken together with the nitrogen atom through which they are attached form a C₅₋₇ atom saturated ring system substituted with R³¹; and
 R³¹ is a benzyloxy group substituted with C₁₋₄ alkyl.

15


(57) A diagnostic agent according to any one of embodiments 55-57 wherein the targeting moiety is a matrix metalloproteinase inhibitor of the formulae (Ia) or (Ib):

20 wherein:

R is -OH;

R² is C₁₋₆ alkyl;

X is C=O;



25 R¹ and R⁴ are taken together to form a bridging group of formula -(CH₂)₃-O-phenyl-CH₂-;

R⁵ is NH(C₁₋₆alkyl), substituted with a bond to the linking group or a bond to the surfactant.

30 (58) A diagnostic agent according to any one of embodiments 55-57 wherein:

R is -OH;

R⁹ is C₁ alkyl substituted with a bond to Ln;

R¹⁰ and R¹¹ taken together with the nitrogen atom to which they are attached form a 5 atom saturated ring system, said ring system is substituted with 0-3 R²⁷;

- 5 R²⁷ is =O, C₁-4 alkyl, or phenyl substituted with R²⁸; and
R²⁸ is a phenoxy group substituted with 0-2 OCH₃ groups.

(59) A diagnostic agent according to any one of embodiments 55-58 wherein

- 10 R is -OH;


R¹ and R² taken together with the nitrogen and carbon atom through which they are attached form a C₅₋₇ atom saturated ring system substituted with one or more substituents selected from the group consisting of: a bond to Ln, a bond to Sf, and -C(=O)-
15 NR²⁹R³⁰;

R²⁹ and R³⁰ taken together with the nitrogen atom through which they are attached form a C₅₋₇ atom saturated ring system substituted with R³¹; and

R³¹ is a benzyloxy group substituted with C₁-4 alkyl.

20

(60) A diagnostic agent according to any one of embodiments 48-59 wherein the linking group is of the formula:



25

W¹ and W² are independently selected at each occurrence from the group: O, S, NH, NHC(=O), C(=O)NH, NR¹⁵C(=O), C(=O)NR¹⁵, C(=O), C(=O)O, OC(=O), NHC(=S)NH, NHC(=O)NH, SO₂, SO₂NH, - (OCH₂CH₂)₇₆₋₈₄, (OCH₂CH₂)_s, (CH₂CH₂O)_{s'}, (OCH₂CH₂CH₂)_{s''},
30 (CH₂CH₂CH₂O)_t, and (aa)_{t'};

aa is independently at each occurrence an amino acid;

- Z is selected from the group: aryl substituted with 0-3 R¹⁶, C₃-10 cycloalkyl substituted with 0-3 R¹⁶, and a 5-10 membered heterocyclic ring system containing 1-4 heteroatoms independently selected from N, S, and O and substituted with 0-3 R¹⁶;
- R¹³, R^{13a}, R¹⁴, R^{14a}, and R¹⁵ are independently selected at each occurrence from the group: H, =O, COOH, SO₃H, PO₃H, C₁-C₅ alkyl substituted with 0-3 R¹⁶, aryl substituted with 0-3 R¹⁶, benzyl substituted with 0-3 R¹⁶, and C₁-C₅ alkoxy substituted with 0-3 R¹⁶, NHC(=O)R¹⁷, C(=O)NHR¹⁷, NHC(=O)NHR¹⁷, NHR¹⁷, R¹⁷, and a bond to the surfactant;
- R¹⁶ is independently selected at each occurrence from the group: a bond to the surfactant, COOR¹⁷, C(=O)NHR¹⁷, NHC(=O)R¹⁷, OH, NHR¹⁷, SO₃H, PO₃H, -OPO₃H₂, -OSO₃H, aryl substituted with 0-3 R¹⁷, C₁-5 alkyl substituted with 0-1 R¹⁸, C₁-5 alkoxy substituted with 0-1 R¹⁸, and a 5-10 membered heterocyclic ring system containing 1-4 heteroatoms independently selected from N, S, and O and substituted with 0-3 R¹⁷;
- R¹⁷ is independently selected at each occurrence from the group: H, alkyl substituted with 0-1 R¹⁸, aryl substituted with 0-1 R¹⁸, a 5-10 membered heterocyclic ring system containing 1-4 heteroatoms independently selected from N, S, and O and substituted with 0-1 R¹⁸, C₃-10 cycloalkyl substituted with 0-1 R¹⁸, polyalkylene glycol substituted with 0-1 R¹⁸, carbohydrate substituted with 0-1 R¹⁸, cyclodextrin substituted with 0-1 R¹⁸, amino acid substituted with 0-1 R¹⁸, polycarboxyalkyl substituted with 0-1 R¹⁸, polyazaalkyl substituted with 0-1 R¹⁸, peptide

substituted with 0-1 R¹⁸, wherein the peptide is comprised of 2-10 amino acids, 3,6-O-disulfo-B-D-galactopyranosyl, bis(phosphonomethyl)glycine, and a bond to the surfactant;

5 R¹⁸ is a bond to the surfactant;

k is selected from 0, 1, and 2;

h is selected from 0, 1, and 2;

h' is selected from 0, 1, and 2;

10 g is selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;

g' is selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;

s is selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;

s' is selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;

s" is selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;

15 t is selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;

t' is selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;

x is selected from 0, 1, 2, 3, 4, and 5; and

x' is selected from 0, 1, 2, 3, 4, and 5.

20 (61) A diagnostic agent according to any one of embodiments 48-60 wherein

W¹ and W² are independently selected at each occurrence from the group: O, NH, NHC(=O), C(=O)NH, NR¹⁵C(=O), C(=O)NR¹⁵, C(=O), C(=O)O, OC(=O), NHC(=S)NH, NHC(=O)NH, SO₂, -

25 (CH₂CH₂O)₇₆₋₈₄, (OCH₂CH₂)_s, (CH₂CH₂O)_{s'}, (OCH₂CH₂CH₂)_{s"}, (CH₂CH₂CH₂O)_t, and (aa)_{t'};

aa is independently at each occurrence an amino acid;

30 Z is selected from the group: aryl substituted with 0-1 R¹⁶, C₃₋₁₀ cycloalkyl substituted with 0-1 R¹⁶, and a 5-10 membered heterocyclic ring system containing 1-4 heteroatoms independently selected from N, S, and O and substituted with 0-1 R¹⁶;

35

R¹³, R^{13a}, R¹⁴, R^{14a}, and R¹⁵ are independently selected at each occurrence from the group: H, =O, COOH, SO₃H, C₁-C₅ alkyl substituted with 0-1 R¹⁶, aryl substituted with 0-1 R¹⁶, benzyl substituted with 0-1 R¹⁶, and C₁-C₅ alkoxy substituted with 0-1 R¹⁶, NHC(=O)R¹⁷, C(=O)NHR¹⁷, NHC(=O)NHR¹⁷, NHR¹⁷, R¹⁷, and a bond to the surfactant;

k is 0 or 1;
s is selected from 0, 1, 2, 3, 4, and 5;
10 s' is selected from 0, 1, 2, 3, 4, and 5;
s" is selected from 0, 1, 2, 3, 4, and 5; and
t is selected from 0, 1, 2, 3, 4, and 5.

(62) A diagnostic agent according to embodiment 60
15 wherein:
W¹ is C(=O)NR¹⁵;
h is 1;
g is 3;
R¹³ and R¹⁴ are independently H;
20 x is 1;
k is 0;
g' is 0;
h' is 1;
W² is NH; and
25 x' is 1.

(63) A diagnostic agent according to embodiment 60
x is 0;
k is 1;
30 Z is aryl substituted with 0-3 R¹⁶;
g' is 1;
W² is NH;
R^{13a} and R^{14a} are independently H;
h' is 1; and

x' is 1.

(64) A diagnostic agent according to embodiment 60

W¹ is C(=O)NR¹⁵;

5 h is 1;

g is 2;

R¹³ and R¹⁴ are independently H;

x is 1;

k is 0;

10 g' is 1;

R^{13a} and R^{14a} are independently H; or C1-5 alkyl substituted
with 0-3 R¹⁶;

R¹⁶ is SO₃H;

W² is NHC(=O) or NH;

15 h' is 1; and

x' is 2.

(65) A diagnostic agent according to embodiment 60

W¹ is C(=O)NH;

20 h is 1;

g is 3;

R¹³ and R¹⁴ are independently H;

k is 0;

g' is 0;

25 x is 1;

W² is -NH(C=O)- or -(OCH₂CH₂)₇₆₋₈₄-;

h' is 2; and

x' is 1.

30 (66) A diagnostic agent according to embodiment 60

x is 0;

k is 0;

g' is 3;

h' is 1;

W^2 is NH; and
 x' is 1.

(67) A diagnostic agent according to embodiment 60
 5 x is 0;
 Z is aryl substituted with 0-3 R^{16} ;
 k is 1;
 g' is 1;
 $R^{13a}R^{14a}$ are independently H;
 10 W^2 is $NHC(=O)$ or $-(OCH_2CH_2)_{76-84}-$; and
 x' is 1.

(68) A diagnostic agent according to embodiment 60
 W^1 is $C=O$;
 15 g is 2;
 R^{13} and R^{14} are independently H;
 k is 0;
 g' is 0;
 h' is 1;
 20 W^2 is NH; and
 x' is 1.

(69) A diagnostic agent according to embodiment 48 wherein the
 linking group is present.

25 (70) A diagnostic agent according to any one of embodiments 48-
 69 wherein

S_f is a surfactant which is a lipid or a compound of the
 30

formula: $A^9 \begin{array}{c} \diagup \\ E^9 - A^{10} \end{array}$;

A^9 is selected from the group: OH and OR^{32} ;

A¹⁰ is OR³²;

R³² is C(=O)C₁₋₂₀ alkyl;

5

E⁹ is C₁₋₁₀ alkylene substituted with 1-3 R³³;

R³³ is independently selected at each occurrence from the group:

10 R³⁵, -PO₃H-R³⁵, =O, -CO₂R³⁴, -C(=O)R³⁴, -C(=O)N(R³⁴)₂,
-CH₂OR³⁴, -OR³⁴, -N(R³⁴)₂, C₁₋₅ alkyl, and C₂₋₄ alkenyl;

R³⁴ is independently selected at each occurrence from the group:

R³⁵, H, C₁₋₆ alkyl, phenyl, benzyl, and trifluoromethyl;

15 R³⁵ is a bond to L_n;

and a pharmaceutically acceptable salt thereof.

20 (71) A diagnostic agent according to any one of
embodiments 48-70 wherein the surfactant is a lipid or a
compound of the

formula:
$$\begin{array}{c} \text{E}^9 - \text{A}^{10} \\ \diagup \\ \text{A}^9 \end{array} ;$$

25

A⁹ is OR³²;

A¹⁰ is OR³²;

30 R³² is C(=O)C₁₋₁₅ alkyl;

E⁹ is C₁₋₄ alkylene substituted with 1-3 R³³;

R^{33} is independently selected at each occurrence from the group:

R^{35} , $-\text{PO}_3\text{H}-R^{35}$, $=\text{O}$, $-\text{CO}_2R^{34}$, $-\text{C}(=\text{O})R^{34}$, $-\text{CH}_2\text{OR}^{34}$, $-\text{OR}^{34}$,
and $\text{C}_1\text{-C}_5$ alkyl;

5


R^{34} is independently selected at each occurrence from the group:

R^{35} , H, $\text{C}_1\text{-C}_6$ alkyl, phenyl, and benzyl; and

R^{35} is a bond to L_n .

10

(72) A diagnostic agent according to any one of embodiments 48-71, wherein



15 wherein:

A^1 is a bond to L_n ;

E^1 is C_1 alkyl substituted by R^{23} ;

A^2 is NH ;

E^2 is C_2 alkyl substituted with $0\text{-}1R^{23}$;

20 A^3 is $-\text{O}-\text{P}(\text{O})(R^{21})-\text{O}-$;

E^3 is C_1 alkyl;

A^4 and A^5 are each $-\text{O}-$;

E^4 and E^6 are each independently C_{1-16} alkyl substituted with $0\text{-}1R^{23}$;


25 E^5 is C_1 alkyl;

A^5 is $-\text{O}-$;

R^{21} is $-\text{OH}$; and


R^{23} is $=\text{O}$.

(73) A diagnostic agent according to embodiment 48 wherein the compound is of the formula:



5

wherein, Q is a compound of Formulae (Ia) or (Ib):




10 R is independently OH or $-CH_2SH$;

R^1 is independently selected at each occurrence from the group:
H, OH, C_1-3 alkyl, C_2-3 alkenyl, C_2-3 alkynyl, and
heterocycle-S- CH_2- ;

15

R^2 is independently C_1-20 alkyl;

X is independently C=O or SO_2 , provided when X is C=O, R^3 is



20

, and when X is SO_2 , R^3 is independently selected
from the group: aryl substituted with 0-2 R^6 , and
heterocycle substituted with 0-2 R^6 ;

R^4 is independently selected at each occurrence from the group:
 C_1-6 alkyl, phenyl, and benzyl;

25

R^5 is independently at each occurrence from the group: $NH(C_1-6$
alkyl), NH -phenyl, and NH -heterocycle; wherein said alkyl,
phenyl and heterocycle groups are optionally substituted
with a bond to L_n ;

30

R⁶ is independently aryloxy substituted with 0-3 R⁷;

R⁷ is independently halogen or methoxy;

5 or alternatively,

R¹ and R⁴ may be taken together to form a bridging group of the formula $-(CH_2)_3-O-phenyl-CH_2-$, optionally substituted with a bond to L_n;

10

or alternatively,

R¹ and R² may be taken together to form a bridging group of the formula $-(CH_2)_3-NH-$, optionally substituted with a bond to L_n; or

15

R¹ and R² taken together with the nitrogen and carbon atom through which they are attached form a C₅₋₇ atom saturated ring system substituted with one or more substituents selected from the group consisting of: a bond to L_n, a bond to Sf, and $-C(=O)-NR^{29}R^{30}$;

20

R⁸ is independently at each occurrence OH or phenyl, optionally substituted with a bond to L_n, provided that when R⁸ is phenyl, R¹⁰ is $-C(=O)-CR^{12}-NH-CH(CH_3)-COOH$;

25

R⁹ and R^{9'} are independently H, C₁₋₆ alkyl optionally substituted with a bond to L_n, or are taken together with the carbon atom to which they are attached to form a 5-7 atom saturated, partially unsaturated or aromatic ring system containing 0-3 heteroatoms selected from O, N, SO₂ and S, said ring system substituted with R⁶ and optionally substituted with a bond to L_n;

30

R¹⁰ and R¹¹ are independently H, or C₁₋₆ alkyl optionally substituted with a bond to L_n, or are taken together with the nitrogen atom to which they are attached to form a 5-7 atom saturated, partially unsaturated or aromatic ring system containing 0-3 heteroatoms selected from O, N, SO₂ and S, said ring system optionally substituted with 0-3 R²⁷ or a bond to L_n;


or alternatively,

R⁹ and R¹⁰ are taken together with the carbon atom to which they are attached to form a 5-7 atom saturated, partially unsaturated or aromatic ring system containing 0-3 heteroatoms selected from O, N, SO₂ and S, said ring system optionally substituted with a bond to L_n;

R¹² is independently C₁₋₂₀ alkyl;

d is selected from 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;

L_n is a linking group having the formula:



W¹ and W² are independently selected at each occurrence from the group: O, S, NH, NHC(=O), C(=O)NH, NR¹⁵C(=O), C(=O)NR¹⁵, C(=O), C(=O)O, OC(=O), NHC(=S)NH, NHC(=O)NH, SO₂, SO₂NH, - (OCH₂CH₂)₇₆₋₈₄, (OCH₂CH₂)_s, (CH₂CH₂O)_{s'}, (OCH₂CH₂CH₂)_{s''}, (CH₂CH₂CH₂O)_t, and (aa)_{t'};

aa is independently at each occurrence an amino acid;

Z is selected from the group: aryl substituted with 0-3 R¹⁶, C₃₋₁₀ cycloalkyl substituted with 0-3 R¹⁶, and a 5-10

membered heterocyclic ring system containing 1-4 heteroatoms independently selected from N, S, and O and substituted with 0-3 R¹⁶;

5 R¹³, R^{13a}, R¹⁴, R^{14a}, and R¹⁵ are independently selected at each occurrence from the group: H, =O, COOH, SO₃H, PO₃H, C₁-C₅ alkyl substituted with 0-3 R¹⁶, aryl substituted with 0-3 R¹⁶, benzyl substituted with 0-3 R¹⁶, and C₁-C₅ alkoxy substituted with 0-3 R¹⁶, NHC(=O)R¹⁷, C(=O)NHR¹⁷,
10 NHC(=O)NHR¹⁷, NHR¹⁷, R¹⁷, and a bond to Sf;

R¹⁶ is independently selected at each occurrence from the group: a bond to Sf, COOR¹⁷, C(=O)NHR¹⁷, NHC(=O)R¹⁷, OH, NHR¹⁷, SO₃H, PO₃H, -OPO₃H₂, -OSO₃H, aryl substituted with 0-3 R¹⁷,
15 C₁-5 alkyl substituted with 0-1 R¹⁸, C₁-5 alkoxy substituted with 0-1 R¹⁸, and a 5-10 membered heterocyclic ring system containing 1-4 heteroatoms independently selected from N, S, and O and substituted with 0-3 R¹⁷;

20 R¹⁷ is independently selected at each occurrence from the group: H, alkyl substituted with 0-1 R¹⁸, aryl substituted with 0-1 R¹⁸, a 5-10 membered heterocyclic ring system containing 1-4 heteroatoms independently selected from N, S, and O and substituted with 0-1 R¹⁸, C₃-10 cycloalkyl
25 substituted with 0-1 R¹⁸, polyalkylene glycol substituted with 0-1 R¹⁸, carbohydrate substituted with 0-1 R¹⁸, cyclodextrin substituted with 0-1 R¹⁸, amino acid substituted with 0-1 R¹⁸, polycarboxyalkyl substituted with 0-1 R¹⁸, polyazaalkyl substituted with 0-1 R¹⁸, peptide
30 substituted with 0-1 R¹⁸, wherein the peptide is comprised of 2-10 amino acids, 3,6-O-disulfo-B-D-galactopyranosyl, bis(phosphonomethyl)glycine, and a bond to Sf;

R¹⁸ is a bond to Sf;

k is selected from 0, 1, and 2;

5 h is selected from 0, 1, and 2;

h' is selected from 0, 1, and 2;

g is selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;

g' is selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;

s is selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;

10 s' is selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;

s" is selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;


t is selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;

t' is selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;

x is selected from 0, 1, 2, 3, 4, and 5;

15 x' is selected from 0, 1, 2, 3, 4, and 5;

Sf is a surfactant which is a lipid or a compound of the



20

A⁹ is selected from the group: OH and OR³²;

A¹⁰ is OR³²;

25 R³² is C(=O)C₁₋₂₀ alkyl;

E⁹ is C₁₋₁₀ alkylene substituted with 1-3 R³³;

R³³ is independently selected at each occurrence from the group:

30 R³⁵, -PO₃H-R³⁵, =O, -CO₂R³⁴, -C(=O)R³⁴, -C(=O)N(R³⁴)₂,
-CH₂OR³⁴, -OR³⁴, -N(R³⁴)₂, C₁-C₅ alkyl, and C₂-C₄ alkenyl;


R^{34} is independently selected at each occurrence from the group:

R³⁵, H, C₁-C₆ alkyl, phenyl, benzyl, and trifluoromethyl;

R^{35} is a bond to L_n ; or

5

Sf is of the formula:



wherein:

A¹ is a bond to Ln;

10 E¹ is C₁ alkyl substituted by R²³;

A^2 is NH;

E^2 is C_2 alkyl sunsttuted wth 0-1R²³;

$$A^3 \text{ is } -O-P(O)(R^{21})-O;$$

E^3 is C_1 alkyl;

15 A^4 and A^5 are each -0-;

E⁴ and E⁶ are each independently C₁₋₁₆ alkyl substituted with 0-1R²³;

E⁵ is C₁ alkyl;

A⁵ is -0-;

20 R^{21} is $-OH$; and

$$R^{23} \text{ is } = 0; \text{ or}$$

a pharmaceutically acceptable salt thereof.


(74) A diagnostic agent according to embodiment 73,

25 wherein:

R is -OH;

R² is C1-6 alkyl;

X is C=0;



R¹ and R⁴ are taken together to form a bridging group of formula
-(CH₂)₃-O-phenyl-CH₂-;

R⁵ is NH(C1-6alkyl), substituted with a bond to the linking
5 group or a bond to the surfactant.

(75) A diagnostic agent according to any one of
embodiments 73-74, wherein:

R is -OH;


10 R⁹ is C₁ alkyl substituted with a bond to Ln;

R¹⁰ and R¹¹ taken together with the nitrogen atom to which they
are attached form a 5 atom saturated ring system, said right
system is substituted with 0-3 R²⁷;

R²⁷ is =O, C1-4 alkyl, or phenyl substituted with R²⁸; and

15 R²⁸ is a phenoxy group substituted with 0-2 OCH₃ groups;

S_f is a surfactant which is a lipid or a compound of the



20

A⁹ is OR³²;

A¹⁰ is OR³²;

25 R³² is C(=O)C1-15 alkyl;

E⁹ is C1-4 alkylene substituted with 1-3 R³³;

R^{33} is independently selected at each occurrence from the group:
 R^{35} , $-PO_3H-R^{35}$, $=O$, $-CO_2R^{34}$, $-C(=O)R^{34}$, $-CH_2OR^{34}$, $-OR^{34}$,
 and C_1 - C_5 alkyl;

5 R^{34} is independently selected at each occurrence from the group:
 R^{35} , H , C_1 - C_6 alkyl, phenyl, and benzyl; and

R^{35} is a bond to L_n .

10 (76) . A diagnostic agent according according to any one
 of embodiments 73-75, wherein:

R is $-OH$;


R^9 is C_1 alkyl substituted with a bond to L_n ;

R^{10} and R^{11} taken together with the nitrogen atom to which they
 15 are attached form a 5 atom saturated ring system, said right
 system is substituted with 0-3 R^{27} ;

R^{27} is $=O$, C_1 -4 alkyl, or phenyl substituted with R^{28} ; and

R^{28} is a phenoxy group substituted with 0-2 OCH_3 groups;

20 S_f is a surfactant which is a lipid or a compound of the
 of the formula:



wherein:

A^1 is a bond to L_n ;

25 E^1 is C_1 alkyl substituted by R^{23} ;

A^2 is NH ;

E^2 is C_2 alkyl substituted with 0-1 R^{23} ;

A^3 is $-O-P(O)(R^{21})-O$;

E^3 is C_1 alkyl;

A⁴ and A⁵ are each -O-;

E⁴ and E⁶ are each independently C₁₋₁₆ alkyl substituted with 0-1R²³;

E⁵ is C₁ alkyl;

5 A⁵ is -O-;

R²¹ is -OH; and

R²³ is =O.

(77) A diagnostic agent according to any one of
10 embodiments 73-76, wherein:

wherein

R is -OH;

R¹ and R² taken together with the nitrogen and carbon atom
through which they are attached form a C₅₋₇ atom saturated ring
15 system substituted with one or more substituents selected from
the group consisting of: a bond to Ln, a bond to Sf, and -C(=O)-
NR²⁹R³⁰;

R²⁹ and R³⁰ taken together with the nitrogen atom through which
they are attached form a C₅₋₇ atom saturated ring system
20 substituted with R³¹; and

R³¹ is a benzyloxy group substituted with C₁₋₄ alkyl.

d is selected from 1, 2, 3, 4, and 5;

25 W is independently selected at each occurrence from the group:

O, NH, NHC(=O), C(=O)NH, NR¹⁵C(=O), C(=O)NR¹⁵, C(=O),
C(=O)O, OC(=O), NHC(=S)NH, NHC(=O)NH, SO₂, (OCH₂CH₂)_s,
(CH₂CH₂O)_s['], (OCH₂CH₂CH₂)_s["], (CH₂CH₂CH₂O)_t, and (aa)_t['];

30 aa is independently at each occurrence an amino acid;

Z is selected from the group: aryl substituted with 0-1 R¹⁶,
C₃₋₁₀ cycloalkyl substituted with 0-1 R¹⁶, and a 5-10
membered heterocyclic ring system containing 1-4

heteroatoms independently selected from N, S, and O and substituted with 0-1 R¹⁶;

5 R¹³, R^{13a}, R¹⁴, R^{14a}, and R¹⁵ are independently selected at each occurrence from the group: H, =O, COOH, SO₃H, C₁-C₅ alkyl substituted with 0-1 R¹⁶, aryl substituted with 0-1 R¹⁶, benzyl substituted with 0-1 R¹⁶, and C₁-C₅ alkoxy substituted with 0-1 R¹⁶, NHC(=O)R¹⁷, C(=O)NHR¹⁷, NHC(=O)NHR¹⁷, NHR¹⁷, R¹⁷, and a bond to Sf;
10
k is 0 or 1;
s is selected from 0, 1, 2, 3, 4, and 5;
s' is selected from 0, 1, 2, 3, 4, and 5;
s" is selected from 0, 1, 2, 3, 4, and 5; and
15 t is selected from 0, 1, 2, 3, 4, and 5.

(78) A diagnostic agent according to according to any one of embodiments 73-77, wherein:

20 W¹ is C(=O)NR¹⁵;
h is 1;
g is 3;
R¹³ and R¹⁴ are independently H;
x is 1;
25 k is 0;
g' is 0;
h' is 1;
W² is NH; and
x' is 1.

30 (79) A diagnostic agent according to embodiment 73, wherein:
x is 0;
k is 1;

Z is aryl substituted with 0-3 R¹⁶;
g' is 1;
W² is NH;
R^{13a} and R^{14a} are independently H;
5 h' is 1; and
x' is 1.

(80) A diagnostic agent according to Embodiment 73,
wherein:
10 W¹ is C(=O)NR¹⁵;
h is 1;
g is 2;
R¹³ and R¹⁴ are independently H;
x is 1;
15 k is 0;
g' is 1;
R^{13a} and R^{14a} are independently H; or C1-5 alkyl substituted
with 0-3 R¹⁶;
R¹⁶ is SO₃H;
20 W² is NHC(=O) or NH;
h' is 1; and
x' is 2.

(81) A diagnostic agent according to Embodiment 73,
25 wherein:
W¹ is C(=O)NH;
h is 1;
g is 3;
R¹³ and R¹⁴ are independently H;
30 k is 0;
g' is 0;
x is 1;
W² is -NH(C=O)- or -(OCH₂CH₂)₇₆₋₈₄-;
h' is 2; and

x' is 1.

(82) A diagnostic agent according to Embodiment 73,
wherein:

5 x is 0;
k is 0;
g' is 3;
h' is 1;
W² is NH; and
10 x' is 1.


(83) A diagnostic agent according to Embodiment 73,
wherein:

x is 0;
15 Z is aryl substituted with 0-3 R¹⁶;
k is 1;
g' is 1;
R^{13a}R^{14a} are independently H;
W² is NHC(=O) or -(OCH₂CH₂)₇₆₋₈₄-; and
20 x' is 1.

(84) A diagnostic agent according to Embodiment 73,
wherein:

W¹ is C=O;
25 g is 2;
R¹³ and R¹⁴ are independently H;
k is 0;
g' is 0;
h' is 1;
30 W² is NH; and
x' is 1.

(85) A diagnostic agent according to Embodiment 1, wherein
35 the compound is selected from the group consisting of:



- (86) A diagnostic agent according to embodiment 48,
 5 wherein: wherein the echogenic gas is a perfluorocarbon gas or sulfur hexafluoride.
- (87) A diagnostic agent according to embodiment 86 wherein said
 10 perfluorocarbon is selected from the group consisting of perfluoromethane, perfluoroethane, perfluoropropane, perfluorobutane, perfluorocyclobutane, perfluoropentane, and perfluorohexane.
- (88) A diagnostic composition comprising a compound according to
 15 embodiment 48 or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier.
- (89) A diagnostic composition comprising a compound
 20 according to embodiment 48 or a pharmaceutically acceptable salt form thereof, an echogenic gas and a pharmaceutically acceptable carrier.
- (90) A diagnostic composition comprising a compound according to
 25 embodiment 48 further comprising: 1,2-dipalmitoyl-sn-glycero-3-phosphatidic acid, 1,2-dipalmitoyl-sn-glycero-3-phosphatidylcholine, and N-(methoxypolyethylene glycol 5000 carbamoyl)-1,2-dipalmitoyl-sn-glycero-3-phosphatidylethanolamine.

- (91) A method of detecting, imaging or monitoring the presence of matrix metalloproteinase in a patient comprising the steps of:
- 5 a) administering to said patient a diagnostic agent of embodiment 1; and
- b) acquiring an image of a site of concentration of said diagnostic agent in the patient by a diagnostic imaging technique.
- 10 (92) A method of detecting, imaging or monitoring the presence of matrix metalloproteinase in a patient comprising the steps of:
- 15 a) administering to said patient a diagnostic agent of embodiment 48; and
- b) acquiring an image of a site of concentration of said diagnostic agent in the patient by a diagnostic imaging technique.
- 20 (93) A method of detecting, imaging or monitoring a pathological disorder associated with matrix metalloproteinase activity in a patient comprising the steps of:
- 25 a) administering to said patient a diagnostic agent of embodiment 1; and
- b) acquiring an image of a site of concentration of said diagnostic agent in the patient by a diagnostic imaging technique.
- 30 (94) A method of detecting, imaging or monitoring a pathological disorder associated with matrix metalloproteinase activity in a patient comprising the steps of:
- a) administering to said patient a diagnostic agent according to embodiment 48; and

- b) acquiring an image of a site of concentration of said diagnostic agent in the patient by a diagnostic imaging technique.

5 (95) A method of detecting, imaging or monitoring atherosclerosis in a patient comprising the steps of:
a) administering a diagnostic agent according to embodiment 1; and
b) acquiring an image of a site of concentration of said
10 diagnostic agent in the body by a diagnostic imaging technique.

(96) A method of detecting, imaging or monitoring atherosclerosis in a patient comprising the steps of:
15 a) administering a diagnostic agent according to embodiment 48; and
b) acquiring an image of a site of concentration of said diagnostic agent in the body by a diagnostic imaging technique.

20 (97) A method according to embodiment 95, wherein the atherosclerosis is coronary atherosclerosis or cerebrovascular atherosclerosis.

25 (98) A method according to embodiment 96, wherein the atherosclerosis is coronary atherosclerosis or cerebrovascular atherosclerosis.

(99) A method of identifying a patient at high risk for
30 transient ischemic attacks or stroke by determining the degree of active atherosclerosis in a patient comprising carrying out the method of embodiment 96.

(100) A method of identifying a patient at high risk for
35 transient ischemic attacks or stroke by determining the

degree of active atherosclerosis in a patient comprising carrying out the method of embodiment 97.

(101) A method of identifying a patient at high risk for acute cardiac ischemia, myocardial infarction or cardiac death by determining the degree of active atherosclerosis by imaging the patient by the method of embodiment 96.

(102) A method of identifying a patient at high risk for acute cardiac ischemia, myocardial infarction or cardiac death by determining the degree of active atherosclerosis by imaging the patient by the method of embodiment 97.

(103) A method of simultaneous imaging of cardiac perfusion and extracellular matrix degradation in a patient comprising the steps of:

- a) administering a diagnostic agent according to embodiment 1, wherein the diagnostic metal is a gamma-emitting radioisotope; and
- b) administering a cardiac perfusion compound, wherein the compound is radiolabeled with a gamma-emitting radioisotope which exhibits a gamma emission energy that is spectrally separable from the gamma emission energy of the diagnostic metal conjugated to the targeting moiety in step (a); and
- c) acquiring, by a diagnostic imaging technique, simultaneous images of the sites of concentration of the spectrally separable gamma-emission energies of the compounds administered in steps (a) and (b).

The present invention may be embodied in other specific forms without departing from the spirit or essential attributes thereof.

DEFINITIONS

The compounds herein described may have asymmetric centers. Unless otherwise indicated, all chiral, diastereomeric and

racemic forms are included in the present invention. Many geometric isomers of olefins, C=N double bonds, and the like can also be present in the compounds described herein, and all such stable isomers are contemplated in the present invention. It
5 will be appreciated that compounds of the present invention contain asymmetrically substituted carbon atoms, and may be isolated in optically active or racemic forms. It is well known in the art how to prepare optically active forms, such as by resolution of racemic forms or by synthesis from optically
10 active starting materials. Two distinct isomers (cis and trans) of the peptide bond are known to occur; both can also be present in the compounds described herein, and all such stable isomers are contemplated in the present invention. The D and L-isomers of a particular amino acid are designated herein using the
15 conventional 3-letter abbreviation of the amino acid, as indicated by the following examples: D-Leu, or L-Leu.

When any variable occurs more than one time in any substituent or in any formula, its definition on each occurrence is independent of its definition at every other occurrence.
20 Thus, for example, if a group is shown to be substituted with 0-2 R⁵², then said group may optionally be substituted with up to two R⁵², and R⁵² at each occurrence is selected independently from the defined list of possible R⁵². Also, by way of example, for the group -N(R⁵³)₂, each of the two R⁵³ substituents on N is
25 independently selected from the defined list of possible R⁵³. Combinations of substituents and/or variables are permissible only if such combinations result in stable compounds. When a bond to a substituent is shown to cross the bond connecting two atoms in a ring, then such substituent may be bonded to any atom
30 on the ring.

The term "metallopharmaceutical" means a pharmaceutical comprising a metal. The metal is the cause of the imageable signal in diagnostic applications and the source of the cytotoxic radiation in radiotherapeutic applications.

Radiopharmaceuticals are metallopharmaceuticals in which the metal is a radioisotope.

By "reagent" is meant a compound of this invention capable of direct transformation into a metallopharmaceutical of this invention. Reagents may be utilized directly for the preparation of the metallopharmaceuticals of this invention or may be a component in a kit of this invention.

The term "inhibitor" means a compound of this invention that inhibits the function of a matrix metalloproteinase.

By "stable compound" or "stable structure" is meant herein a compound that is sufficiently robust to survive isolation to a useful degree of purity from a reaction mixture, and formulation into an efficacious pharmaceutical agent.

The term "substituted", as used herein, means that one or more hydrogens on the designated atom or group is replaced with a selection from the indicated group, provided that the designated atom's or group's normal valency is not exceeded, and that the substitution results in a stable compound. When a substituent is keto (i.e., =O), then 2 hydrogens on the atom are replaced.

The term "bond", as used herein, means either a single or double bond.

The term "salt", as used herein, is used as defined in the CRC Handbook of Chemistry and Physics, 65th Edition, CRC Press, Boca Raton, Fla, 1984, as any substance which yields ions, other than hydrogen or hydroxyl ions. As used herein, "pharmaceutically acceptable salts" refer to derivatives of the disclosed compounds modified by making acid or base salts. Examples of pharmaceutically acceptable salts include, but are not limited to, mineral or organic acid salts of basic residues such as amines; alkali or organic salts of acidic residues such as carboxylic acids; and the like.

The phrase "pharmaceutically acceptable" is employed herein to refer to those compounds, materials, compositions, and/or dosage forms which are, within the scope of sound medical judgment, suitable for use in contact with the tissues of human

beings and animals without excessive toxicity, irritation, allergic response, or other problem or complication, commensurate with a reasonable benefit/risk ratio.

As used herein, "pharmaceutically acceptable salts" refer to derivatives of the disclosed compounds wherein the parent compound is modified by making acid or base salts thereof. Examples of pharmaceutically acceptable salts include, but are not limited to, mineral or organic acid salts of basic residues such as amines; alkali or organic salts of acidic residues such as carboxylic acids; and the like. The pharmaceutically acceptable salts include the conventional non-toxic salts or the quaternary ammonium salts of the parent compound formed, for example, from non-toxic inorganic or organic acids. For example, such conventional non-toxic salts include those derived from inorganic acids such as hydrochloric, hydrobromic, sulfuric, sulfamic, phosphoric, nitric and the like; and the salts prepared from organic acids such as acetic, propionic, succinic, glycolic, stearic, lactic, tartaric, citric, ascorbic, pamoic, maleic, hydroxymaleic, phenylacetic, glutamic, benzoic, salicylic, sulfanilic, 2-acetoxybenzoic, fumaric, toluenesulfonic, methanesulfonic, ethane disulfonic, oxalic, isethionic, and the like.

The pharmaceutically acceptable salts of the present invention can be synthesized from the parent compound which contains a basic or acidic moiety by conventional chemical methods. Generally, such salts can be prepared by reacting the free acid or base forms of these compounds with a stoichiometric amount of the appropriate base or acid in water or in an organic solvent, or in a mixture of the two; generally, nonaqueous media like ether, ethyl acetate, ethanol, isopropanol, or acetonitrile are preferred. Lists of suitable salts are found in *Remington's Pharmaceutical Sciences*, 17th ed., Mack Publishing Company, Easton, PA, 1985, p. 1418, the disclosure of which is hereby incorporated by reference.

As used herein, "alkyl" is intended to include both branched and straight-chain saturated aliphatic hydrocarbon

groups having the specified number of carbon atoms, examples of which include, but are not limited to, methyl, ethyl, n-propyl, i-propyl, n-butyl, i-butyl, sec-butyl, t-butyl, pentyl, hexyl, heptyl, octyl, nonyl, and decyl; "cycloalkyl" or "carbocycle" is intended to include saturated and partially unsaturated ring groups, including mono-, bi- or poly-cyclic ring systems, such as cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclooctyl and adamantyl; "bicycloalkyl" or "bicyclic" is intended to include saturated bicyclic ring groups such as [3.3.0]bicyclooctane, [4.3.0]bicyclononane, [4.4.0]bicyclodecane (decalin), [2.2.2]bicyclooctane, and so forth.

As used herein, the term "alkene" or "alkenyl" is intended to include hydrocarbon chains having the specified number of carbon atoms of either a straight or branched configuration and one or more unsaturated carbon-carbon bonds which may occur in any stable point along the chain, such as ethenyl, propenyl, and the like.

As used herein, the term "alkyne" or "alkynyl" is intended to include hydrocarbon chains having the specified number of carbon atoms of either a straight or branched configuration and one or more unsaturated carbon-carbon triple bonds which may occur in any stable point along the chain, such as propargyl, and the like.

As used herein, "aryl" or "aromatic residue" is intended to mean phenyl or naphthyl, which when substituted, the substitution can be at any position.

As used herein, the term "heterocycle" or "heterocyclic system" is intended to mean a stable 5- to 7- membered monocyclic or bicyclic or 7- to 10-membered bicyclic heterocyclic ring which is saturated partially unsaturated or unsaturated (aromatic), and which consists of carbon atoms and from 1 to 4 heteroatoms independently selected from the group consisting of N, O and S and including any bicyclic group in which any of the above-defined heterocyclic rings is fused to a benzene ring. The nitrogen and sulfur heteroatoms may

optionally be oxidized. The heterocyclic ring may be attached to its pendant group at any heteroatom or carbon atom which results in a stable structure. The heterocyclic rings described herein may be substituted on carbon or on a nitrogen atom if the resulting compound is stable. If specifically noted, a nitrogen in the heterocycle may optionally be quaternized. It is preferred that when the total number of S and O atoms in the heterocycle exceeds 1, then these heteroatoms are not adjacent to one another. It is preferred that the total number of S and O atoms in the heterocycle is not more than 1. As used herein, the term "aromatic heterocyclic system" is intended to mean a stable 5- to 7- membered monocyclic or bicyclic or 7- to 10-membered bicyclic heterocyclic aromatic ring which consists of carbon atoms and from 1 to 4 heteroatoms independently selected from the group consisting of N, O and S. It is preferred that the total number of S and O atoms in the aromatic heterocycle is not more than 1.

Examples of heterocycles include, but are not limited to, 1H-indazole, 2-pyrrolidonyl, 2H,6H-1,5,2-dithiazinyl, 2H-pyrrolyl, 3H-indolyl, 4-piperidonyl, 4aH-carbazole, 4H-quinoliziny, 6H-1,2,5-thiadiazinyl, acridinyl, azocinyl, benzimidazolyl, benzofuranyl, benzothiofuranyl, benzothiophenyl, benzoxazolyl, benzthiazolyl, benztriazolyl, benztetrazolyl, benzisoxazolyl, benzisothiazolyl, benzimidazalonyl, carbazolyl, 4aH-carbazolyl, \square -carbolinyl, chromanyl, chromenyl, cinnolinyl, decahydroquinolinyl, 2H,6H-1,5,2-dithiazinyl, dihydrofuro[2,3-b]tetrahydrofuran, furanyl, furazanyl, imidazolidinyl, imidazolyl, imidazolyl, 1H-indazolyl, indolenyl, indolinyl, indoliziny, indolyl, isobenzofuranyl, isochromanyl, isoindazolyl, isoindolinyl, isoindolyl, isoquinolinyl, isothiazolyl, isoxazolyl, morpholinyl, naphthyridinyl, octahydroisoquinolinyl, oxadiazolyl, 1,2,3-oxadiazolyl, 1,2,4-oxadiazolyl, 1,2,5-oxadiazolyl, 1,3,4-oxadiazolyl, oxazolidinyl., oxazolyl, oxazolidinylperimidinyl, phenanthridinyl, phenanthrolinyl, phenarsazinyl, phenazinyl, phenothiazinyl, phenoxathiinyl,

phenoxazinyl, phthalazinyl, piperazinyl, piperidinyl,
pteridinyl, piperidonyl, 4-piperidonyl, pteridinyl, purinyl,
pyranyl, pyrazinyl, pyrazolidinyl, pyrazolinyl, pyrazolyl,
pyridazinyl, pyridooxazole, pyridoimidazole, pyridothiazole,
5 pyridinyl, pyridyl, pyrimidinyl, pyrrolidinyl, pyrrolinyl,
pyrrolyl, quinazolinyl, quinolinyl, 4*H*-quinolizinyll,
quinoxalinyl, quinuclidinyl, carbolinyl, tetrahydrofuranyl,
tetrahydroisoquinolinyl, tetrahydroquinolinyl,
6*H*-1,2,5-thiadiazinyl, 1,2,3-thiadiazolyl, 1,2,4-thiadiazolyl,
10 1,2,5-thiadiazolyl, 1,3,4-thiadiazolyl, thianthrenyl, thiazolyl,
thienyl, thienothiazolyl, thienooxazolyl, thienoimidazolyl,
thiophenyl, triazinyl, 1,2,3-triazolyl, 1,2,4-triazolyl,
1,2,5-triazolyl, 1,3,4-triazolyl, xanthenyl. Preferred
heterocycles include, but are not limited to, pyridinyl,
15 furanyl, thienyl, pyrrolyl, pyrazolyl, imidazolyl, indolyl,
benzimidazolyl, 1*H*-indazolyl, oxazolidinyl, benzotriazolyl,
benzisoxazolyl, oxindolyl, benzoxazolinyl, or isatinoyl. Also
included are fused ring and spiro compounds containing, for
example, the above heterocycles.

20 As used herein, the term "alkaryl" means an aryl group
bearing an alkyl group of 1-10 carbon atoms; the term "aralkyl"
means an alkyl group of 1-10 carbon atoms bearing an aryl group;
the term "arylalkaryl" means an aryl group bearing an alkyl
group of 1-10 carbon atoms bearing an aryl group; and the term
25 "heterocycloalkyl" means an alkyl group of 1-10 carbon atoms
bearing a heterocycle.

A "polyalkylene glycol" is a polyethylene glycol,
polypropylene glycol or polybutylene glycol having a molecular
weight of less than about 5000, terminating in either a hydroxy
30 or alkyl ether moiety.

A "carbohydrate" is a polyhydroxy aldehyde, ketone, alcohol
or acid, or derivatives thereof, including polymers thereof
having polymeric linkages of the acetal type.

A "cyclodextrin" is a cyclic oligosaccharide. Examples of
35 cyclodextrins include, but are not limited to, α -cyclodextrin,
hydroxyethyl- α -cyclodextrin, hydroxypropyl- α -cyclodextrin,

□-cyclodextrin, hydroxypropyl-□-cyclodextrin, carboxymethyl-□-cyclodextrin, dihydroxypropyl-□-cyclodextrin, hydroxyethyl-□-cyclodextrin, 2,6 di-O-methyl-□-cyclodextrin, sulfated-□-cyclodextrin, □-cyclodextrin, 5 hydroxypropyl-□-cyclodextrin, dihydroxypropyl-□-cyclodextrin, hydroxyethyl-□-cyclodextrin, and sulfated □-cyclodextrin.

As used herein, the term "polycarboxyalkyl" means an alkyl group having between two and about 100 carbon atoms and a plurality of carboxyl substituents; and the term "polyazaalkyl" 10 means a linear or branched alkyl group having between two and about 100 carbon atoms, interrupted by or substituted with a plurality of amine groups.

A "reducing agent" is a compound that reacts with a radionuclide, which is typically obtained as a relatively 15 unreactive, high oxidation state compound, to lower its oxidation state by transferring electron(s) to the radionuclide, thereby making it more reactive. Reducing agents useful in the preparation of radiopharmaceuticals and in diagnostic kits useful for the preparation of said radiopharmaceuticals include 20 but are not limited to stannous chloride, stannous fluoride, formamidine sulfinic acid, ascorbic acid, cysteine, phosphines, and cuprous or ferrous salts. Other reducing agents are described in Brodack et. al., PCT Application 94/22496, which is incorporated herein by reference.

25 A "transfer ligand" is a ligand that forms an intermediate complex with a metal ion that is stable enough to prevent unwanted side-reactions but labile enough to be converted to a metallopharmaceutical. The formation of the intermediate complex is kinetically favored while the formation of the 30 metallopharmaceutical is thermodynamically favored. Transfer ligands useful in the preparation of metallopharmaceuticals and in diagnostic kits useful for the preparation of diagnostic radiopharmaceuticals include but are not limited to gluconate, glucoheptonate, mannitol, glucarate, 35 N,N,N',N'-ethylenediaminetetraacetic acid, pyrophosphate and

methylenediphosphonate. In general, transfer ligands are comprised of oxygen or nitrogen donor atoms.

The term "donor atom" refers to the atom directly attached to a metal by a chemical bond.

5 "Ancillary" or "co-ligands" are ligands that are incorporated into a radiopharmaceutical during its synthesis. They serve to complete the coordination sphere of the radionuclide together with the chelator or radionuclide bonding unit of the reagent. For radiopharmaceuticals comprised of a
10 binary ligand system, the radionuclide coordination sphere is composed of one or more chelators or bonding units from one or more reagents and one or more ancillary or co-ligands, provided that there are a total of two types of ligands, chelators or bonding units. For example, a radiopharmaceutical comprised of
15 one chelator or bonding unit from one reagent and two of the same ancillary or co-ligands and a radiopharmaceutical comprised of two chelators or bonding units from one or two reagents and one ancillary or co-ligand are both considered to be comprised of binary ligand systems. For radiopharmaceuticals comprised of
20 a ternary ligand system, the radionuclide coordination sphere is composed of one or more chelators or bonding units from one or more reagents and one or more of two different types of ancillary or co-ligands, provided that there are a total of three types of ligands, chelators or bonding units. For
25 example, a radiopharmaceutical comprised of one chelator or bonding unit from one reagent and two different ancillary or co-ligands is considered to be comprised of a ternary ligand system.

Ancillary or co-ligands useful in the preparation of
30 radiopharmaceuticals and in diagnostic kits useful for the preparation of said radiopharmaceuticals are comprised of one or more oxygen, nitrogen, carbon, sulfur, phosphorus, arsenic, selenium, and tellurium donor atoms. A ligand can be a transfer ligand in the synthesis of a radiopharmaceutical and also serve
35 as an ancillary or co-ligand in another radiopharmaceutical. Whether a ligand is termed a transfer or ancillary or co-ligand

depends on whether the ligand remains in the radionuclide coordination sphere in the radiopharmaceutical, which is determined by the coordination chemistry of the radionuclide and the chelator or bonding unit of the reagent or reagents.

5 A "chelator" or "bonding unit" is the moiety or group on a reagent that binds to a metal ion through the formation of chemical bonds with one or more donor atoms.

 The term "binding site" means the site in vivo or in vitro that binds a biologically active molecule.

10 A "diagnostic kit" or "kit" comprises a collection of components, termed the formulation, in one or more vials which are used by the practicing end user in a clinical or pharmacy setting to synthesize diagnostic radiopharmaceuticals. The kit provides all the requisite components to synthesize and use the
15 diagnostic radiopharmaceutical except those that are commonly available to the practicing end user, such as water or saline for injection, a solution of the radionuclide, equipment for heating the kit during the synthesis of the radiopharmaceutical, if required, equipment necessary for administering the
20 radiopharmaceutical to the patient such as syringes and shielding, and imaging equipment.

 X-ray contrast agent pharmaceuticals, ultrasound contrast agent pharmaceuticals and metallopharmaceuticals for magnetic resonance imaging contrast are provided to the end user in their
25 final form in a formulation contained typically in one vial, as either a lyophilized solid or an aqueous solution. The end user reconstitutes the lyophilized with water or saline and withdraws the patient dose or just withdraws the dose from the aqueous solution formulation as provided.

30 A "lyophilization aid" is a component that has favorable physical properties for lyophilization, such as the glass transition temperature, and is added to the formulation to improve the physical properties of the combination of all the components of the formulation for lyophilization.

35 A "stabilization aid" is a component that is added to the metallopharmaceutical or to the diagnostic kit either to

stabilize the metallopharmaceutical or to prolong the shelf-life of the kit before it must be used. Stabilization aids can be antioxidants, reducing agents or radical scavengers and can provide improved stability by reacting preferentially with species that degrade other components or the metallopharmaceutical.

A "solubilization aid" is a component that improves the solubility of one or more other components in the medium required for the formulation.

A "bacteriostat" is a component that inhibits the growth of bacteria in a formulation either during its storage before use or after a diagnostic kit is used to synthesize a radiopharmaceutical.

The following abbreviations are used herein:

Acm	acetamidomethyl
b-Ala, beta-Ala	
or bAla	3-aminopropionic acid
ATA	2-aminothiazole-5-acetic acid or 2-aminothiazole-5-acetyl group
Boc	t-butyloxycarbonyl
CBZ, Cbz or Z	Carbobenzyloxy
Cit	citrulline
Dap	2,3-diaminopropionic acid
DCC	dicyclohexylcarbodiimide
DIEA	diisopropylethylamine
DMAP	4-dimethylaminopyridine
EOE	ethoxyethyl
HBTU	2-(1H-Benzotriazol-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate
hynic	boc-hydrazinonicotinyl group or 2-[[[5-[carbonyl]-2-pyridinyl]hydrazono]methyl]-benzenesulfonic acid,
NMeArg or MeArg	a-N-methyl arginine
NMeAsp	a-N-methyl aspartic acid
NMM	N-methylmorpholine

OcHex	O-cyclohexyl
OBzl	O-benzyl
oSu	O-succinimidyl
TBTU	2-(1H-Benzotriazol-1-yl)-1,1,3,3-
5	tetramethyluronium tetrafluoroborate
THF	tetrahydrofuranyl
THP	tetrahydropyranyl
Tos	tosyl
Tr	trityl

10

The following conventional three-letter amino acid abbreviations are used herein; the conventional one-letter amino acid abbreviations are NOT used herein:

15	Ala	=	alanine
	Arg	=	arginine
	Asn	=	asparagine
	Asp	=	aspartic acid
	Cys	=	cysteine
20	Gln	=	glutamine
	Glu	=	glutamic acid
	Gly	=	glycine
	His	=	histidine
	Ile	=	isoleucine
25	Leu	=	leucine
	Lys	=	lysine
	Met	=	methionine
	Nle	=	norleucine
	Orn	=	ornithine
30	Phe	=	phenylalanine
	Phg	=	phenylglycine
	Pro	=	proline
	Sar	=	sarcosine
	Ser	=	serine
35	Thr	=	threonine
	Trp	=	tryptophan

Tyr = tyrosine
Val = valine

5 The ultrasound contrast agents of the present invention
comprise a plurality of matrix metalloproteinase inhibiting
moieties attached to or incorporated into a microbubble of a
biocompatible gas, a liquid carrier, and a surfactant
microsphere, further comprising an optional linking moiety, L_n ,
between the targeting moieties and the microbubble. In this
10 context, the term liquid carrier means aqueous solution and the
term surfactant means any amphiphilic material which produces a
reduction in interfacial tension in a solution. A list of
suitable surfactants for forming surfactant microspheres is
disclosed in EP0727225A2, herein incorporated by reference. The
15 term surfactant microsphere includes nanospheres, liposomes,
vesicles and the like. The biocompatible gas can be air, or a
fluorocarbon, such as a C₃-C₅ perfluoroalkane, which provides
the difference in echogenicity and thus the contrast in
ultrasound imaging. The gas is encapsulated or contained in the
20 microsphere to which is attached the biodirecting group,
optionally via a linking group. The attachment can be covalent,
ionic or by van der Waals forces. Specific examples of such
contrast agents include lipid encapsulated perfluorocarbons with
a plurality of MMP inhibiting compounds.

25 X-ray contrast agents of the present invention are
comprised of one or more matrix metalloproteinase inhibiting
targeting moieties attached to one or more X-ray absorbing or
"heavy" atoms of atomic number 20 or greater, further comprising
an optional linking moiety, L_n , between the targeting moieties
30 and the X-ray absorbing atoms. The frequently used heavy atom in
X-ray contrast agents is iodine. Recently, X-ray contrast agents
comprised of metal chelates (Wallace, R., U.S. 5,417,959) and
polychelates comprised of a plurality of metal ions (Love, D.,
U.S. 5,679,810) have been disclosed. More recently, multinuclear
35 cluster complexes have been disclosed as X-ray contrast agents
(U.S. 5,804,161, PCT WO91/14460, and PCT WO 92/17215).

MRI contrast agents of the present invention are comprised of one or more matrix metalloproteinase inhibiting targeting moieties attached to one or more paramagnetic metal ions, further comprising an optional linking moiety, L_n , between the targeting moieties and the paramagnetic metal ions. The paramagnetic metal ions are present in the form of metal complexes or metal oxide particles. U.S. 5,412,148, and 5,760,191, describe examples of chelators for paramagnetic metal ions for use in MRI contrast agents. U.S. 5,801,228, U.S. 5,567,411, and U.S. 5,281,704, describe examples of polychelants useful for complexing more than one paramagnetic metal ion for use in MRI contrast agents. U.S. 5,520,904, describes particulate compositions comprised of paramagnetic metal ions for use as MRI contrast agents.

Matrix metalloproteinases (MMPs) are a family of structurally related zinc-containing enzymes that mediate the integrity of extracellular matrix (Whittaker, M. et al, Chem. Rev., 1999, 99, 2735-2776). They are excreted by a variety of connective tissue and pro-inflammatory cells, such as, fibroblasts, osteoblasts, macrophages, neutrophils, lymphocytes and endothelial cells. There is now a body of evidence that matrix metalloproteinases (MMPs) are important in the uncontrolled breakdown of connective tissue, including proteoglycan and collagen, leading to resorption of the extracellular matrix. This is a feature of a number of cardiovascular pathological conditions, such as atherosclerosis, heart failure, restenosis and reperfusion injury. Normally these catabolic enzymes are tightly regulated at the level of their synthesis, as well as, at their level of extracellular activity through the action of specific inhibitors, such as alpha-2-macroglobulins and TIMP (tissue inhibitor of metalloproteinase), which form inactive complexes with the MMPs. Therefore, extracellular matrix degradation and remodeling are regulated by the relative expression of TIMPs and MMPs. The MMPs are classified into several families based on their domain structure: matrilysin (minimal domain, MMP-7), collagenase


(hemopexin domain, MMP-1; MMP-8, MMP-13), gelatinase (fibronectin domain, MMP-2, MMP-9), stromelysin (hemopexin domain, MMP-3, MMP-10, MMP-11), metalloelastase (MMP-12). In addition, the transmembrane domain family (MT-MMPs) has been recently discovered and comprises MMP-14 through MMP-17.

MMP proteolytic activity and extracellular matrix degradation is dependent on the comparative balance between MMPs and TIMPs. TIMPs and synthetic small molecules or matrix metalloproteinase inhibitors have therapeutic potential for diseases involving elevated levels of MMP activity (Whittaker, M. et al, Chem. Rev., 1999, 99, 2735-2776; Babine, R. E. et al, Chem. Rev., 1997, 97, 1359; De, B. et al, Ann. N.Y. Acad. Sci., 1999, 878, 40-60; Summers, J.B. et al, Annual Reports in Med. Chem., 1998, 33, 131).

A functional group, such as -CONH-OH, -COOH, or -SH, is necessary for a molecule to be an effective inhibitor of MMPs. This functional group is involved in the chelation of the active site zinc ion, and is commonly referred to as the zinc binding group or ZBG. The hydroxamate, for example, is a bidentate ligand for zinc.


The pharmaceuticals of the present invention have the formulae, $(Q)d-L_n-(Ch-X)$, $(Q)d-L_n-(Ch-X^1)d'$, $(Q)d-L_n-(X^2)d''$, and $(Q)d-L_n-(X^3)$, wherein Q represents a compound that inhibits a matrix metalloproteinase, d is 1-10, $d' = 1-100$, L_n represents an optional linking group, Ch represents a metal chelator or bonding moiety, X represents a radioisotope, X^1 represents paramagnetic metal ion, X^2 represents a paramagnetic metal ion or heavy atom containing insoluble solid particle, d'' is 1-100, and X^3 represents a surfactant microsphere of an echogenic gas.

One class of compounds of the present invention is comprised of one or more inhibitors, Q, which are succinyl hydroxamates. A generic structure of succinyl hydroxamate is shown below (1).




(1)

The ethylene spacer between the ZBG (-CONH-OH) and the succinyl
 5 amide is essential for potent activity. Substitution at P₁ tends
 to confer broad-spectrum activity on the MMPis. Substituents at
 this position, in general, tend to point away from the enzyme.
 Moieties capable of hydrogen bonding and lipophilic substituents
 at the P₁ position α to the hydroxamate (Johnson, W.H. et al, J.
 10 Enz. Inhib., 1987, 2, 1) tend to enhance activity (2).
 Incorporation of a hydroxyl group (Beckett, P.R., et al, Drug
 Discovery Today, 1996, 1, 16) at that position improves oral
 activity in some case (3).





(2)



(3)


Substituents at the P₁' position on the succinyl hydroxamates tend
 to impart selectivity to the MMPis. The S₁' pocket is deep for
 MMP-2, MMP-3, MMP-8 and MMP-9 and occluded (short) for MMP-1 and
 MMP-7. A long alkyl substituent at the P₁' position, for
 20 example, imparts selectivity (Miller, A. et al, Bioorg. Med.
 Chem. Lett., 1997, 7, 193) for MMP-2 over MMP-1 and MMP-3 (4 and
 5).



(4)

(5)


Substituents at the P_2' position also point away from the enzyme. The P_1 and the P_2' positions can be linked (Xue, C-B. et al, J. Med. Chem., 1998, 41, 1745; Steinman, D.H. et al, Bioorg. Med. Chem. Lett., 1998, 8, 2087) to form a macrocycle (6). Compounds such as (6) also exhibit nanomolar activity.



(6)

The nature of the macrocycle also imparts some selective inhibition among the MMPs. The P_2' and the P_3' positions may be cyclized to form lactams. The size of the lactam governs the selectivity.


The P_3' position is a relatively open area in the succinyl hydroxamates, and a wide range of substituents (for example (7)) may be introduced (Sheppard, G.S. et al, Bioorg. Med. Chem. Lett., 1998, 8, 3251) at this position. This position also offers the flexibility of attaching the optional linker, L_n , the chelator(s), Ch , for the imageable moieties X and X^1 , and the imageable moieties, X^2 and X^3 .



(7)


Other succinyl hydroxamates with modified P_2' and P_3' positions, such as (8) also have shown potent inhibition of MMPs. Those

compounds and syntheses of them are further described in the following patent applications which are hereby incorporated by reference into this patent application: U.S. Patent Application Serial Nos. 09/165,747, 08/743,439, 09/134,484, 09/247,675, 5 09/335,086, 09/312,066, 09/311,168, 60/127,594, and 60/127,635.




(8)

10 Another class of compounds of the present invention is comprised of one or more inhibitors, Q, which are sulfonamide hydroxamates, such as (9) and (10). Modification of the isopropyl substituent in (10) results in deep pocket MMP selectivity, for example MMP-2 vs MMP-1 (Santos, O. et al., J. Clin. Exp. Metastasis, 1997, 15, 499; MacPherson, L.J. et al, J. 15 Med. Chem., 1997, 40, 2525).




(9)




(10)

20 Additional examples of inhibitors, Q, include the derivatized 'alanine' hydroxamates, such as compounds (11) and (12), which show selectivity for MMP-2 and MMP-9 over the other MMPs. Those compounds and syntheses of them are further described in the following patent applications which are hereby incorporated by

reference into this patent application: U.S. Patent Application Serial Nos. 09/165,747, 08/743,439, 09/134,484, 09/247,675, 09/335,086, 09/312,066, 09/311,168, 60/127,594, and 60/127,635.




(11)




(12)

compound (13), which shows selectivity for MMP-2 and MMP-9, and in which the alpha position has a quaternary carbon and the molecule does not contain any stereo centers (Lovejoy, B. et al., Nature Struct. Biol., 1999, 6, 217);




(13)

compound (14), which uses a carboxylic acid as the ZBG, and exhibits significant selectivity for MMP-2 (vs MMP-1), when X = butyl vs X = H (Sahoo, S.P. et al, Bioorg. Med. Chem. Lett., 1995, 5, 2441); and



(14)

succinyl thiols such as (15) (Levin, J.I. et al, Bioorg. Med. Chem. Lett., 1998, 8, 1163), in which the P₃' position may be
 5 utilized to attach the optional linker, L_n, the chelator(s), C_n, for the imageable moieties X and X¹, and the imageable moieties, X² and X³.



(15)

10

Preferred pharmaceuticals of the present invention are comprised of inhibitors, Q, which exhibit selectivity for MMP-1,
 15 MMP-2, MMP-3, MMP-9, or MMP-14 alone or in combination over the other MMPs. Examples of preferred moieties, Q, include compounds 4, 5, 6, 8, 9, 10, 11, 12, and 13.

Most preferred are comprised of inhibitors, Q, which exhibit selectivity for MMP-2, MMP-9, or MMP-14 alone or in
 20 combination over the other MMPs. Examples of the most preferred moieties, Q, include compounds 6, 8, 11, and 12.

The pharmaceuticals of the present invention can be synthesized by several approaches. One approach involves the synthesis of the targeting MMP inhibiting moiety, Q, and direct
 25 attachment of one or more moieties, Q, to one or more metal

chelators or bonding moieties, C_h , or to a paramagnetic metal ion or heavy atom containing solid particle, or to an echogenic gas microbubble. Another approach involves the attachment of one or more moieties, Q , to the linking group, L_n , which is then
5 attached to one or more metal chelators or bonding moieties, C_h , or to a paramagnetic metal ion or heavy atom containing solid particle, or to an echogenic gas microbubble. Another approach, useful in the synthesis of pharmaceuticals wherein d is 1, involves the synthesis of the moiety, $Q-L_n$, together, by
10 incorporating residue bearing L_n into the synthesis of the MMP inhibitor, Q . The resulting moiety, $Q-L_n$, is then attached to one or more metal chelators or bonding moieties, C_h , or to a paramagnetic metal ion or heavy atom containing solid particle, or to an echogenic gas microbubble. Another approach involves
15 the synthesis of an inhibitor, Q , bearing a fragment of the linking group, L_n , one or more of which are then attached to the remainder of the linking group and then to one or more metal chelators or bonding moieties, C_h , or to a paramagnetic metal ion or heavy atom containing solid particle, or to an echogenic
20 gas microbubble.

The MMP inhibiting moieties, Q , optionally bearing a linking group, L_n , or a fragment of the linking group, can be synthesized using standard synthetic methods known to those skilled in the art. Preferred methods include but are not
25 limited to those methods described below.

Generally, peptides, polypeptides and peptidomimetics are elongated by deprotecting the alpha-amine of the C-terminal residue and coupling the next suitably protected amino acid through a peptide linkage using the methods described. This
30 deprotection and coupling procedure is repeated until the desired sequence is obtained. This coupling can be performed with the constituent amino acids in a stepwise fashion, or condensation of fragments (two to several amino acids), or combination of both processes, or by solid phase peptide
35 synthesis according to the method originally described by

Merrifield, J. Am. Chem. Soc., 85, 2149-2154 (1963), the disclosure of which is hereby incorporated by reference.

The peptides, polypeptides and peptidomimetics may also be synthesized using automated synthesizing equipment. In addition
5 to the foregoing, procedures for peptide, polypeptide and peptidomimetic synthesis are described in Stewart and Young, "Solid Phase Peptide Synthesis", 2nd ed, Pierce Chemical Co., Rockford, IL (1984); Gross, Meienhofer, Udenfriend, Eds., "The Peptides: Analysis, Synthesis, Biology, Vol. 1, 2, 3, 5, and 9,
10 Academic Press, New York, (1980-1987); Bodanszky, "Peptide Chemistry: A Practical Textbook", Springer-Verlag, New York (1988); and Bodanszky et al. "The Practice of Peptide Synthesis" Springer-Verlag, New York (1984), the disclosures of which are hereby incorporated by reference.

15 The coupling between two amino acid derivatives, an amino acid and a peptide, polypeptide or peptidomimetic, two peptide, polypeptide or peptidomimetic fragments, or the cyclization of a peptide, polypeptide or peptidomimetic can be carried out using standard coupling procedures such as the azide method, mixed
20 carbonic acid anhydride (isobutyl chloroformate) method, carbodiimide (dicyclohexylcarbodiimide, diisopropylcarbodiimide, or water-soluble carbodiimides) method, active ester (p-nitrophenyl ester, N-hydroxysuccinic imido ester) method, Woodward reagent K method, carbonyldiimidazole method,
25 phosphorus reagents such as BOP-Cl, or oxidation-reduction method. Some of these methods (especially the carbodiimide) can be enhanced by the addition of 1-hydroxybenzotriazole. These coupling reactions may be performed in either solution (liquid phase) or solid phase.

30 The functional groups of the constituent amino acids or amino acid mimetics must be protected during the coupling reactions to avoid undesired bonds being formed. The protecting groups that can be used are listed in Greene, "Protective Groups in Organic Synthesis" John Wiley & Sons, New York (1981) and
35 "The Peptides: Analysis, Synthesis, Biology, Vol. 3, Academic

Press, New York (1981), the disclosure of which is hereby incorporated by reference.

The alpha-carboxyl group of the C-terminal residue is usually protected by an ester that can be cleaved to give the carboxylic acid. These protecting groups include: 1) alkyl esters such as methyl and t-butyl, 2) aryl esters such as benzyl and substituted benzyl, or 3) esters which can be cleaved by mild base treatment or mild reductive means such as trichloroethyl and phenacyl esters. In the solid phase case, the C-terminal amino acid is attached to an insoluble carrier (usually polystyrene). These insoluble carriers contain a group which will react with the carboxyl group to form a bond which is stable to the elongation conditions but readily cleaved later. Examples of which are: oxime resin (DeGrado and Kaiser (1980) *J. Org. Chem.* 45, 1295-1300) chloro or bromomethyl resin, hydroxymethyl resin, and aminomethyl resin. Many of these resins are commercially available with the desired C-terminal amino acid already incorporated.

The alpha-amino group of each amino acid must be protected. Any protecting group known in the art can be used. Examples of these are: 1) acyl types such as formyl, trifluoroacetyl, phthalyl, and p-toluenesulfonyl; 2) aromatic carbamate types such as benzyloxycarbonyl (Cbz) and substituted benzyloxycarbonyls, 1-(p-biphenyl)-1-methylethoxycarbonyl, and 9-fluorenylmethyloxycarbonyl (Fmoc); 3) aliphatic carbamate types such as tert-butyloxycarbonyl (Boc), ethoxycarbonyl, diisopropylmethoxycarbonyl, and allyloxycarbonyl; 4) cyclic alkyl carbamate types such as cyclopentyloxycarbonyl and adamantyloxycarbonyl; 5) alkyl types such as triphenylmethyl and benzyl; 6) trialkylsilane such as trimethylsilane; and 7) thiol containing types such as phenylthiocarbonyl and dithiasuccinoyl. The preferred alpha-amino protecting group is either Boc or Fmoc. Many amino acid or amino acid mimetic derivatives suitably protected for peptide synthesis are commercially available.

The alpha-amino protecting group is cleaved prior to the coupling of the next amino acid. When the Boc group is used, the methods of choice are trifluoroacetic acid, neat or in dichloromethane, or HCl in dioxane. The resulting ammonium salt
5 is then neutralized either prior to the coupling or in situ with basic solutions such as aqueous buffers, or tertiary amines in dichloromethane or dimethylformamide. When the Fmoc group is used, the reagents of choice are piperidine or substituted piperidines in dimethylformamide, but any secondary amine or
10 aqueous basic solutions can be used. The deprotection is carried out at a temperature between 0 °C and room temperature.

Any of the amino acids or amino acid mimetics bearing side chain functionalities must be protected during the preparation of the peptide using any of the above-identified groups. Those
15 skilled in the art will appreciate that the selection and use of appropriate protecting groups for these side chain functionalities will depend upon the amino acid or amino acid mimetic and presence of other protecting groups in the peptide, polypeptide or peptidomimetic. The selection of such a
20 protecting group is important in that it must not be removed during the deprotection and coupling of the alpha-amino group.

For example, when Boc is chosen for the alpha-amine protection the following protecting groups are acceptable:
p-toluenesulfonyl (tosyl) moieties and nitro for arginine;
25 benzyloxycarbonyl, substituted benzyloxycarbonyls, tosyl or trifluoroacetyl for lysine; benzyl or alkyl esters such as cyclopentyl for glutamic and aspartic acids; benzyl ethers for serine and threonine; benzyl ethers, substituted benzyl ethers or 2-bromobenzyloxycarbonyl for tyrosine; *p*-methylbenzyl,
30 *p*-methoxybenzyl, acetamidomethyl, benzyl, or *t*-butylsulfonyl for cysteine; and the indole of tryptophan can either be left unprotected or protected with a formyl group.

When Fmoc is chosen for the alpha-amine protection usually tert-butyl based protecting groups are acceptable. For
35 instance, Boc can be used for lysine, tert-butyl ether for

serine, threonine and tyrosine, and tert-butyl ester for glutamic and aspartic acids.

Once the elongation of the peptide, polypeptide or peptidomimetic, or the elongation and cyclization of a cyclic peptide or peptidomimetic is completed all of the protecting groups are removed. For the liquid phase synthesis the protecting groups are removed in whatever manner as dictated by the choice of protecting groups. These procedures are well known to those skilled in the art.

When a solid phase synthesis is used to synthesize a cyclic peptide or peptidomimetic, the peptide or peptidomimetic should be removed from the resin without simultaneously removing protecting groups from functional groups that might interfere with the cyclization process. Thus, if the peptide or peptidomimetic is to be cyclized in solution, the cleavage conditions need to be chosen such that a free α -carboxylate and a free α -amino group are generated without simultaneously removing other protecting groups. Alternatively, the peptide or peptidomimetic may be removed from the resin by hydrazinolysis, and then coupled by the azide method. Another very convenient method involves the synthesis of peptides or peptidomimetics on an oxime resin, followed by intramolecular nucleophilic displacement from the resin, which generates a cyclic peptide or peptidomimetic (Osapay, Profit, and Taylor (1990) *Tetrahedron Letters* 43, 6121-6124). When the oxime resin is employed, the Boc protection scheme is generally chosen. Then, the preferred method for removing side chain protecting groups generally involves treatment with anhydrous HF containing additives such as dimethyl sulfide, anisole, thioanisole, or p-cresol at 0 °C. The cleavage of the peptide or peptidomimetic can also be accomplished by other acid reagents such as trifluoromethanesulfonic acid/trifluoroacetic acid mixtures.

Unusual amino acids used in this invention can be synthesized by standard methods familiar to those skilled in the art ("The Peptides: Analysis, Synthesis, Biology, Vol. 5, pp. 342-449, Academic Press, New York (1981)). N-Alkyl amino acids

can be prepared using procedures described in previously (Cheung et al., (1977) *Can. J. Chem.* 55, 906; Freidinger et al., (1982) *J. Org. Chem.* 48, 77 (1982)), which are incorporated herein by reference.


5 The attachment of linking groups, L_n , to the MMP inhibitors, Q; chelators or bonding units, Ch , to the inhibitors, Q, or to the linking groups, L_n ; and inhibitors bearing a fragment of the linking group to the remainder of the linking group, in combination forming the moiety, $(Q)_d-L_n$, and
10 then to the moiety Ch ; can all be performed by standard techniques. These include, but are not limited to, amidation, esterification, alkylation, and the formation of ureas or thioureas. Procedures for performing these attachments can be found in Brinkley, M., *Bioconjugate Chemistry* 1992, 3(1), which
15 is incorporated herein by reference.

A number of methods can be used to attach the MMP inhibitors, Q, to paramagnetic metal ion or heavy atom containing solid particles, X^2 , by one of skill in the art of the surface modification of solid particles. In general, the
20 targeting moiety Q or the combination $(Q)_d-L_n$ is attached to a coupling group that react with a constituent of the surface of the solid particle. The coupling groups can be any of a number of silanes which react with surface hydroxyl groups on the solid particle surface, as described in co-pending U.S. Patent
25 Application Serial No. 09/356,178 and can also include polyphosphonates, polycarboxylates, polyphosphates or mixtures thereof which couple with the surface of the solid particles, as described in U.S. 5,520,904.

A number of reaction schemes can be used to attach the MMP
30 inhibitors, Q, to the surfactant microsphere, X^3 . These are illustrated in following reaction schemes where S_f represents a surfactant moiety that forms the surfactant microsphere.

Acylation Reaction:

35



Y is a leaving group or active ester

5 Disulfide Coupling:



Sulfonamide Coupling:

10



Reductive Amidation:



In these reaction schemes, the substituents Sf and Q can be reversed as well.

20 The linking group L_n can serve several roles. First it provides a spacing group between the metal chelator or bonding moiety, Ch , the paramagnetic metal ion or heavy atom containing solid particle, x^2 , and the surfactant microsphere, x^3 , and the one or more of the MMP inhibitors, Q, so as to minimize the

25 possibility that the moieties Ch-X , Ch-X^1 , x^2 , and x^3 , will interfere with the interaction of the recognition sequences of Q with MMPs associated with cardiovascular pathologies. The necessity of incorporating a linking group in a reagent is dependent on the identity of Q, Ch-X , Ch-X^1 , x^2 , and x^3 . If Ch-X ,

30 Ch-X^1 , x^2 , and x^3 , cannot be attached to Q without substantially diminishing its ability to inhibit MMPs, then a linking group is used. A linking group also provides a means of independently attaching multiple inhibitors, Q, to one group that is attached to Ch-X , Ch-X^1 , x^2 , or x^3 .

The linking group also provides a means of incorporating a pharmacokinetic modifier into the pharmaceuticals of the present invention. The pharmacokinetic modifier serves to direct the biodistribution of the injected pharmaceutical other than by the interaction of the targeting moieties, Q, with the MMPs expressed in the cardiovascular pathologies. A wide variety of functional groups can serve as pharmacokinetic modifiers, including, but not limited to, carbohydrates, polyalkylene glycols, peptides or other polyamino acids, and cyclodextrins. The modifiers can be used to enhance or decrease hydrophilicity and to enhance or decrease the rate of blood clearance. The modifiers can also be used to direct the route of elimination of the pharmaceuticals. Preferred pharmacokinetic modifiers are those that result in moderate to fast blood clearance and enhanced renal excretion.

The metal chelator or bonding moiety, Ch, is selected to form stable complexes with the metal ion chosen for the particular application. Chelators or bonding moieties for diagnostic radiopharmaceuticals are selected to form stable complexes with the radioisotopes that have imageable gamma ray or positron emissions, such as ^{99m}Tc , ^{95}Tc , ^{111}In , ^{62}Cu , ^{60}Cu , ^{64}Cu , ^{67}Ga , ^{68}Ga , ^{86}Y .

Chelators for technetium, copper and gallium isotopes are selected from diaminedithiols, monoamine-monoamidedithiols, triamide-monothiols, monoamine-diamide-monothiols, diaminedioximes, and hydrazines. The chelators are generally tetradentate with donor atoms selected from nitrogen, oxygen and sulfur. Preferred reagents are comprised of chelators having amine nitrogen and thiol sulfur donor atoms and hydrazine bonding units. The thiol sulfur atoms and the hydrazines may bear a protecting group which can be displaced either prior to using the reagent to synthesize a radiopharmaceutical or preferably in situ during the synthesis of the radiopharmaceutical.

Exemplary thiol protecting groups include those listed in Greene and Wuts, "Protective Groups in Organic Synthesis" John

Wiley & Sons, New York (1991), the disclosure of which is hereby incorporated by reference. Any thiol protecting group known in the art can be used. Examples of thiol protecting groups include, but are not limited to, the following: acetamidomethyl, 5 benzamidomethyl, 1-ethoxyethyl, benzoyl, and triphenylmethyl.

Exemplary protecting groups for hydrazine bonding units are hydrazones which can be aldehyde or ketone hydrazones having substituents selected from hydrogen, alkyl, aryl and heterocycle. Particularly preferred hydrazones are described in 10 co-pending U.S.S.N. 08/476,296 the disclosure of which is herein incorporated by reference in its entirety.

The hydrazine bonding unit when bound to a metal radionuclide is termed a hydrazido, or diazenido group and serves as the point of attachment of the radionuclide to the remainder of the radiopharmaceutical. A diazenido group can be 15 either terminal (only one atom of the group is bound to the radionuclide) or chelating. In order to have a chelating diazenido group at least one other atom of the group must also be bound to the radionuclide. The atoms bound to the metal are 20 termed donor atoms.

Chelators for ^{111}In and ^{86}Y are selected from cyclic and acyclic polyaminocarboxylates such as DTPA, DOTA, D03A, 2-benzyl-DOTA, alpha-(2-phenethyl)1,4,7,10-tetraazacyclododecane-1-acetic-4,7,10-tris(methylacetic)acid, 2-benzyl- 25 cyclohexyldiethylenetriaminepentaacetic acid, 2-benzyl-6-methyl-DTPA, and 6,6"-bis[N,N,N",N"-tetra(carboxymethyl)aminomethyl)-4'-(3-amino-4-methoxyphenyl)-2,2':6',2"-terpyridine. Procedures for synthesizing these chelators that are not commercially available can be found in Brechbiel, M. and Gansow, O., *J. Chem. Soc. Perkin Trans.* 1992, 1, 1175; Brechbiel, M. and Gansow, O., 30 *Bioconjugate Chem.* 1991, 2, 187; Deshpande, S., et. al., *J. Nucl. Med.* 1990, 31, 473; Kruper, J., U.S. Patent 5,064,956, and Toner, J., U.S. Patent 4,859,777, the disclosures of which are hereby incorporated by reference in their entirety.

35 The coordination sphere of metal ion includes all the ligands or groups bound to the metal. For a transition metal

radionuclide to be stable it typically has a coordination number (number of donor atoms) comprised of an integer greater than or equal to 4 and less than or equal to 8; that is there are 4 to 8 atoms bound to the metal and it is said to have a complete
5 coordination sphere. The requisite coordination number for a stable radionuclide complex is determined by the identity of the radionuclide, its oxidation state, and the type of donor atoms. If the chelator or bonding unit does not provide all of the atoms necessary to stabilize the metal radionuclide by
10 completing its coordination sphere, the coordination sphere is completed by donor atoms from other ligands, termed ancillary or co-ligands, which can also be either terminal or chelating.

A large number of ligands can serve as ancillary or co-ligands, the choice of which is determined by a variety of
15 considerations such as the ease of synthesis of the radiopharmaceutical, the chemical and physical properties of the ancillary ligand, the rate of formation, the yield, and the number of isomeric forms of the resulting radiopharmaceuticals, the ability to administer said ancillary or co-ligand to a
20 patient without adverse physiological consequences to said patient, and the compatibility of the ligand in a lyophilized kit formulation. The charge and lipophilicity of the ancillary ligand will effect the charge and lipophilicity of the radiopharmaceuticals. For example, the use of
25 4,5-dihydroxy-1,3-benzene disulfonate results in radiopharmaceuticals with an additional two anionic groups because the sulfonate groups will be anionic under physiological conditions. The use of N-alkyl substituted
3,4-hydroxypyridinones results in radiopharmaceuticals with
30 varying degrees of lipophilicity depending on the size of the alkyl substituents.

Preferred technetium radiopharmaceuticals of the present invention are comprised of a hydrazido or diazenido bonding unit and an ancillary ligand, A_{L1} , or a bonding unit and two types of
35 ancillary A_{L1} and A_{L2} , or a tetradentate chelator comprised of two nitrogen and two sulfur atoms. Ancillary ligands A_{L1} are

comprised of two or more hard donor atoms such as oxygen and amine nitrogen (sp^3 hybridized). The donor atoms occupy at least two of the sites in the coordination sphere of the radionuclide metal; the ancillary ligand A_{L1} serves as one of the three ligands in the ternary ligand system. Examples of ancillary ligands A_{L1} include but are not limited to dioxygen ligands and functionalized aminocarboxylates. A large number of such ligands are available from commercial sources.

Ancillary dioxygen ligands include ligands that coordinate to the metal ion through at least two oxygen donor atoms. Examples include but are not limited to: glucoheptonate, gluconate, 2-hydroxyisobutyrate, lactate, tartrate, mannitol, glucarate, maltol, Kojic acid, 2,2-bis(hydroxymethyl)propionic acid, 4,5-dihydroxy-1,3-benzene disulfonate, or substituted or unsubstituted 1,2 or 3,4 hydroxypyridinones. (The names for the ligands in these examples refer to either the protonated or non-protonated forms of the ligands.)

Functionalized aminocarboxylates include ligands that have a combination of amine nitrogen and oxygen donor atoms. Examples include but are not limited to: iminodiacetic acid, 2,3-diaminopropionic acid, nitrilotriacetic acid, N,N'-ethylenediamine diacetic acid, N,N,N'-ethylenediamine triacetic acid, hydroxyethylethylenediamine triacetic acid, and N,N'-ethylenediamine bis-hydroxyphenylglycine. (The names for the ligands in these examples refer to either the protonated or non-protonated forms of the ligands.)

A series of functionalized aminocarboxylates are disclosed by Bridger et. al. in U.S. Patent 5,350,837, herein incorporated by reference, that result in improved rates of formation of technetium labeled hydrazino modified proteins. We have determined that certain of these aminocarboxylates result in improved yields of the radiopharmaceuticals of the present invention. The preferred ancillary ligands A_{L1} functionalized aminocarboxylates that are derivatives of glycine; the most preferred is tricine (tris(hydroxymethyl)methylglycine).

The most preferred technetium radiopharmaceuticals of the present invention are comprised of a hydrazido or diazenido bonding unit and two types of ancillary designated A_{L1} and A_{L2} , or a diaminedithiol chelator. The second type of ancillary
5 ligands A_{L2} are comprised of one or more soft donor atoms selected from the group: phosphine phosphorus, arsine arsenic, imine nitrogen (sp^2 hybridized), sulfur (sp^2 hybridized) and carbon (sp hybridized); atoms which have p-acid character. Ligands A_{L2} can be monodentate, bidentate or tridentate, the
10 denticity is defined by the number of donor atoms in the ligand. One of the two donor atoms in a bidentate ligand and one of the three donor atoms in a tridentate ligand must be a soft donor atom. We have disclosed in co-pending U.S.S.N. 08/415,908, and U.S.S.N. 60/013360 and 08/646,886, the disclosures of which are
15 herein incorporated by reference in their entirety, that radiopharmaceuticals comprised of one or more ancillary or co-ligands A_{L2} are more stable compared to radiopharmaceuticals that are not comprised of one or more ancillary ligands, A_{L2} ; that is, they have a minimal number of isomeric forms, the
20 relative ratios of which do not change significantly with time, and that remain substantially intact upon dilution.

The ligands A_{L2} that are comprised of phosphine or arsine donor atoms are trisubstituted phosphines, trisubstituted arsines, tetrasubstituted diphosphines and tetrasubstituted
25 diarsines. The ligands A_{L2} that are comprised of imine nitrogen are unsaturated or aromatic nitrogen-containing, 5 or 6-membered heterocycles. The ligands that are comprised of sulfur (sp^2 hybridized) donor atoms are thiocarbonyls, comprised of the moiety $C=S$. The ligands comprised of carbon (sp
30 hybridized) donor atoms are isonitriles, comprised of the moiety CNR , where R is an organic radical. A large number of such ligands are available from commercial sources. Isonitriles can be synthesized as described in European Patent 0107734 and in U.S. Patent No. 4,988,827, herein incorporated by reference.

Preferred ancillary ligands A_{L2} are trisubstituted phosphines and unsaturated or aromatic 5 or 6 membered heterocycles. The most preferred ancillary ligands A_{L2} are trisubstituted phosphines and unsaturated 5 membered
5 heterocycles.

The ancillary ligands A_{L2} may be substituted with alkyl, aryl, alkoxy, heterocycle, aralkyl, alkaryl and arylalkaryl groups and may or may not bear functional groups comprised of heteroatoms such as oxygen, nitrogen, phosphorus or sulfur.
10 Examples of such functional groups include but are not limited to: hydroxyl, carboxyl, carboxamide, nitro, ether, ketone, amino, ammonium, sulfonate, sulfonamide, phosphonate, and phosphonamide. The functional groups may be chosen to alter the lipophilicity and water solubility of the ligands which may
15 affect the biological properties of the radiopharmaceuticals, such as altering the distribution into non-target tissues, cells or fluids, and the mechanism and rate of elimination from the body.

Chelators for magnetic resonance imaging contrast agents
20 are selected to form stable complexes with paramagnetic metal ions, such as Gd(III), Dy(III), Fe(III), and Mn(II), are selected from cyclic and acyclic polyaminocarboxylates such as DTPA, DOTA, DO3A, 2-benzyl-DOTA, alpha-(2-phenethyl)1,4,7,10-tetraazacyclododecane-1-acetic-4,7,10-tris(methylacetic)acid, 2-
25 benzyl-cyclohexyldiethylenetriaminepentaacetic acid, 2-benzyl-6-methyl-DTPA, and 6,6"-bis[N,N,N",N"-tetra(carboxymethyl)aminomethyl]-4'-(3-amino-4-methoxyphenyl)-2,2':6',2"-terpyridine.

There are three key features of the pharmaceuticals of the
30 present invention that determine their efficacy: MMP selectivity, inhibitory potency, typically expressed as the K_i value, and the rate of clearance from the blood. Preferred pharmaceuticals of the present invention are comprised of inhibitors, Q, which exhibit selectivity for MMP-1, MMP-2, MMP-
35 3, MMP-9, or MMP-14 alone or in combination over the other MMPs. Most preferred are comprised of inhibitors, Q, which exhibit

selectivity for MMP-2, MMP-9, or MMP-14 alone or in combination over the other MMPs. K_i values for the preferred pharmaceuticals of the present invention are <100 nM for one or more of MMP-1, MMP-2, MMP-3, MMP-9, or MMP-14. K_i values for the most preferred pharmaceuticals of the present invention are <10 nM for one or more of MMP-2, MMP-9, or MMP-14.

The rate of clearance from the blood is of particular importance for cardiac imaging procedures, since the cardiac blood pool is large compared to the disease foci that one desires to image. For an effective cardiac imaging agent, the target to background ratios (disease foci-to-blood and disease foci-to-muscle) need to be greater or equal to 1.5, preferably greater or equal to 2.0, and more preferably even greater. Preferred pharmaceuticals of the present invention have blood clearance rates that result in $< 10\%$ i.d./g at 2 hours post-injection, measured in a mouse model, or $< 0.5\%$ i.d./g at 2 hours post-injection, measured in a dog model. Most preferred pharmaceuticals of the present invention have blood clearance rates that result in $< 3\%$ i.d./g at 2 hours post-injection, measured in a mouse model, or $< 0.05\%$ i.d./g at 2 hours post-injection, measured in a dog model.

The technetium radiopharmaceuticals of the present invention comprised of a hydrazido or diazenido bonding unit can be easily prepared by admixing a salt of a radionuclide, a reagent of the present invention, an ancillary ligand AL_1 , an ancillary ligand AL_2 , and a reducing agent, in an aqueous solution at temperatures from 0 to 100 °C. The technetium radiopharmaceuticals of the present invention comprised of a tetradentate chelator having two nitrogen and two sulfur atoms can be easily prepared by admixing a salt of a radionuclide, a reagent of the present invention, and a reducing agent, in an aqueous solution at temperatures from 0 to 100 °C.

When the bonding unit in the reagent of the present invention is present as a hydrazone group, then it must first be converted to a hydrazine, which may or may not be protonated, prior to complexation with the metal radionuclide. The

conversion of the hydrazone group to the hydrazine can occur either prior to reaction with the radionuclide, in which case the radionuclide and the ancillary or co-ligand or ligands are combined not with the reagent but with a hydrolyzed form of the reagent bearing the chelator or bonding unit, or in the presence of the radionuclide in which case the reagent itself is combined with the radionuclide and the ancillary or co-ligand or ligands. In the latter case, the pH of the reaction mixture must be neutral or acidic.

10 Alternatively, the radiopharmaceuticals of the present invention comprised of a hydrazido or diazenido bonding unit can be prepared by first admixing a salt of a radionuclide, an ancillary ligand AL_1 , and a reducing agent in an aqueous solution at temperatures from 0 to 100 °C to form an
15 intermediate radionuclide complex with the ancillary ligand AL_1 then adding a reagent of the present invention and an ancillary ligand AL_2 and reacting further at temperatures from 0 to 100 °C.

 Alternatively, the radiopharmaceuticals of the present
20 invention comprised of a hydrazido or diazenido bonding unit can be prepared by first admixing a salt of a radionuclide, an ancillary ligand AL_1 , a reagent of the present invention, and a reducing agent in an aqueous solution at temperatures from 0 to 100 °C to form an intermediate radionuclide complex, and then
25 adding an ancillary ligand AL_2 and reacting further at temperatures from 0 to 100 °C.

 The technetium radionuclides are preferably in the chemical form of pertechnetate or perrhenate and a pharmaceutically acceptable cation. The pertechnetate salt form is preferably
30 sodium pertechnetate such as obtained from commercial Tc-99m generators. The amount of pertechnetate used to prepare the radiopharmaceuticals of the present invention can range from 0.1 mCi to 1 Ci, or more preferably from 1 to 200 mCi.

 The amount of the reagent of the present invention used to
35 prepare the technetium radiopharmaceuticals of the present invention can range from 0.01 µg to 10 mg, or more preferably

from 0.5 μ g to 200 μ g. The amount used will be dictated by the amounts of the other reactants and the identity of the radiopharmaceuticals of the present invention to be prepared.

The amounts of the ancillary ligands A_{L1} used can range
5 from 0.1 mg to 1 g, or more preferably from 1 mg to 100 mg. The exact amount for a particular radiopharmaceutical is a function of identity of the radiopharmaceuticals of the present invention to be prepared, the procedure used and the amounts and identities of the other reactants. Too large an amount of A_{L1}
10 will result in the formation of by-products comprised of technetium labeled A_{L1} without a biologically active molecule or by-products comprised of technetium labeled biologically active molecules with the ancillary ligand A_{L1} but without the ancillary ligand A_{L2} . Too small an amount of A_{L1} will result in
15 other by-products such as technetium labeled biologically active molecules with the ancillary ligand A_{L2} but without the ancillary ligand A_{L1} , or reduced hydrolyzed technetium, or technetium colloid.

The amounts of the ancillary ligands A_{L2} used can range
20 from 0.001 mg to 1 g, or more preferably from 0.01 mg to 10 mg. The exact amount for a particular radiopharmaceutical is a function of the identity of the radiopharmaceuticals of the present invention to be prepared, the procedure used and the amounts and identities of the other reactants. Too large an
25 amount of A_{L2} will result in the formation of by-products comprised of technetium labeled A_{L2} without a biologically active molecule or by-products comprised of technetium labeled biologically active molecules with the ancillary ligand A_{L2} but without the ancillary ligand A_{L1} . If the reagent bears one or
30 more substituents that are comprised of a soft donor atom, as defined above, at least a ten-fold molar excess of the ancillary ligand A_{L2} to the reagent of formula 2 is required to prevent the substituent from interfering with the coordination of the ancillary ligand A_{L2} to the metal radionuclide.

In another embodiment of the current invention, a scintigraphic image of a radiolabeled MMPI compound would be acquired at the same time as a scintigraphic image of a radiolabeled cardiac perfusion imaging agent. This simultaneous dual isotope imaging would be done by utilizing radioisotopes of the MMPI and perfusion imaging agents which had spectrally separable gamma emission energies. For example, a Tc99m cardiac perfusion imaging agent (such as Tc99m-Sestamibi) or Tl201 (as Thallous Chloride), and an In111-labeled MMPI compound would be imaged simultaneously with a standard gamma camera. This is possible because the Tc99m gamma energy of ~140KeV or the Tl201 gamma energy of ~80KeV are easily separable from the In111 gamma energies of ~160KeV and 250KeV. This simultaneous imaging of cardiac perfusion and extracellular matrix degradation (as evidenced by MMPI compound localization) is extremely useful for improved anatomic assessment of the location of MMPI compound distribution in the heart based on the comparison to the perfusion distribution seen on the Tc99m-Sestamibi or Tl201 image. In addition, the simultaneous imaging of perfusion and extracellular matrix degradation allows a more complete assessment of the underlying cardiac disease, both in terms of blood flow alterations and biochemical changes, in a single imaging session on a patient.

The simultaneous dual-isotope imaging of cardiac perfusion and extracellular matrix degradation allows the localization of sites of vulnerable plaque and cardiac perfusion to be visualized during one imaging session. In addition, the simultaneous imaging of tissue changes associated with congestive heart failure (from the MMPI imaging agent) and coronary artery disease (from the perfusion imaging agent) is extremely useful in characterizing the underlying causes of CHF.

The simultaneous imaging of different radioisotopically-labeled radiopharmaceuticals in patients has been reported. For example, Antunes, et al., Am J. Cardiol. 1992; 70: 426-431, have demonstrated that it is possible to image myocardial infarction with an In111-antimyosin antibody along with the imaging of

cardiac perfusion with Tl201. However, the dual isotope imaging of the present invention is new, because it is the first reported approach to the simultaneous, dual isotope imaging of a radiolabeled MMPI compound and a cardiac perfusion imaging compound. The combination of MMPI scintigraphic imaging with perfusion imaging provides the imaging physician with an extraordinary amount of clinical information regarding ischemic coronary artery disease or congestive heart failure in one imaging session.

Suitable reducing agents for the synthesis of the radiopharmaceuticals of the present invention include stannous salts, dithionite or bisulfite salts, borohydride salts, and formamidinesulfinic acid, wherein the salts are of any pharmaceutically acceptable form. The preferred reducing agent is a stannous salt. The amount of a reducing agent used can range from 0.001 mg to 10 mg, or more preferably from 0.005 mg to 1 mg.

The specific structure of a radiopharmaceutical of the present invention comprised of a hydrazido or diazenido bonding unit will depend on the identity of the reagent of the present invention used, the identity of any ancillary ligand A_{L1} , the identity of any ancillary ligand A_{L2} , and the identity of the radionuclide. Radiopharmaceuticals comprised of a hydrazido or diazenido bonding unit synthesized using concentrations of reagents of $<100 \mu\text{g/mL}$, will be comprised of one hydrazido or diazenido group. Those synthesized using $>1 \text{ mg/mL}$ concentrations will be comprised of two hydrazido or diazenido groups from two reagent molecules. For most applications, only a limited amount of the biologically active molecule can be injected and not result in undesired side-effects, such as chemical toxicity, interference with a biological process or an altered biodistribution of the radiopharmaceutical. Therefore, the radiopharmaceuticals which require higher concentrations of the reagents comprised in part of the biologically active molecule, will have to be diluted or purified after synthesis to avoid such side-effects.

The identities and amounts used of the ancillary ligands AL_1 and AL_2 will determine the values of the variables y and z . The values of y and z can independently be an integer from 1 to 2. In combination, the values of y and z will result in a
5 technetium coordination sphere that is made up of at least five and no more than seven donor atoms. For monodentate ancillary ligands AL_2 , z can be an integer from 1 to 2; for bidentate or tridentate ancillary ligands AL_2 , z is 1. The preferred combination for monodentate ligands is y equal to 1 or 2 and z
10 equal to 1. The preferred combination for bidentate or tridentate ligands is y equal to 1 and z equal to 1.

The indium, copper, gallium, and yttrium radiopharmaceuticals of the present invention can be easily prepared by admixing a salt of a radionuclide and a reagent of
15 the present invention, in an aqueous solution at temperatures from 0 to 100 °C. These radionuclides are typically obtained as a dilute aqueous solution in a mineral acid, such as hydrochloric, nitric or sulfuric acid. The radionuclides are combined with from one to about one thousand equivalents of the
20 reagents of the present invention dissolved in aqueous solution. A buffer is typically used to maintain the pH of the reaction mixture between 3 and 10.

The gadolinium, dysprosium, iron and manganese metallopharmaceuticals of the present invention can be easily
25 prepared by admixing a salt of the paramagnetic metal ion and a reagent of the present invention, in an aqueous solution at temperatures from 0 to 100 °C. These paramagnetic metal ions are typically obtained as a dilute aqueous solution in a mineral acid, such as hydrochloric, nitric or sulfuric acid. The
30 paramagnetic metal ions are combined with from one to about one thousand equivalents of the reagents of the present invention dissolved in aqueous solution. A buffer is typically used to maintain the pH of the reaction mixture between 3 and 10.

The total time of preparation will vary depending on the
35 identity of the metal ion, the identities and amounts of the reactants and the procedure used for the preparation. The

preparations may be complete, resulting in > 80% yield of the radiopharmaceutical, in 1 minute or may require more time. If higher purity metallopharmaceuticals are needed or desired, the products can be purified by any of a number of techniques well known to those skilled in the art such as liquid chromatography, solid phase extraction, solvent extraction, dialysis or ultrafiltration.

Buffers useful in the preparation of metallopharmaceuticals and in diagnostic kits useful for the preparation of said radiopharmaceuticals include but are not limited to phosphate, citrate, sulfosalicylate, and acetate. A more complete list can be found in the United States Pharmacopeia.

Lyophilization aids useful in the preparation of diagnostic kits useful for the preparation of radiopharmaceuticals include but are not limited to mannitol, lactose, sorbitol, dextran, Ficoll, and polyvinylpyrrolidone (PVP).

Stabilization aids useful in the preparation of metallopharmaceuticals and in diagnostic kits useful for the preparation of radiopharmaceuticals include but are not limited to ascorbic acid, cysteine, monothioglycerol, sodium bisulfite, sodium metabisulfite, gentisic acid, and inositol.

Solubilization aids useful in the preparation of metallopharmaceuticals and in diagnostic kits useful for the preparation of radiopharmaceuticals include but are not limited to ethanol, glycerin, polyethylene glycol, propylene glycol, polyoxyethylene sorbitan monooleate, sorbitan monooleate, polysorbates, poly(oxyethylene)poly(oxypropylene)poly(oxyethylene) block copolymers (Pluronic) and lecithin. Preferred solubilizing aids are polyethylene glycol, and Pluronic.

Bacteriostats useful in the preparation of metallopharmaceuticals and in diagnostic kits useful for the preparation of radiopharmaceuticals include but are not limited to benzyl alcohol, benzalkonium chloride, chlorbutanol, and methyl, propyl or butyl paraben.

A component in a diagnostic kit can also serve more than one function. A reducing agent can also serve as a stabilization aid, a buffer can also serve as a transfer ligand, a lyophilization aid can also serve as a transfer, ancillary or co-ligand and so forth.

The diagnostic radiopharmaceuticals are administered by intravenous injection, usually in saline solution, at a dose of 1 to 100 mCi per 70 kg body weight, or preferably at a dose of 5 to 50 mCi. Imaging is performed using known procedures.

The magnetic resonance imaging contrast agents of the present invention may be used in a similar manner as other MRI agents as described in U.S. Patent 5,155,215; U.S. Patent 5,087,440; Margerstadt et al., Magn. Reson. Med., 1986, 3, 808; Runge et al., Radiology, 1988, 166, 835; and Bousquet et al., Radiology, 1988, 166, 693. Generally, sterile aqueous solutions of the contrast agents are administered to a patient intravenously in dosages ranging from 0.01 to 1.0 mmol/kg body weight.


For use as X-ray contrast agents, the compositions of the present invention should generally have a heavy atom concentration of 1 mM to 5 M, preferably 0.1 M to 2 M. Dosages, administered by intravenous injection, will typically range from 0.5 mmol/kg to 1.5 mmol/kg, preferably 0.8 mmol/kg to 1.2 mmol/kg. Imaging is performed using known techniques, preferably X-ray computed tomography.

The ultrasound contrast agents of the present invention are administered by intravenous injection in an amount of 10 to 30 μ L of the echogenic gas per kg body weight or by infusion at a rate of approximately 3 μ L/kg/min. Imaging is performed using known techniques of sonography.

Other features of the invention will become apparent in the course of the following descriptions of exemplary embodiments which are given for illustration of the invention and are not intended to be limiting thereof.


Example 1

Synthesis of 2-[[5-(3-{2-[(6-Hydroxycarbamoyl-7-isobutyl-8-oxo-2-oxa-9-aza-bicyclo[10.2.2]hexadeca-1(15),12(16),13-triene-10-carbonyl)-amino]-acetylamino}-propylcarbamoyl)-pyridin-2-yl]-hydrazonomethyl}-benzenesulfonic acid



5

A. Preparation of [3-(2-Benzyloxycarbonylamino-acetylamino)-propyl]-carbamic acid *tert*-butyl ester




10

To 3 grams of (3-Amino-propyl)-carbamic acid *tert*-butyl ester in 15 ml of dimethylformamide was added 3 grams of N-benzyloxycarbonyl glycine, 4.7 mL of N-methylmorpholine and 5.06 grams of TBTU. The reaction was cooled to 0 degrees C for 30 minutes then allowed to stir at room temperature overnight. The volatiles were removed under reduced pressure and the resulting material was dissolved in ethyl acetate and washed with 10% citric acid. The aqueous was extracted an additional two times with ethyl acetate, combined and washed with water, saturated aqueous sodium bicarbonate, water, brine and dried over MgSO₄. The volatiles were removed under reduced pressure and the resulting material was crystallized from EtOAc / hexane affording 4.55 grams of the desired product as a tan solid. LRMS found 388.3 = (M+Na)⁺


25

B. Preparation of [3-(2-Amino-acetylamino)-propyl]-carbamic acid *tert*-butyl ester




To 4.16 grams of the compound from Example 1A in 25 mL of methanol was added 0.5 grams of 10% Pd-C. The reaction was stirred under H₂ (balloon) for 2 hours. The reaction was
 5 filtered through a 0.45 μ m PTFE filter and the volatiles were removed under reduced pressure affording 2.5 grams of the desired product. LRMS found 232.3 (M+H)⁺1.

C. Preparation of 3-{2-[(6-Benzyloxycarbamoyl-7-isobutyl-8-oxo-2-oxa-9-aza-bicyclo[10.2.2]hexadeca-1(15),12(16),13-triene-10-carbonyl)-amino}-propyl)-carbamic acid *tert*-butyl
 10 ester




To 0.25 grams of 6-Benzyloxycarbamoyl-7-isobutyl-9-oxo-2-oxa-9-aza-bicyclo[10.2.2]hexadeca-1(15),12(16),13-triene-10-carboxylic acid in 10 mL of dimethylformamide was added 0.17 ml N-methylmorpholine and 0.217 grams of TBTU. After 10 minutes 0.359 grams of the compound from Example 1B was added. The reaction was allowed to stir at room temperature overnight, then
 20 it was heated at 70 degrees C for 30 minutes. The volatiles were removed under reduced pressure and the resulting material was dissolved in EtOAc, washed with 10 % aqueous citric acid, water, saturated NaHCO₃, brine and dried over MgSO₄. The resulting material was chromatographed on silica gel eluting
 25 with 2% MeOH/ CHCl₃ affording 0.274 grams of the desired product.

- D. Preparation of 3-{2-[(6-Hydroxycarbamoyl-7-isobutyl-8-oxo-2-oxa-9-aza-bicyclo[10.2.2]hexadeca-1(15),12(16),13-triene-10-carbonyl)-amino]-propyl)-carbamic acid tert-butyl ester



- 5 To 0.035 grams of the compound from Example 1C in 5 mL of methanol was added 0.050 grams of 5% Pd/BaSO₄. The reaction was stirred under hydrogen (balloon) for 2 hours, then filtered through a 0.45 μm PTFE filter and the volatiles were removed under reduced pressure affording 0.031 grams of the desired
10 compound. LRMS found 604.4 (M-H)⁻¹.

- E. Preparation of 7-Isobutyl-8-oxo-2-oxa-9-aza-bicyclo[10.2.2]hexadeca-1(15),12(16),13-triene-6,10-dicarboxylic acid 10-[[[3-amino-propylcarbonyl)-methyl]-amide} 6-hydroxyamide trifluoroacetic acid salt
15



- To 0.025 grams of the compound from Example 1D was added 1 mL of trifluoroacetic acid. The reaction was stirred 1 hour and
20 the volatiles were removed under reduced pressure affording 0.017 grams of the desired compound. LRMS found 506.4 (M+H)⁺¹.

F. Preparation of 2-[[5-(3-{2-[(6-Hydroxycarbamoyl-7-isobutyl-8-oxo-2-oxa-9-aza-bicyclo[10.2.2]hexadeca-1(15),12(16),13-triene-10-carbonyl)-amino]-acetylamino}-propylcarbamoyl)-pyridin-2-yl]-hydrazonomethyl]-benzenesulfonic acid


5

To a stirred solution of 0.050 grams of the compound from Example 1E was added 0.031 mL of N-methylmorpholine and 0.035 grams of 6-[N"-(2-Sodio-sulfo-benzylidene)-hydrazino]-nicotinic acid 2,5-dioxo-pyrrolidin-1-yl ester. The reaction was stirred at ambient temperature overnight. Volatiles were removed under reduced pressure and the resulting material was purified by reverse phase HPLC affording 0.08 grams of the desired compound. LRMS found 807 (M-H)⁻¹.

15


Example 2

Synthesis of 2-[[5-(4-{[(6-Hydroxycarbamoyl-7-isobutyl-8-oxo-2-oxa-9-aza-bicyclo[10.2.2]hexadeca-1(15),12(16),13-triene-10-carbonyl)-amino]-methyl}-benzylcarbamoyl)-pyridin-2-yl]-hydrazonomethyl]-benzenesulfonic acid



20

A. Preparation of (4-Aminomethyl-benzyl)carbamic acid tert-butyl ester




To a stirred solution of 5.3 grams of p-xylenediamine 10 in 20 mL of dimethylformamide was added a solution of 2.12 grams of di-tert-butyl-dicarbonate in 50 mL of dimethylformamide by syringe pump over 1 hour. After stirring an additional 10 minutes the volatiles were removed under reduced pressure and

the resulting material was chromatographed on silica gel eluting with 5% MeOH/CHCl₃ affording 2 grams of the desired compound.


LRMS found 237.2 (M+H)⁺1.

- 5 B. Preparation of (4-[[[6-Benzyloxycarbamoyl-7-isobutyl-8-oxo-2-oxa-9-aza-bicyclo[10.2.2]hexadeca-1(15),12(16),13-triene-10-carbonyl)-amino]-methyl)-benzyl)-carbamic acid tert-butyl ester



- 10 To 0.20 grams of 6-Benzyloxycarbamoyl-7-isobutyl-9-oxo-2-oxa-9-aza-bicyclo[10.2.2]hexadeca-1(15),12(16),13-triene-10-carboxylic acid in 5 mL of dimethylformamide was added 0.18 mL of n-methylmorpholine and 0.173 grams of TBTU. After stirring 20 minutes 0.293 grams of the compound from Example 2A was
- 15 added. After stirring at ambient temperature overnight the reaction was heated to 80 degrees C for 30 minutes. The volatiles were removed under reduced pressure and the resulting material was dissolved in EtOAc, washed with 10 % aqueous citric acid, water, saturated NaHCO₃, brine and dried over MgSO₄. The
- 20 volatiles were removed under reduced pressure affording 0.296 grams of the desired compound. LRMS found 699.4 (M-H)⁻1.


- C. Preparation of (4-[[[6-Hydroxycarbamoyl-7-isobutyl-8-oxo-2-oxa-9-aza-bicyclo[10.2.2]hexadeca-1(15),12(16),13-triene-10-carbonyl)-amino]-methyl]-benzyl)-carbamic acid tert-butyl ester



5

To 0.275 grams of the compound from Example 2B in 20 mL of methanol was added 0.50 grams of pre-hydrogenated 5% Pd-BaSO₄. The reaction was stirred 3 hours under H₂ (Balloon) at which time an additional portion of 0.25 grams of 5% Pd-BaSO₄ was added and the stirring was continued for another hour. The mixture was filtered through a 0.45 μm PTFE filter and the volatiles were removed under reduced pressure affording 0.24 grams of the desired compound. LRMS found 609.4 (M-H)⁻¹.

- 15 D. Preparation of 7-Isobutyl-8-oxo-2-oxa-9-aza-bicyclo[10.2.2]hexadeca-1(15),12(16),13-triene-6,10-dicarboxylic acid 10-(4-aminomethyl-benzylamice) 6-hydroxyamide trifluoroacetic acid salt



20 To 0.225 grams of the compound from Example 2C in 5 mL of CH₂Cl₂ was added 2 mL of trifluoroacetic acid. The reaction was stirred one hour at ambient temperature. The volatiles were removed under reduced pressure affording the desired compound. LRMS found 509.4 (M-H)⁻¹.

25

E. Preparation of 2-([5-(4-((6-Hydroxycarbamoyl-7-isobutyl-8-oxo-2-oxa-9-aza-bicyclo[10.2.2]hexadeca-1(15),12(16),13-triene-10-carbonyl)-amino)-methyl)-benzylcarbamoyl)-pyridin-2-yl]-hydrazonomethyl)-benzenesulfonic acid

5


To 0.050 grams of the compound from Example 2D in 1 mL of dimethylformamide was added 0.031 mL of N-methylmorpholine and 0.035 grams of 6-[N''-(2-Sodio-sulfo-benzylidene)-hydrazino]-nicotinic acid 2,5-dioxo-pyrrolidin-1-yl ester. After stirring overnight at ambient temperature the volatiles were removed under reduced pressure and the resulting material was purified by reverse phase HPLC affording 0.06 grams of the desired compound. LRMS found 814 (M+H)⁺1.

15

Example 3

Synthesis of 2-[7-((N-[3-(2-([7-(N-hydroxycarbamoyl) (3S,6R,7S)-4-aza-6-(2-methylpropyl)-11-oxa-5-oxobicyclo[10.2.2]hexadeca-1(15),12(16),13-trien-3-yl]carbonylamino)acetylaminopropyl]carbamoyl)methyl)-1,4,7,10-tetraaza-4,10-bis(carboxymethyl)cyclododecyl]acetic acid

20



A solution of the commercially (Macrocyclics) available DOTA tri-t-butyl ester (1.5 mmol) and Hunig's base (6 mmol) in anhydrous DMF are treated with HBTU (1.25 mmol) and allowed to react for 15 min at ambient temperatures under nitrogen. 2-([7-(N-Hydroxycarbamoyl) (3S,6R,7S)-4-aza-6-(2-methylpropyl)-11-oxa-5-oxobicyclo[10.2.2]hexadeca-1(15),12(16),13-trien-3-yl]carbonylamino)-N-(3-aminopropyl)acetamide TFA salt (1 mmol) is added to this solution and stirring is continued at ambient temperatures under nitrogen for 18 h. The DMF is removed under

30


vacuum and the resulting residue is triturated in ethyl acetate or diethyl ether and filtered. If necessary, the crude is purified by preparative HPLC on a C18 column using a water:ACN:0.1% TFA gradient and the product fraction is

5 lyophilized to give the DOTA-conjugate. The DOTA conjugate is stirred in degassed TFA at room temperature under nitrogen for 2 h. The solution is concentrated and the resulting residue is purified by preparative HPLC on a C18 column using a water:ACN:0.1% TFA gradient. The product fraction is

10 lyophilized to give the title compound.

Example 4

Synthesis of 2-{7-[(N-{[4-({[7-(N-hydroxycarbamoyl)(3S,6R,7S)-4-
 15 aza-6-(2-methylpropyl)-11-oxa-5-oxobicyclo[10.2.2]hexadeca-
 1(15),12(16),13-trien-3-yl]-
 carbonylamino)methyl]phenyl)methyl}carbamoyl)methyl]-1,4,7,10-
 tetraaza-4,10-bis(carboxymethyl)cyclododecyl}acetic acid



20 A solution of the commercially (Macrocylics) available DOTA tri-t-butyl ester (1.5 mmol) and Hunig's base (6 mmol) in anhydrous DMF are treated with HBTU (1.25 mmol) and allowed to react for 15 min at ambient temperatures under nitrogen. [7-(N-Hydroxycarbamoyl)(3S,6R,7S)-4-aza-6-

25 (2-methylpropyl)-11-oxa-5-oxobicyclo[10.2.2]hexadeca-
 1(15),12(16),13-trien-3-yl]-N-{[4-
 (aminomethyl)phenyl)methyl}carboxamide TFA salt
 (1 mmol) is added to this solution and stirring is continued at ambient temperatures under nitrogen for 18 h. The DMF is


30 removed under vacuum and the resulting residue is triturated in ethyl acetate or diethyl ether and filtered. If necessary, the

crude is purified by preparative HPLC on a C18 column using a water:ACN:0.1% TFA gradient and the product fraction is lyophilized to give the DOTA-conjugate.


The DOTA conjugate is stirred in degassed TFA at room temperature under nitrogen for 2 h. The solution is concentrated and the resulting residue is purified by preparative HPLC on a C18 column using a water:ACN:0.1% TFA gradient. The product fraction is lyophilized to give the title compound.

Example 5

Synthesis of 2-(7-([N-(1-(N-[3-(2-([7-(N-hydroxycarbamoyl) (3S, 6R, 7S)-4-aza-6-(2-methylpropyl)-11-oxa-5-oxobicyclo[10.2.2]hexadeca-1(15), 12(16), 13-trien-3-yl]carbonylamino}acetylamino)propyl]carbamoyl)-2-sulfoethyl)carbamoyl)methyl]-1, 4, 7, 10-tetraaza-4, 10-bis(carboxymethyl)cyclododecyl)acetic acid



A. Preparation of N-[3-(2-([7-(N-hydroxycarbamoyl) (3S, 6R, 7S)-4-aza-6-(2-methylpropyl)-11-oxa-5-oxobicyclo[10.2.2]hexadeca-1(15), 12(16), 13-trien-3-yl]carbonylamino}acetylamino)propyl]-2-aminopropanesulfonic acid



2-([7-(N-Hydroxycarbamoyl) (3S, 6R, 7S)-4-aza-6-(2-methylpropyl)-11-oxa-5-oxobicyclo[10.2.2]hexadeca-1(15), 12(16), 13-trien-3-

yl]carbonylamino)-N-(3-aminopropyl)acetamide TFA salt (1 mmol) is dissolved in anhydrous DMF, and treated with the N-hydroxysuccinimide ester (1.5 mmol) of Boc-cysteic acid (as described in *Liebigs Ann. Chem.* 1979, 776-783) and Hunig's base.

5 The solution is stirred at ambient temperatures under nitrogen for 18 h, and the DMF is removed under vacuum. The resulting residue is purified by preparative HPLC on a C18 column using a water:ACN:0.1% TFA gradient. The product fraction is lyophilized to give a solid which is dissolved in degassed TFA

10 and stirred at ambient temperatures for 30 min. The solution is concentrated under vacuum, and the resulting residue is dissolved in 50% ACN and lyophilized to give the boc deprotected product N-[3-(2-([7-(N-hydroxycarbamoyl) (3S,6R,7S)-4-aza-6-(2-methylpropyl)-11-oxa-5-oxobicyclo[10.2.2]hexadeca-

15 1(15),12(16),13-trien-3-yl]carbonylamino)acetylaminopropyl]-2-aminopropanesulfonic acid.

B. Preparation of 2-(7-([N-(1-{N-[3-(2-([7-(N-hydroxycarbamoyl) (3S,6R,7S)-4-aza-6-(2-methylpropyl)-11-oxa-5-

20 oxobicyclo[10.2.2]hexadeca-1(15),12(16),13-trien-3-yl]carbonylamino)acetylaminopropyl]carbamoyl)-2-sulfoethyl)carbamoyl)methyl)-1,4,7,10-tetraaza-4,10-bis(carboxymethyl)cyclododecyl)acetic acid.

25 A solution of the commercially (Macrocylics) available DOTA tri-t-butyl ester (1.5 mmol) and Hunig's base (6 mmol) in anhydrous DMF are treated with HBTU (1.25 mmol) and allowed to react for 15 min at ambient temperatures under nitrogen. N-[3-(2-([7-(N-Hydroxycarbamoyl) (3S,6R,7S)-4-aza-6-(2-methylpropyl)-

30 11-oxa-5-oxobicyclo[10.2.2]hexadeca-1(15),12(16),13-trien-3-yl]carbonylamino)acetylaminopropyl]-2-aminopropanesulfonic acid is added to this solution and stirring is continued at ambient temperatures under nitrogen for 18 h. The DMF is removed under vacuum and the resulting residue is triturated in ethyl acetate


35 or diethyl ether and filtered. If necessary, the crude is purified by preparative HPLC on a C18 column using a

water:ACN:0.1% TFA gradient and the product fraction is lyophilized to give the DOTA-conjugate.


The DOTA conjugate is stirred in degassed TFA at room temperature under nitrogen for 2 h. The solution is concentrated and the resulting residue is purified by preparative HPLC on a C18 column using a water:ACN:0.1% TFA gradient. The product fraction is lyophilized to give the title compound.

Example 6

Synthesis of 2-[7-({N-[1-(N-{[4-({[7-(N-hydroxycarbamoyl) (3S,6R,7S)-4-aza-6-(2-methylpropyl)-11-oxa-5-oxobicyclo[10.2.2]hexadeca-1(15),12(16),13-trien-3-yl]-carbonylamino)methyl}phenyl)methyl}carbamoyl)-2-sulfoethyl]carbamoyl)methyl)-1,4,7,10-tetraaza-4,10-bis(carboxymethyl)cyclododecyl]acetic acid



A. Preparation of N-{[4-({[7-(N-hydroxycarbamoyl) (3S,6R,7S)-4-aza-6-(2-methylpropyl)-11-oxa-5-oxobicyclo[10.2.2]hexadeca-1(15),12(16),13-trien-3-yl]-carbonylamino)methyl}phenyl)methyl}-2-aminopropanesulfonic acid



[7-(N-Hydroxycarbamoyl) (3S, 6R, 7S)-4-aza-6-(2-methylpropyl)-11-oxa-5-oxobicyclo[10.2.2]hexadeca-1(15), 12(16), 13-trien-3-yl]-N-{{4-(aminomethyl)phenyl}methyl}carboxamide TFA salt

(1 mmol) is dissolved in anhydrous DMF, and treated with the N-
5 hydroxysuccinimide ester (1.5 mmol) of Boc-cysteic acid (as described in *Liebigs Ann. Chem.* 1979, 776-783) and Hunig's base. The solution is stirred at ambient temperatures under nitrogen for 18 h, and the DMF is removed under vacuum. The resulting residue is purified by preparative HPLC on a C18 column using a
10 water:ACN:0.1% TFA gradient. The product fraction is lyophilized to give a solid, which is dissolved in degassed TFA and stirred at ambient temperatures for 30 min. The solution is concentrated under vacuum, and the resulting residue is dissolved in 50% ACN and lyophilized to give the boc deprotected
15 product N-{{4-([7-(N-hydroxycarbamoyl) (3S, 6R, 7S)-4-aza-6-(2-methylpropyl)-11-oxa-5-oxobicyclo[10.2.2]hexadeca-1(15), 12(16), 13-trien-3-yl]-carbonylamino)methyl}phenyl)methyl}-2-aminopropanesulfonic acid.

20 B. Preparation of 2-[7-({N-[1-(N-{{4-([7-(N-hydroxycarbamoyl) (3S, 6R, 7S)-4-aza-6-(2-methylpropyl)-11-oxa-5-oxobicyclo[10.2.2]hexadeca-1(15), 12(16), 13-trien-3-yl]-carbonylamino)methyl}phenyl)methyl}carbamoyl)-2-sulfoethyl]carbamoyl)methyl)-1,4,7,10-tetraaza-4,10-
25 bis(carboxymethyl)cyclododecyl]acetic acid.

A solution of the commercially (Macrocylics) available DOTA tri-t-butyl ester (1.5 mmol) and Hunig's base (6 mmol) in anhydrous DMF are treated with HBTU (1.25 mmol) and allowed to react for 15 min at ambient temperatures under nitrogen. N-{{4-
30 ([7-(N-hydroxycarbamoyl) (3S, 6R, 7S)-4-aza-6-(2-methylpropyl)-11-oxa-5-oxobicyclo[10.2.2]hexadeca-1(15), 12(16), 13-trien-3-yl]-carbonylamino)methyl}phenyl)methyl}-2-aminopropanesulfonic acid is added to this solution and stirring is continued at ambient temperatures under nitrogen for 18 h. The DMF is removed under
35 vacuum and the resulting residue is triturated in ethyl acetate or diethyl ether and filtered. If necessary, the crude is

purified by preparative HPLC on a C18 column using a water:ACN:0.1% TFA gradient and the product fraction is lyophilized to give the DOTA-conjugate.


The DOTA conjugate is stirred in degassed TFA at room temperature under nitrogen for 2 h. The solution is concentrated and the resulting residue is purified by preparative HPLC on a C18 column using a water:ACN:0.1% TFA gradient. The product fraction is lyophilized to give the title compound.

10

Example 7

Synthesis of 2-({2-[(N-[3-(2-([7-(N-hydroxycarbamoyl) (3S,6R,7S)-4-aza-6-(2-methylpropyl)-11-oxa-5-oxobicyclo[10.2.2]hexadeca-1(15),12(16),13-trien-3-yl]carbonylamino}acetyl amino)propyl]carbamoyl)methyl] (carboxymethyl) amino}ethyl) {2-[bis(carboxymethyl) amino]ethyl} amino}acetic acid

15




20

To a solution of 2-([7-(N-hydroxycarbamoyl) (3S,6R,7S)-4-aza-6-(2-methylpropyl)-11-oxa-5-oxobicyclo[10.2.2]hexadeca-1(15),12(16),13-trien-3-yl]carbonylamino)-N-(3-aminopropyl)acetamide TFA salt (1 mmol) in DMF (20 mL) is added triethylamine (3 mmol). This solution is added dropwise over 4 h to a solution of diethylenetriaminepentaacetic dianhydride (3 mmol) in DMF (20 mL) and methyl sulfoxide (20 mL). The reaction mixture is then stirred for 16 h, concentrated to an oil under high vacuum and purified by preparative HPLC to give the title compound.

30

Example 8

Synthesis of 2-[(2-[(N-[[4-([7-(N-hydroxycarbamoyl) (3S, 6R, 7S)-
4-aza-6-(2-methylpropyl)-11-oxa-5-oxobicyclo[10.2.2]hexadeca-
1(15), 12(16), 13-trien-3-yl]-
5 carbonylamino)methyl]phenyl)methyl]carbamoyl)methyl](carboxymeth-
yl)amino)ethyl]{2-[bis(carboxymethyl)amino]ethyl}amino}acetic
acid




10 To a solution of [7-(N-hydroxycarbamoyl) (3S, 6R, 7S)-4-aza-6-
(2-methylpropyl)-11-oxa-5-oxobicyclo[10.2.2]hexadeca-
1(15), 12(16), 13-trien-3-yl]-N-[[4-
(aminomethyl)phenyl]methyl]carboxamide
TFA salt (1 mmol) in DMF (20 mL) is added triethylamine (3
15 mmol). This solution is added dropwise over 4 h to a solution
of diethylenetriaminepentaacetic dianhydride (3 mmol) in DMF (20
mL) and methyl sulfoxide (20 mL). The reaction mixture is then
stirred for 16 h, concentrated to an oil under high vacuum and
purified by preparative HPLC to give the title compound.

20

Example 9


Synthesis of N-[3-(2-([7-(N-hydroxycarbamoyl) (3S, 6R, 7S)-4-aza-6-
(2-methylpropyl)-11-oxa-5-oxobicyclo[10.2.2]hexadeca-
1(15), 12(16), 13-trien-3-yl]carbonylamino)acetylaminopropyl]-
25 4,5-bis[2-(ethoxyethylthio)acetylaminopentanamide



To a solution of 2-[[7-(N-hydroxycarbamoyl) (3S,6R,7S)-4-
 aza-6-(2-methylpropyl)-11-oxa-5-oxobicyclo[10.2.2]hexadeca-
 1(15),12(16),13-trien-3-yl]carbonylamino}-N-(3-
 5 aminopropyl)acetamide TFA salt (1 mmol) and triethylamine (3
 mmol) in DMF is added 2,3,5,6-tetrafluorophenyl 4,5-bis(S-1-
 ethoxyethyl-mercapto-acetamido)pentanoate (1.1 mmol), and the
 reaction mixture is stirred for 18 hours. DMF is removed in
 vacuo and the crude residue is triturated with ethyl acetate.
 10 The product is filtered, dried, and if necessary, further
 purified by preparative HPLC to give the title compound.

Example 10

Synthesis of N-[[4-([7-(N-hydroxycarbamoyl) (3S,6R,7S)-4-aza-6-
 15 (2-methylpropyl)-11-oxa-5-oxobicyclo[10.2.2]hexadeca-
 1(15),12(16),13-trien-3-yl]carbonylamino)methyl]-phenyl]methyl]-
 4,5-bis[2-(ethoxyethylthio)acetylaminol]-pentanamide




To a solution of [7-(N-hydroxycarbamoyl) (3S,6R,7S)-4-aza-6-
 (2-methylpropyl)-11-oxa-5-oxobicyclo[10.2.2]hexadeca-
 1(15),12(16),13-trien-3-yl]-N-[[4-(aminomethyl)phenyl]-
 methyl]carboxamide TFA salt (1 mmol) and triethylamine (3 mmol)
 25 in DMF is added 2,3,5,6-tetrafluorophenyl 4,5-bis(S-1-
 ethoxyethyl-mercapto-acetamido)pentanoate (1.1 mmol), and the
 reaction mixture is stirred for 18 hours. DMF is removed in

vacuo and the crude residue is triturated with ethyl acetate. The product is filtered, dried, and if necessary, further purified by preparative HPLC to give the title compound.

5

Example 11

Synthesis of 1-(1,2-Dipalmitoyl-sn-glycero-3-phosphoethanolamino)- α,ω -dicarbonylPEG₃₄₀₀-2-[[7-(N-hydroxycarbamoyl) (3S,6R,7S)-4-aza-6-(2-methylpropyl)-11-oxa-5-oxobicyclo[10.2.2]hexadeca-1(15),12(16),13-trien-3-yl]carbonylamino}-N-(3-aminopropyl)acetamide conjugate




To solution of the commercially available (Shearwater Polymers) succinimidyl ester, DSPE-PEG-NHS ester (1 mmol) in 25 ml chloroform is added 2-[[7-(N-hydroxycarbamoyl) (3S,6R,7S)-4-aza-6-(2-methylpropyl)-11-oxa-5-oxobicyclo[10.2.2]hexadeca-1(15),12(16),13-trien-3-yl]carbonylamino}-N-(3-aminopropyl)acetamide TFA salt (1 mmol). Sodium carbonate (1 mmol) and sodium sulfate (1 mmol) are added and the solution is stirred at room temperature under nitrogen for 18 h. The solvent is removed in vacuo and the crude product is purified using preparative HPLC to obtain the title compound.

25

Example 12

Synthesis of 1-(1,2-Dipalmitoyl-sn-glycero-3-phosphoethanolamino)- α,ω -dicarbonylPEG₃₄₀₀-[7-(N-hydroxycarbamoyl) (3S,6R,7S)-4-aza-6-(2-methylpropyl)-11-oxa-5-oxobicyclo[10.2.2]hexadeca-1(15),12(16),13-trien-3-yl]-N-[[4-(aminomethyl)phenyl]methyl]carboxamide conjugate

30



To solution of the commercially available (Shearwater Polymers) succinimidyl ester, DSPE-PEG-NHS ester (1 mmol) in 25 ml chloroform is added [7-(N-hydroxycarbamoyl) (3S,6R,7S)-4-aza-6-(2-methylpropyl)-11-oxa-5-oxobicyclo[10.2.2]hexadeca-1(15),12(16),13-trien-3-yl]-N-([4-(aminomethyl)phenyl]methyl)carboxamide TFA salt (1 mmol). Sodium carbonate (1 mmol) and sodium sulfate (1 mmol) are added and the solution is stirred at room temperature under nitrogen for 18 h. The solvent is removed in vacuo and the crude product is purified using preparative HPLC to obtain the title compound.

Examples 13 and 14

Synthesis of ^{99m}Tc Complexes

To a lyophilized vial containing 4.84 mg TPPTS, 6.3 mg tricine, 40 mg mannitol, succinic acid buffer, pH 4.8, and 0.1% Pluronic F-64 surfactant, was added 0.75 -1.1 mL sterile water for injection, 0.2 - 0.45 mL (20 - 40 μg) of the compounds of Examples 1 and 2, respectively, in deionized water or 50% aqueous ethanol, and 0.2 - 0.4 mL of $^{99m}\text{TcO}_4^-$ (50-120 mCi) in saline. The reconstituted kit was heated in a 100 $^\circ\text{C}$ water bath for 10 - 15 minutes, and was allowed to cool 10 minutes at room temperature. A sample was then analyzed by HPLC.

HPLC Method for Example 13

Column: Zorbax C18 , 25 cm x 4.6 mm

Flow rate : 1.0 mL/min

Solvent A: 10 mM sodium phosphate buffer, pH 6.0

Solvent B : 100 % CH_3CN

t (min)	0	20	21	30	31	40
% Solvent B	0	25	75	75	0	0

HPLC Method for Example 14

5 Column: Zorbax C18 , 25 cm x 4.6 mm

Flow rate : 1.0 mL/min

Solvent A: 10 mM sodium phosphate buffer, pH 6.0

Solvent B : 100 % CH₃CN

t (min)	0	20	21	26	27	40
10 % Solvent B	0	25	75	75	0	0


Example#	Reagent Ex. #	Ret. Time (min)	%Yield
13	1	7.8	79
14	2	16.7	81

Example 15

15


Synthesis of 2-[2-({5-[N-(5-(N-hydroxycarbamoyl) (5R)-5-{3-[4-(3,4-dimethoxyphenoxy)phenyl]-3-methyl-2-oxopyrrolidinyl}pentyl)carbamoyl] (2-pyridyl)}amino) (1Z)-2-azavinyl]benzenesulfonic acid

20



The title compound can be synthesized as shown in Scheme I from the starting materials described in the following patent applications which are hereby incorporated by reference into this patent application: U.S. Patent Application Serial Nos. 09/165,747, 08/743,439, 09/134,484, 09/247,675, 09/335,086, 09/312,066, 09/311,168, 60/127,594, and 60/127,635.

Scheme I




5

Example 16

Synthesis of 2-(2-([5-(N-{3-[3-(N-hydroxycarbonyl) (4S)-4-({4-
 [(4-methylphenyl)methoxy]piperidyl)carbonyl]piperidyl]-3-
 oxopropyl}carbonyl) (2-pyridyl)]amino) (1Z)-2-
 azaviny]benzenesulfonic acid


10



The title compound can be synthesized as shown in Scheme II from the starting materials described in the following patent


5 applications which are hereby incorporated by reference into this patent application: U.S. Patent Application Serial Nos. 09/165,747, 08/743,439, 09/134,484, 09/247,675, 09/335,086, 09/312,066, 09/311,168, 60/127,594, and 60/127,635.

10 Scheme II



Example 17

Synthesis of N-BOC-Glycine-(3-carbobenzyloxyamido)propylamide:




5

Di-isopropylethylamine (7.0 mL, 40 mmol) was added to a suspension of N-t-butyloxycarbonylglycine N-hydroxysuccinimide ester (5.56 g, 20 mmol) and N-carbobenzyloxy-1,3-diaminopropane hydrochloride (5.0 g, 20 mmol) in dichloromethane (50 ml). The solution became clear over several minutes. After 30 minutes, additional of N-t-butyloxycarbonylglycine N-hydroxysuccinimide ester (0.275 g, 1 mmol) was added. The solution was extracted with water, followed by saturated aqueous NaHCO₃, then by 0.5 N HCl. The dichloromethane solution was filtered through a short column of Na₂SO₄ and evaporated in vacuo to obtain 4.1 g (56%) of N-BOC-Glycine-(3-carbobenzyloxyamido)propylamide. MS: m/e = 366 (M+H⁺), 310 (M-C₄H₉+H⁺), 266 (M-BOC+H⁺).

10

15

Synthesis of Glycine-(3-carbobenzyloxyamido)propylamide:




20

N-BOC-Glycine-(3-carbobenzyloxyamido)propylamide (260 mg, 0.71 mmol) was dissolved in dichloromethane (3 mL) and trifluoroacetic acid (0.5 mL) added. After 20 minutes, the solution was evaporated in vacuo, dissolved in dichloromethane (2 mL) and evaporated in vacuo to obtain the crude product which was used directly in the next reaction. MS: m/e = 266 (M+H⁺).


25

Synthesis of 17a:




Di-isopropylethylamine (0.25 mL, 1.4 mmol) was added to a mixture of 1 (300 mg, 0.69 mmol) and HBTU (270 mg, 0.71 mmol) in dichloromethane (5 mL). Dimethylformamide (1 mL) was added to obtain a clear solution. After 30 minutes, a solution of Glycine-(3-carbobenzyloxyamido)propylamide (~0.71 mmol) and di-isopropylethylamine (0.25 mL, 1.4 mmol) in dichloromethane (1 mL) was added. The reaction mixture was extracted twice with 0.5 N HCl, once each with saturated aqueous NaCl, 1.0 N NaOH, saturated aqueous NaCl, and saturated aqueous NaHCO₃. The dichloromethane solution was filtered through a short column of Na₂SO₄ and evaporated *in vacuo* to obtain crude 17b (599 mg, 127%). MS: m/e = 681 (M+H⁺).

15 1:



Synthesis of 17b:




20

Trifluoroacetic acid (1 mL) was added to a solution of 17a (599 mg) in dichloromethane (5 mL) and allowed to stand at room temperature overnight. The solution was evaporated *in vacuo*, dissolved in

dichloromethane (2 mL) and evaporated *in vacuo* to obtain 3 (714 mg).
MS: $m/e = 625$ ($M+H^+$).


Synthesis of 17c:



5

A mixture of 3 (~0.3 g, ~ 0.5 mmol) and 10% Pd/C (25 mg) in ethanol (5 mL) was stirred under hydrogen (1 atm) for 2.5 hours. Disappearance of 17b was accompanied by the appearance of two peaks in the HPLC-MS chromatogram, both of which exhibited base peaks at $m/e = 491$ amu, consistent with ($M+H^+$) for the loss of carbobenzyloxy group from 26b. The reaction mixture was filtered through Celite and evaporated *in vacuo* to obtain 17c.


15 Synthesis of 17d:



A mixture of N,N-bis[2-bis(1,1'-dimethylethoxy)-2-oxoethyl]-amino]ethyl]glycine (487 mg, 0.788 mmol), HBTU (288 mg, 0.760 mmol) and di-isopropylethyl amine (0.4 mL, 2.3 mmol) in dimethylformamide (4 mL) was stirred at room temperature. A solution of 4 (~0.5 mmol) in dimethylformamide (2 mL) was added in one portion. After 2 hours, ~2/3 of the solution was removed, partitioned between dichloromethane and 0.5 M HCl. The organic phase extracted once with 0.5 M HCl, then with saturated aqueous NaCl, filtered through a column of Na₂SO₄, and evaporated *in vacuo* to obtain crude 17d. MS: m/e 1090 ($M+H^+$), 546 ($M+2H$)⁺

25

Synthesis of 17e




5

The remaining 1/3 of the reaction mixture 17d was treated with HBTU (75 mg, 0.20 mmol) and allowed to stir for 20 minutes. A solution prepared from hydroxylamine hydrochloride (50 mg, 0.70 mmol) and diisopropylethylamine (0.15 mL, 0.86 mmol) in dimethylformamide (0.5 mL) was added in one portion. The reaction mixture was partitioned between dichloromethane and 0.5 M HCl. The organic phase was extracted with saturated aqueous NaCl, then with saturated aqueous NaHCO₃. The NaHCO₃ phase was back-extracted with dichloromethane, the combined organic extracts filtered through a column of Na₂SO₄, and evaporated in vacuo. The crude product was purified by reverse-phase HPLC to obtain 14 mg of 27e. MS: m/e 1105 (M+H⁺), 553 (M+2H)²⁺

10

15

Synthesis of 17f:



20

A solution of 17e in trifluoroacetic acid (0.5 mL) and dichloromethane (2 mL) was allowed to stand at room temperature overnight. The solution was evaporated in vacuo, dissolved in acetonitrile-water and purified by reverse-phase HPLC to obtain 3.4 mg of 17f. MS: m/e 881 (M+H⁺), 441 (M+2H)²⁺

25

Utility

The pharmaceuticals of the present invention are useful for imaging of cardiovascular disease processes involving the degradation of the extracellular matrix. The
5 radiopharmaceuticals of the present invention comprised of a gamma emitting isotope are useful for imaging of cardiovascular pathological processes involving the degradation of the extracellular matrix, including atherosclerosis, congestive heart failure, and restenosis of blood vessels after
10 angioplasty.

The compounds of the present invention comprised of one or more paramagnetic metal ions selected from gadolinium, dysprosium, iron, and manganese, are useful as contrast agents for magnetic resonance imaging (MRI) of cardiovascular
15 pathological processes involving extracellular matrix degradation.

The compounds of the present invention comprised of one or more heavy atoms with atomic number of 20 or greater are useful as X-ray contrast agents for X-ray imaging of cardiovascular
20 pathological processes involving extracellular matrix degradation.

The compounds of the present invention comprised of an echogenic gas containing surfactant microsphere are useful as ultrasound contrast agents for sonography of cardiovascular
25 pathological processes involving extracellular matrix degradation.

Representative compounds of the present invention were tested in the one or more of the following in vitro assays and
30 were found to be active.

Matrix Metalloproteinase Assays for MMP-1 (collagenase-1), MMP-2 (gelatinase A), MMP-3 (stromelysin-1), MMP-8 (collagenase-2), MMP-9 (gelatinase B), MMP-13 (collagenase-3), MMP-14 (membrane
35 type 1 MMP), MMP-15 (membrane type 2 MMP), and MMP-16 (membrane type 3 MMP).

A. Reagents

1. MCA peptide substrate: Mca-Pro-Leu-Gly-Leu-Dpa-Ala-NH₂.
Peptide stocks are stored at -70 C in DMSO at 20 mM. Dilute
peptide in 1 x reaction buffer to a working concentration of 14
5 uM on day of use.

2. Enzyme buffer. 50 mM Tricine, 0.05% Brij-35, 400 mM NaCl, 10
mM CaCl₂, 0.02% NaN₃, pH 7.5.

10 3. Reaction buffer. 50 mM Tricine, 10 mM CaCl₂, 0.02% NaN₃, pH
7.5

4. Compounds. Stock compounds are at 10 mM in DMSO. Dilutions
are done in buffer.

15 5. Plates. microfluor W flat bottom plates (Dynex Inc.
Cat.#7905).

B. Assay

20 1. To 96 well fluorescent assay plates add 2 uL of DMSO control
or compound dilutions to wells.

2. Add 20 uL of EDTA (0.5M) to each quench well.

25 3. Add 50 uL of enzyme at the appropriate concentration.

4. Add 150 uL of the MCA peptide at final concentration of 10
uM.

30 5. Incubate each plate for 1 hour at room temperature on an
orbital shaker.

35 6. Add 20 uL of EDTA (0.5 M) to each test well to quench the
reaction.

7. Read each plate at 330 nm excitation, 440 nm emission (Dynx
plate reader).

8. Subtract each quench value from the corresponding test value.

9. % inhibition = $100 - (\text{sample fluorescence} / \text{control fluorescence}) \times 100$.

5

TACE Assay

A. Reagents

10 1. MCA Peptide substrate: Mca-PLAQAV(Dpa)RSSSR-NH₂. Peptide stocks are stored at -70 C in DMSO at 20 mM. Dilute peptide stock in reaction buffer to a working concentration of 20 uM on day of use.

15 2. Reaction buffer. 50 mM Tricine, 100 mM NaCl, 10 mM CaCl₂, 1 mM ZnCl₂, pH 7.5.

3. Compounds. Stock compounds are at 10 mM in DMSO

20 4. Plates. black Packard Optiplate (Cat.# HTRF-96)

5. Cytofluor Multi-well Plate Reader (Series 4000)

B. Assay

25

1. Initiate assay by adding 2 nM TACE to buffered solutions containing 10 μ M MCA peptide substrate in the presence of increasing concentrations of compound.

30 2. Add 20 uL of EDTA (0.5M) to each quench well.

3. Total volume is 300 uL in each well.

35 4. Incubate the reaction mixtures for 1 hour at 28 C on an orbital shaker.

5. Add 20 uL of EDTA (0.5M) to each test well to quench the reaction.

6. Read each plate at 330 nm excitation, 395 nm emission
7. Subtract each quench value from the corresponding test value.
- 5 8. % inhibition = $100 - (\text{sample fluorescence} / \text{control fluorescence}) \times 100$

In vivo Models

- 10 Cardiovascular disease models that can be used to assess the diagnostic radiopharmaceuticals, magnetic resonance, X-ray and ultrasound contrast agents of the present invention are reviewed in J. Nucl. Cardiol., 1998, 5, 167-83. There are several well established rabbit models of atherosclerosis; one
15 model produces predominantly proliferating smooth muscle cells by balloon deendothelialization of infradiaphragmatic abdominal aorta to simulate restenotic lesions; another model that produces simulated advanced human atherosclerotic plaque by balloon deendothelialization followed by a high cholesterol
20 diet.

- 25 A model of congestive heart failure is described in Am. J. Physiol., 1998, 274, H1516-23. In general, Yorkshire pigs are randomly assigned to undergo 3 wks of rapid atrial pacing at 240 beats/min. or to be sham controls. The pigs are chronically instrumented to measure left ventricular function in the conscious state. The pigs are anesthetized.

- 30 A shielded stimulating electrode is sutured onto the left atrium, connected to a modified programmable pace maker and buried in a subcutaneous pocket. The pericardium is closed loosely, the thoracotomy is closed, and the pleural space is evacuated of air. After a recovery period of 7-10 days, the pacemaker is activated in the animals selected to undergo chronic rapid pacing. The animals are sedated, the pacemaker is deactivated (pacing groups only. After a 30 min stabilization
35 period, indexes of LV function and geometry are determined (by echocardiography as a control) by injecting the radiolabeled compound. For biodistribution, the animals are anesthetized,

the heart extirpate and the LV apex and midventricular regions are evaluated.


A rat model of reversible coronary occlusion and reperfusion is described in McNulty et al., J. Am. Physiol.,
5 1996, H2283-9.

Obviously, numerous modifications and variations of the present invention are possible in light of the above teachings. It is therefore to be understood that within the scope of the
10 appended claims, the invention may be practiced otherwise that as specifically described herein.

WHAT IS CLAIMED IS:

1. A diagnostic agent comprising a diagnostic metal and a compound, wherein the compound comprises:
 - 5 iv) 1-10 targeting moieties;
 - v) a chelator; and
 - vi) 0-1 linking groups between the targeting moiety and chelator;wherein the targeting moiety is a matrix metalloproteinase
10 inhibitor; and
wherein the chelator is capable of conjugating to the diagnostic metal.
2. A diagnostic agent according to claim 1, wherein the
15 targeting moiety is a matrix metalloproteinase inhibitor having an inhibitory constant K_i of <1000 nM.
3. A diagnostic agent according to claim 1, wherein the
targeting moiety is a matrix metalloproteinase inhibitor having
20 an inhibitory constant K_i of <100 nM.
4. A diagnostic agent according to claim 1, comprising 1-5 targeting moieties.
- 25 5. A diagnostic agent according to claim 1, comprising one targeting moiety.
6. A diagnostic agent of claim 1, wherein the targeting moiety is an inhibitor of one or more matrix metalloproteinases
30 selected from the group consisting of MMP-1, MMP-2, MMP-3, MMP-9, and MMP-14.
7. A diagnostic agent of claim 6, wherein the targeting moiety is an inhibitor of one or more matrix metalloproteinases
35 selected from the group consisting of MMP-2, MMP-9, and MMP-14.

8. A diagnostic agent according to claim 1 wherein the targeting moiety is a matrix metalloproteinase inhibitor of the formulae (Ia) or (Ib):




R is independently OH or -CH₂SH;

10 R¹ is independently selected at each occurrence from the group:
H, OH, C₁₋₃ alkyl, C₂₋₃ alkenyl, C₂₋₃ alkynyl, and
heterocycle-S-CH₂-;

R² is independently C₁₋₂₀ alkyl;

15 X is independently C=O or SO₂, provided when X is C=O, R³ is



, and when X is SO₂, R³ is independently selected
from the group: aryl substituted with 0-2 R⁶, and
heterocycle substituted with 0-2 R⁶;

20 R⁴ is independently selected at each occurrence from the group:
C₁₋₆ alkyl, phenyl, and benzyl;

25 R⁵ is independently at each occurrence from the group: NH(C₁₋₆
alkyl), NH-phenyl, and NH-heterocycle; wherein said alkyl,
phenyl and heterocycle groups are optionally substituted
with a bond to the linking group or a bond to the chelator;

R⁶ is independently aryloxy substituted with 0-3 R⁷;

30 R⁷ is independently halogen or methoxy;

or alternatively,

5 R¹ and R⁴ may be taken together to form a bridging group of the formula -(CH₂)₃-O-phenyl-CH₂-, optionally substituted with a bond to the linking group or a bond to the chelator;

or alternatively,

10

R¹ and R² may be taken together to form a bridging group of the formula -(CH₂)₃-NH-, optionally substituted with a bond to the linking group or a bond to the chelator; or


15 R¹ and R² taken together with the nitrogen and carbon atom through which they are attached form a C₅₋₇ atom saturated ring system substituted with one or more substituents selected from the group consisting of: a bond to Ln, a bond to Ch, and -C(=O)-NR²⁹R³⁰;

20 R⁸ is independently at each occurrence OH or phenyl, optionally substituted with a bond to the linking group or a bond to the chelator, provided that when R⁸ is phenyl, R¹⁰ is -C(=O)-CR¹²-NH-CH(CH₃)-COOH;

25 R⁹ and R^{9'} are independently H, C₁₋₆ alkyl optionally substituted with a bond to the linking group or a bond to the chelator, or are taken together with the carbon atom to which R⁹ and R^{9'} are attached to form a 5-7 atom saturated, partially unsaturated or aromatic ring system containing 0-30 3 heteroatoms selected from O, N, SO₂ and S, said ring system substituted with R⁶ and optionally substituted with a bond to the linking group or a bond to the chelator;

- R¹⁰ and R¹¹ are independently H, or C₁₋₆ alkyl optionally substituted with a bond to the linking group or a bond to the chelator, or are taken together with the nitrogen atom to which they are attached to form a 5-7 atom saturated, partially unsaturated or aromatic ring system containing 0-3 heteroatoms selected from O, N, SO₂ and S, said ring system optionally substituted with 0-3 R²⁷, a bond to the linking group or a bond to the chelator;
- or alternatively,
- R⁹ and R¹⁰ are taken together with the carbon atom to which they are attached to form a 5-7 atom saturated, partially unsaturated or aromatic ring system containing 0-3 heteroatoms selected from O, N, SO₂ and S, said ring system optionally substituted with a bond to the linking group or a bond to the chelator; and
- R¹² is independently C₁₋₂₀ alkyl;
- R²⁷ is =O, C₁₋₄ alkyl, or phenyl substituted with R²⁸;
- R²⁸ is a phenoxy group substituted with 0-2 OCH₃ groups;
- R²⁹ and R³⁰ taken together with the nitrogen atom through which they are attached form a C₅₋₇ atom saturated ring system substituted with R³¹; and
- R³¹ is a benzyloxy group substituted with C₁₋₄ alkyl.

9. A diagnostic agent according to claim 8 wherein the targeting moiety is a matrix metalloproteinase inhibitor of the formulae (Ia) or (Ib):



R is OH;

R¹ is independently selected at each occurrence from the group:
5 H, OH, C₁₋₃ alkyl, C₂₋₃ alkenyl, C₂₋₃ alkynyl, and
heterocycle-S-CH₂-;

R² is independently C₁₋₆ alkyl;

10 X is C=O;

R⁴ is independently selected at each occurrence from the group:
C₁₋₆ alkyl, phenyl, and benzyl;

15 R⁵ is independently at each occurrence from the group: NH(C₁₋₆
alkyl), NH-phenyl, and NH-heterocycle; wherein said alkyl,
phenyl and heterocycle groups are optionally substituted
with a bond to the linking group or a bond to the chelator;

20 R⁶ is independently aryloxy substituted with 0-3 R⁷;

R⁷ is independently halogen or methoxy;

or alternatively,

25

R¹ and R⁴ may be taken together to form a bridging group of the
formula -(CH₂)₃-O-phenyl-CH₂-, optionally substituted with a
bond to the linking group or a bond to the chelator;

30 or alternatively,

R¹ and R² may be taken together to form a bridging group of the
formula -(CH₂)₃-NH-, optionally substituted with a bond to
the linking group or a bond to the chelator; or

R¹ and R² taken together with the nitrogen and carbon atom through which they are attached form a C₅₋₇ atom saturated ring system substituted with one or more substituents
5 selected from the group consisting of: a bond to Ln, a bond to Ch, and -C(=O)-NR²⁹R³⁰;

R⁸ is OH;

10 R⁹ and R^{9'} are independently H, C₁₋₆ alkyl optionally substituted with a bond to the linking group or a bond to the chelator, or are taken together with the carbon atom to which R⁹ and R^{9'} are attached to form a 5-7 atom saturated, partially unsaturated or aromatic ring system containing 0-
15 1 heteroatoms selected from O, N, , said ring system optionally substituted with a bond to the linking group or a bond to the chelator;

R¹⁰ and R¹¹ are independently H, or C₁₋₆ alkyl optionally
20 substituted with a bond to the linking group or a bond to the chelator, or are taken together with the nitrogen atom to which they are attached to form a 5-7 atom saturated, partially unsaturated or aromatic ring system containing 0-1 heteroatoms selected from O, N, , said ring system optionally substituted
25 with 0-3 R²⁷, a bond to the linking group or a bond to the chelator;

or alternatively,

30 R⁹ and R¹⁰ are taken together with the carbon atom to which they are attached to form a 5-7 atom saturated, partially unsaturated or aromatic ring system containing 0-1 heteroatoms selected from O, N, , said ring system

optionally substituted with a bond to the linking group or a bond to the chelator; and

R¹² is independently C₁₋₆ alkyl;

5 R²⁷ is =O, C₁₋₄ alkyl, or phenyl substituted with R²⁸;

R²⁸ is a phenoxy group substituted with 0-2 OCH₃ groups;

R²⁹ and R³⁰ taken together with the nitrogen atom through which they are attached form a C₅₋₇ atom saturated ring system substituted with R³¹; and

10 R³¹ is a benzyloxy group substituted with C₁₋₄ alkyl.


10. A diagnostic agent according to claim 8 wherein the targeting moiety is a matrix metalloproteinase inhibitor of the formulae (Ia) or (Ib):

15 wherein:

R is -OH;

R² is C₁₋₆ alkyl;

X is C=O;



20 R¹ and R⁴ are taken together to form a bridging group of formula -(CH₂)₃-O-phenyl-CH₂-;

R⁵ is NH(C₁₋₆alkyl), substituted with a bond to the linking group or a bond to the chelator.

25 11. A diagnostic agent according to claim 8, wherein:

R is -OH;

R⁹ is C₁ alkyl substituted with a bond to Ln;

R¹⁰ and R¹¹ taken together with the nitrogen atom to which they are attached form a 5 atom saturated ring system, said right

30 system is substituted with 0-3 R²⁷;

R²⁷ is =O, C₁₋₄ alkyl, or phenyl substituted with R²⁸; and

R²⁸ is a phenoxy group substituted with 0-2 OCH₃ groups.


12. A diagnostic agent according to claim 8 wherein the R is -OH;

5 R¹ and R² taken together with the nitrogen and carbon atom through which they are attached form a C₅₋₇ atom saturated ring system substituted with one or more substituents selected from the group consisting of: a bond to Ln, a bond to Ch, and -C(=O)-NR²⁹R³⁰;

10 R²⁹ and R³⁰ taken together with the nitrogen atom through which they are attached form a C₅₋₇ atom saturated ring system substituted with R³¹; and

R³¹ is a benzyloxy group substituted with C₁₋₄ alkyl.

15 13. A diagnostic agent according to claim 1 wherein the linking group is of the formula:



20 W¹ and W² are independently selected at each occurrence from the group: O, S, NH, NHC(=O), C(=O)NH, NR¹⁵C(=O), C(=O)NR¹⁵, C(=O), C(=O)O, OC(=O), NHC(=S)NH, NHC(=O)NH, SO₂, SO₂NH, - (OCH₂CH₂)₇₆₋₈₄, (OCH₂CH₂)_s, (CH₂CH₂O)_{s'}, (OCH₂CH₂CH₂)_{s''}, (CH₂CH₂CH₂O)_t, and (aa)_{t'};

25 aa is independently at each occurrence an amino acid;

Z is selected from the group: aryl substituted with 0-3 R¹⁶, C₃₋₁₀ cycloalkyl substituted with 0-3 R¹⁶, and a 5-10 membered heterocyclic ring system containing 1-4 heteroatoms independently selected from N, S, and O and substituted with 0-3 R¹⁶;

30

R¹³, R^{13a}, R¹⁴, R^{14a}, and R¹⁵ are independently selected at each occurrence from the group: H, =O, COOH, SO₃H, PO₃H, C₁-C₅ alkyl substituted with 0-3 R¹⁶, aryl substituted with 0-3 R¹⁶, benzyl substituted with 0-3 R¹⁶, and C₁-C₅ alkoxy substituted with 0-3 R¹⁶, NHC(=O)R¹⁷, C(=O)NHR¹⁷, NHC(=O)NHR¹⁷, NHR¹⁷, R¹⁷, and a bond to the chelator;

R¹⁶ is independently selected at each occurrence from the group: a bond to the chelator, COOR¹⁷, C(=O)NHR¹⁷, NHC(=O)R¹⁷, OH, NHR¹⁷, SO₃H, PO₃H, -OPO₃H₂, -OSO₃H, aryl substituted with 0-3 R¹⁷, C₁-5 alkyl substituted with 0-1 R¹⁸, C₁-5 alkoxy substituted with 0-1 R¹⁸, and a 5-10 membered heterocyclic ring system containing 1-4 heteroatoms independently selected from N, S, and O and substituted with 0-3 R¹⁷;

R¹⁷ is independently selected at each occurrence from the group: H, alkyl substituted with 0-1 R¹⁸, aryl substituted with 0-1 R¹⁸, a 5-10 membered heterocyclic ring system containing 1-4 heteroatoms independently selected from N, S, and O and substituted with 0-1 R¹⁸, C₃-10 cycloalkyl substituted with 0-1 R¹⁸, polyalkylene glycol substituted with 0-1 R¹⁸, carbohydrate substituted with 0-1 R¹⁸, cyclodextrin substituted with 0-1 R¹⁸, amino acid substituted with 0-1 R¹⁸, polycarboxyalkyl substituted with 0-1 R¹⁸, polyazaalkyl substituted with 0-1 R¹⁸, peptide substituted with 0-1 R¹⁸, wherein the peptide is comprised of 2-10 amino acids, 3,6-O-disulfo-B-D-galactopyranosyl, bis(phosphonomethyl)glycine, and a bond to the chelator;

R¹⁸ is a bond to the chelator;

k is selected from 0, 1, and 2;

h is selected from 0, 1, and 2;
 h' is selected from 0, 1, and 2;
 g is selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;
 g' is selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;
 5 s is selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;
 s' is selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;
 s" is selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;
 t is selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;
 t' is selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;
 10 x is selected from 0, 1, 2, 3, 4, and 5; and
 x' is selected from 0, 1, 2, 3, 4, and 5.

14. A diagnostic agent according to claim 13 wherein
 w¹ and w² are independently selected at each occurrence from
 15 the group: O, NH, NHC(=O), C(=O)NH, NR¹⁵C(=O), C(=O)NR¹⁵,
 C(=O), C(=O)O, OC(=O), NHC(=S)NH, NHC(=O)NH, SO₂, -
 (CH₂CH₂O)₇₆₋₈₄, (OCH₂CH₂)_s, (CH₂CH₂O)_{s'}, (OCH₂CH₂CH₂)_{s"},
 (CH₂CH₂CH₂O)_t, and (aa)_{t'};

20 aa is independently at each occurrence an amino acid;

Z is selected from the group: aryl substituted with 0-1 R¹⁶,
 C₃-10 cycloalkyl substituted with 0-1 R¹⁶, and a 5-10
 membered heterocyclic ring system containing 1-4
 25 heteroatoms independently selected from N, S, and O and
 substituted with 0-1 R¹⁶;

R¹³, R^{13a}, R¹⁴, R^{14a}, and R¹⁵ are independently selected at each
 occurrence from the group: H, =O, COOH, SO₃H, C₁-C₅ alkyl
 30 substituted with 0-1 R¹⁶, aryl substituted with 0-1 R¹⁶,
 benzyl substituted with 0-1 R¹⁶, and C₁-C₅ alkoxy
 substituted with 0-1 R¹⁶, NHC(=O)R¹⁷, C(=O)NHR¹⁷,
 NHC(=O)NHR¹⁷, NHR¹⁷, R¹⁷, and a bond to the chelator;

k is 0 or 1;
s is selected from 0, 1, 2, 3, 4, and 5;
s' is selected from 0, 1, 2, 3, 4, and 5;
s" is selected from 0, 1, 2, 3, 4, and 5; and
5 t is selected from 0, 1, 2, 3, 4, and 5.

15 A diagnostic agent according to claim 13 wherein
wherein:

W¹ is C(=O)NR¹⁵;
10 h is 1;
g is 3;
R¹³ and R¹⁴ are independently H;
x is 1;
k is 0;
15 g' is 0;
h' is 1;
W² is NH; and
x' is 1.

20 16. A diagnostic agent according to claim 13 wherein
x is 0;
k is 1;
Z is aryl substituted with 0-3 R¹⁶;
g' is 1;
25 W² is NH;
R^{13a} and R^{14a} are independently H;
h' is 1; and
x' is 1.

30 17. A diagnostic agent according to claim 13 wherein
W¹ is C(=O)NR¹⁵;
h is 1;
g is 2;
R¹³ and R¹⁴ are independently H;
35 x is 1;

- k is 0;
g' is 1;
R^{13a} and R^{14a} are independently H; or C1-5' alkyl substituted
with 0-3 R¹⁶;
5 R¹⁶ is SO₃H;
W² is NHC(=O) or NH;
h' is 1; and
x' is 2.
- 10 18. A diagnostic agent according to claim 13 wherein
W¹ is C(=O)NH;
h is 1;
g is 3;
R¹³ and R¹⁴ are independently H;
15 k is 0;
g' is 0;
x is 1;
W² is -NH(C=O)- or -(OCH₂CH₂)₇₆₋₈₄-;
h' is 2; and
20 x' is 1.
19. A diagnostic agent according to claim 13 wherein
x is 0;
k is 0;
25 g' is 3;
h' is 1;
W² is NH; and
x' is 1.
- 30 20. A diagnostic agent according to claim 13 wherein
x is 0;
Z is aryl substituted with 0-3 R¹⁶;
k is 1;
g' is 1;

R^{13} and R^{14} are independently H;

W^2 is $NHC(=O)$ or $-(OCH_2CH_2)_{76-84}-$; and

x' is 1.

5 21. A diagnostic agent according to claim 13 wherein

W^1 is $C=O$;

g is 2;

R^{13} and R^{14} are independently H;

k is 0;

10 g' is 0;


h' is 1;



W^2 is NH ; and

x' is 1.

15 22. A compound according to claim 1 wherein the linking group is absent.

20 23. A diagnostic agent according to claim 1 wherein the chelator is a metal bonding unit having a formula selected from the group:





5 A¹, A², A³, A⁴, A⁵, A⁶, A⁷, and A⁸ are independently selected at each occurrence from the group: N, NR²⁶, NR¹⁹, NR¹⁹R²⁰, S, SH, -S(Pg), O, OH, PR¹⁹, PR¹⁹R²⁰, -O-P(O)(R²¹)-O-, P(O)R²¹R²², a bond to the targeting moiety and a bond to the linking group;

10 Pg is a thiol protecting group;

15 E¹, E², E³, E⁴, E⁵, E⁶, E⁷, and E⁸ are independently a bond, CH, or a spacer group independently selected at each occurrence from the group: C₁-C₁₆ alkyl substituted with 0-3 R²³, aryl substituted with 0-3 R²³, C₃-₁₀ cycloalkyl substituted with 0-3 R²³, heterocyclo-C₁-₁₀ alkyl substituted with 0-3 R²³, wherein the heterocyclo group is a 5-10 membered heterocyclic ring system containing 1-4 heteroatoms independently selected from N, S, and O, C₆-₁₀ aryl-C₁-₁₀ alkyl substituted with 0-3 R²³, C₁-₁₀ alkyl-C₆-₁₀ aryl-substituted with 0-3 R²³, and a 5-10 membered heterocyclic

20

ring system containing 1-4 heteroatoms independently
selected from N, S, and O and substituted with 0-3 R²³;

R¹⁹ and R²⁰ are each independently selected from the group: a
5 bond to the linking group, a bond to the targeting moiety,
hydrogen, C₁-C₁₀ alkyl substituted with 0-3 R²³, aryl
substituted with 0-3 R²³, C₁-10 cycloalkyl substituted with
0-3 R²³, heterocyclo-C₁-10 alkyl substituted with 0-3 R²³,
wherein the heterocyclo group is a 5-10 membered
10 heterocyclic ring system containing 1-4 heteroatoms
independently selected from N, S, and O, C₆-10 aryl-C₁-10
alkyl substituted with 0-3 R²³, C₁-10 alkyl-C₆-10 aryl-
substituted with 0-3 R²³, a 5-10 membered heterocyclic ring
system containing 1-4 heteroatoms independently selected
15 from N, S, and O and substituted with 0-3 R²³, and an
electron, provided that when one of R¹⁹ or R²⁰ is an
electron, then the other is also an electron;

R²¹ and R²² are each independently selected from the group: a
20 bond to the linking group, a bond to the targeting moiety,
-OH, C₁-C₁₀ alkyl substituted with 0-3 R²³, C₁-C₁₀ alkyl
substituted with 0-3 R²³, aryl substituted with 0-3 R²³,
C₃-10 cycloalkyl substituted with 0-3 R²³,
heterocyclo-C₁-10 alkyl substituted with 0-3 R²³, wherein
25 the heterocyclo group is a 5-10 membered heterocyclic ring
system containing 1-4 heteroatoms independently selected
from N, S, and O, C₆-10 aryl-C₁-10 alkyl substituted with
0-3 R²³, C₁-10 alkyl-C₆-10 aryl- substituted with 0-3 R²³,
and a 5-10 membered heterocyclic ring system containing 1-4
30 heteroatoms independently selected from N, S, and O and
substituted with 0-3 R²³;

- R^{23} is independently selected at each occurrence from the group:
 a bond to the linking group, a bond to the targeting moiety, =O, F, Cl, Br, I, $-CF_3$, $-CN$, $-CO_2R^{24}$, $-C(=O)R^{24}$,
 $-C(=O)N(R^{24})_2$, $-CHO$, $-CH_2OR^{24}$, $-OC(=O)R^{24}$, $-OC(=O)OR^{24a}$,
 5 $-OR^{24}$, $-OC(=O)N(R^{24})_2$, $-NR^{25}C(=O)R^{24}$, $-NR^{25}C(=O)OR^{24a}$,
 $-NR^{25}C(=O)N(R^{24})_2$, $-NR^{25}SO_2N(R^{24})_2$, $-NR^{25}SO_2R^{24a}$, $-SO_3H$,
 $-SO_2R^{24a}$, $-SR^{24}$, $-S(=O)R^{24a}$, $-SO_2N(R^{24})_2$, $-N(R^{24})_2$,
 $-NHC(=S)NHR^{24}$, $=NOR^{24}$, NO_2 , $-C(=O)NHOR^{24}$, $-C(=O)NHN(R^{24})_2$,
 $-OCH_2CO_2H$, 2-(1-morpholino)ethoxy, C₁-C₅ alkyl, C₂-C₄
 10 alkenyl, C₃-C₆ cycloalkyl, C₃-C₆ cycloalkylmethyl, C₂-C₆
 alkoxyalkyl, aryl substituted with 0-2 R^{24} , and a 5-10
 membered heterocyclic ring system containing 1-4
 heteroatoms independently selected from N, S, and O; and
 wherein at least one of A^1 , A^2 , A^3 , A^4 , A^5 , A^6 , A^7 , A^8 or R^{23} is
 15 a bond to the linking group or targeting moiety;
 R^{24} , R^{24a} , and R^{25} are independently selected at each occurrence
 from the group: a bond to the linking group, a bond to the
 targeting moiety, H, C₁-C₆ alkyl, phenyl, benzyl, C₁-C₆ alkoxy,
 halide, nitro, cyano, and trifluoromethyl; and
 20 R^{26} is a co-ordinate bond to a metal or a hydrazine protecting
 group; or a pharmaceutically acceptable salt thereof.

24. A diagnostic agent according to claim 23 wherein:
 25 A^1 , A^2 , A^3 , A^4 , A^5 , A^6 , A^7 , and A^8 are independently selected at
 each occurrence from the group: NR^{19} , $NR^{19}R^{20}$, S, SH, OH,
 a bond to the targeting moiety and a bond to the linking
 group;
 30 E^1 , E^2 , E^3 , E^4 , E^5 , E^6 , E^7 , and E^8 are independently a bond,
 CH, or a spacer group independently selected at each
 occurrence from the group: C₁-C₁₀ alkyl substituted with
 0-3 R^{23} , aryl substituted with 0-3 R^{23} , C₃-10 cycloalkyl

substituted with 0-3 R^{23} , and a 5-10 membered heterocyclic ring system containing 1-4 heteroatoms independently selected from N, S, and O and substituted with 0-3 R^{23} ;

5 wherein at least one of A^1 , A^2 , A^3 , A^4 , A^5 , A^6 , A^7 , A^8 and R^{23} is a bond to the linking group or the targeting moiety;


R^{19} , and R^{20} are each independently selected from the group: a bond to the targeting moiety, a bond to the linking group,
10 hydrogen, C_1 - C_{10} alkyl substituted with 0-3 R^{23} , aryl substituted with 0-3 R^{23} , a 5-10 membered heterocyclic ring system containing 1-4 heteroatoms independently selected from N, S, and O and substituted with 0-3 R^{23} , and an
15 electron, provided that when one of R^{19} or R^{20} is an electron, then the other is also an electron;

R^{23} is independently selected at each occurrence from the group:
a bond to the targeting moiety, a bond to the linking group, =O, F, Cl, Br, I, $-CF_3$, $-CN$, $-CO_2R^{24}$, $-C(=O)R^{24}$,
20 $-C(=O)N(R^{24})_2$, $-CH_2OR^{24}$, $-OC(=O)R^{24}$, $-OC(=O)OR^{24a}$, $-OR^{24}$,
 $-OC(=O)N(R^{24})_2$, $-NR^{25}C(=O)R^{24}$, $-NR^{25}C(=O)OR^{24a}$,
 $-NR^{25}C(=O)N(R^{24})_2$, $-NR^{25}SO_2N(R^{24})_2$, $-NR^{25}SO_2R^{24a}$, $-SO_3H$,
 $-SO_2R^{24a}$, $-S(=O)R^{24a}$, $-SO_2N(R^{24})_2$, $-N(R^{24})_2$, $-NHC(=S)NHR^{24}$,
 $=NOR^{18}$, $-C(=O)NHN(R^{18})R^{18a}$, $-OCH_2CO_2H$, and
25 2-(1-morpholino)ethoxy; and

R^{24} , R^{24a} , and R^{25} are independently selected at each occurrence from the group: a bond to the linking group, H, and C_1 - C_6 alkyl.

30

25. A diagnostic agent according to claim 23 wherein the chelator is of the formula:



A¹ is a bond to the linking group;

5 A², A⁴, and A⁶ are each N;

A³, A⁵, A⁷ and A⁸ are each OH;

E¹, E², and E⁴ are C₂ alkyl;


10

E³, E⁵, E⁷, and E⁸ are C₂ alkyl substituted with 0-1 R²³;

R²³ is =O.

15 26. A diagnostic agent according to claim 23 wherein the chelator is of the formula:

Ch is



20 wherein:

A⁵ is a bond to Ln;

A¹, A³, A⁷ and A⁸ are each OH;


A², A⁴ and A⁶ are each NH;

E¹, E³, E⁵, E⁷, and E⁸ are C₂ alkyl substituted with 0-1 R²³;

25 E², and E⁴, are C₂ alkyl;

R²³ is =O.

27. A diagnostic agent according to claim 23 wherein the chelator is of the formula:



A¹, A², A³ and A⁴ are each N;

A⁵, A⁶ and A⁸ are each OH;

10

A⁷ is a bond to L_n;


E¹, E², E³, E⁴ are each independently C₂ alkyl; and

E⁵, E⁶, E⁷, E⁸ are each independently C₂ alkyl substituted with
0-1 R²³;

15

R²³ is =O.

28. A diagnostic agent according to claim 23 wherein the



20

A¹ is NR²⁶;

R²⁶ is a co-ordinate bond to a metal or a hydrazine protecting group;;

E^1 is a bond;

A² is NHR19;

5

R¹⁹ is a heterocycle substituted with R²³, the heterocycle being selected from pyridine and pyrimidine;


10 R^{23} is selected from a bond to the linking group, $C(=O)NHR^{24}$ and $C(=O)R^{24}$; and

10

R²⁴ is a bond to the linking group.

29. A diagnostic agent according to claim 23 wherein the
15 chelator is of the formula:

15



wherein:

A^1 and A^5 are each $-S(Pg)$;

Pg is a thiol protecting group;

20 E¹ and E⁴ are C₂ alkyl substituted with 0-1 R²³;

$$R^{23} \text{ is } = 0;$$


A^2 and A^4 are each $-NH$;

$$E^2 \text{ is } CH_2;$$

E^3 is C_{1-3} alkyl substituted with 0-1 R^{23} ;

25 A^3 is a bond to Ln.

30. A diagnostic agent according to claim 23 wherein the chelator is of the formula:




wherein:

- A¹ is a bond to Ln;
 E¹ is C₁ alkyl substituted by R²³;
 5 A² is NH;
 E² is C₂ alkyl substituted with 0-1R²³;
 A³ is -O-P(O)(R²¹)-O-;
 E³ is C₁ alkyl;
 A⁴ and A⁵ are each -O-;
 10 E⁴ and E⁶ are each independently C₁₋₁₆ alkyl substituted with 0-1R²³;
 E⁵ is C₁ alkyl;
 R²¹ is -OH; and
 R²³ is =O.


15

31. A diagnostic agent according to claim 1 having the formula:



20

wherein, Q is a compound of Formulae (Ia) or (Ib):




25 R is independently OH or -CH₂SH;

R¹ is independently selected at each occurrence from the group:
 H, OH, C₁₋₃ alkyl, C₂₋₃ alkenyl, C₂₋₃ alkynyl, and
 heterocycle-S-CH₂-;

R² is independently C₁₋₂₀ alkyl;

X is independently C=O or SO₂, provided when X is C=O, R³ is

5  , and when X is SO₂, R³ is independently selected from the group: aryl substituted with 0-2 R⁶, and heterocycle substituted with 0-2 R⁶;

10 R⁴ is independently selected at each occurrence from the group:
C₁₋₆ alkyl, phenyl, and benzyl;

R⁵ is independently at each occurrence from the group: NH(C₁₋₆ alkyl), NH-phenyl, and NH-heterocycle; wherein said alkyl, phenyl and heterocycle groups are optionally substituted
15 with a bond to L_n;

R⁶ is independently aryloxy substituted with 0-3 R⁷;

20 R⁷ is independently halogen or methoxy;
or alternatively,

R¹ and R⁴ may be taken together to form a bridging group of the formula -(CH₂)₃-O-phenyl-CH₂-, optionally substituted with a
25 bond to L_n;

or alternatively,

30 R¹ and R² may be taken together to form a bridging group of the formula -(CH₂)₃-NH-, optionally substituted with a bond to L_n; or

R¹ and R² taken together with the nitrogen and carbon atom through which they are attached form a C₅₋₇ atom saturated ring system substituted with one or more substituents selected from the group consisting of: a bond to L_n, a bond to Ch, and -C(=O)-NR²⁹R³⁰;

R⁸ is independently at each occurrence OH or phenyl, optionally substituted with a bond to L_n, provided that when R⁸ is phenyl, R¹⁰ is -C(=O)-CR¹²-NH-CH(CH₃)-COOH;

R⁹ and R^{9'} are independently H, C₁₋₆ alkyl optionally substituted with a bond to L_n, or are taken together with the carbon atom to which they are attached to form a 5-7 atom saturated, partially unsaturated or aromatic ring system containing 0-3 heteroatoms selected from O, N, SO₂ and S, said ring system substituted with R⁶ and optionally substituted with a bond to L_n;

R¹⁰ and R¹¹ are independently H, or C₁₋₆ alkyl optionally substituted with a bond to L_n, or are taken together with the nitrogen atom to which they are attached to form a 5-7 atom saturated, partially unsaturated or aromatic ring system containing 0-3 heteroatoms selected from O, N, SO₂ and S, said ring system optionally substituted with 0-3 R²⁷ or a bond to L_n;


or alternatively,

R⁹ and R¹⁰ are taken together with the carbon atom to which they are attached to form a 5-7 atom saturated, partially unsaturated or aromatic ring system containing 0-3 heteroatoms selected from O, N, SO₂ and S, said ring system optionally substituted with a bond to L_n;

R¹² is independently C₁₋₂₀ alkyl;

d is selected from 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;

5 L_n is a linking group having the formula:



W¹ and W² are independently selected at each occurrence from the
 10 group: O, S, NH, NHC(=O), C(=O)NH, NR¹⁵C(=O), C(=O)NR¹⁵,
 C(=O), C(=O)O, OC(=O), NHC(=S)NH, NHC(=O)NH, SO₂, SO₂NH, -
 (OCH₂CH₂)₇₆₋₈₄, (OCH₂CH₂)_S, (CH₂CH₂O)_{S'}, (OCH₂CH₂CH₂)_{S''},
 (CH₂CH₂CH₂O)_t, and (aa)_{t'};

15 aa is independently at each occurrence an amino acid;

Z is selected from the group: aryl substituted with 0-3 R¹⁶,
 C₃₋₁₀ cycloalkyl substituted with 0-3 R¹⁶, and a 5-10
 membered heterocyclic ring system containing 1-4
 20 heteroatoms independently selected from N, S, and O and
 substituted with 0-3 R¹⁶;

R¹³, R^{13a}, R¹⁴, R^{14a}, and R¹⁵ are independently selected at each
 occurrence from the group: H, =O, COOH, SO₃H, PO₃H, C₁₋₅
 25 alkyl substituted with 0-3 R¹⁶, aryl substituted with 0-3
 R¹⁶, benzyl substituted with 0-3 R¹⁶, and C₁₋₅ alkoxy
 substituted with 0-3 R¹⁶, NHC(=O)R¹⁷, C(=O)NHR¹⁷,
 NHC(=O)NHR¹⁷, NHR¹⁷, R¹⁷, and a bond to C_H;

30 R¹⁶ is independently selected at each occurrence from the group:
 a bond to C_H, COOR¹⁷, C(=O)NHR¹⁷, NHC(=O)R¹⁷, OH, NHR¹⁷,
 SO₃H, PO₃H, -OPO₃H₂, -OSO₃H, aryl substituted with 0-3 R¹⁷,

C₁₋₅ alkyl substituted with 0-1 R¹⁸, C₁₋₅ alkoxy substituted with 0-1 R¹⁸, and a 5-10 membered heterocyclic ring system containing 1-4 heteroatoms independently selected from N, S, and O and substituted with 0-3 R¹⁷;

5

R¹⁷ is independently selected at each occurrence from the group:

H, alkyl substituted with 0-1 R¹⁸, aryl substituted with 0-1 R¹⁸, a 5-10 membered heterocyclic ring system containing 1-4 heteroatoms independently selected from N,
10 S, and O and substituted with 0-1 R¹⁸, C₃₋₁₀ cycloalkyl substituted with 0-1 R¹⁸, polyalkylene glycol substituted with 0-1 R¹⁸, carbohydrate substituted with 0-1 R¹⁸, cyclodextrin substituted with 0-1 R¹⁸, amino acid substituted with 0-1 R¹⁸, polycarboxyalkyl substituted with
15 0-1 R¹⁸, polyazaalkyl substituted with 0-1 R¹⁸, peptide substituted with 0-1 R¹⁸, wherein the peptide is comprised of 2-10 amino acids, 3,6-O-disulfo-B-D-galactopyranosyl, bis(phosphonomethyl)glycine, and a bond to C_H;

20 R¹⁸ is a bond to C_H;

k is selected from 0, 1, and 2;

h is selected from 0, 1, and 2;

h' is selected from 0, 1, and 2;

25 g is selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;

g' is selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;

s is selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;

s' is selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;

s" is selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;


30 t is selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;

t' is selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;

x is selected from 0, 1, 2, 3, 4, and 5;

x' is selected from 0, 1, 2, 3, 4, and 5;

Ch is a metal bonding unit having a formula selected from the group:



A¹, A², A³, A⁴, A⁵, A⁶, A⁷, and A⁸ are independently selected at each occurrence from the group: N, NR²⁶, NR¹⁹, NR¹⁹R²⁰, S, SH, -S(Pg), O, OH, PR¹⁹, PR¹⁹R²⁰, -O-P(O)(R²¹)-O-,

P(O)R²¹R²², a bond to the targeting moiety and a bond to the linking group;

Pg is a thiol protecting group;

5

E¹, E², E³, E⁴, E⁵, E⁶, E⁷, and E⁸ are independently a bond, CH, or a spacer group independently selected at each occurrence from the group: C₁-C₁₆ alkyl substituted with 0-3 R²³, aryl substituted with 0-3 R²³, C₃-₁₀ cycloalkyl substituted with 0-3 R²³, heterocyclo-C₁-₁₀ alkyl substituted with 0-3 R²³, wherein the heterocyclo group is a 5-10 membered heterocyclic ring system containing 1-4 heteroatoms independently selected from N, S, and O, C₆-₁₀ aryl-C₁-₁₀ alkyl substituted with 0-3 R²³, C₁-₁₀ alkyl-C₆-₁₀ aryl-substituted with 0-3 R²³, and a 5-10 membered heterocyclic ring system containing 1-4 heteroatoms independently selected from N, S, and O and substituted with 0-3 R²³;

R¹⁹ and R²⁰ are each independently selected from the group: a bond to the linking group, a bond to the targeting moiety, hydrogen, C₁-C₁₀ alkyl substituted with 0-3 R²³, aryl substituted with 0-3 R²³, C₁-₁₀ cycloalkyl substituted with 0-3 R²³, heterocyclo-C₁-₁₀ alkyl substituted with 0-3 R²³, wherein the heterocyclo group is a 5-10 membered heterocyclic ring system containing 1-4 heteroatoms independently selected from N, S, and O, C₆-₁₀ aryl-C₁-₁₀ alkyl substituted with 0-3 R²³, C₁-₁₀ alkyl-C₆-₁₀ aryl-substituted with 0-3 R²³, a 5-10 membered heterocyclic ring system containing 1-4 heteroatoms independently selected from N, S, and O and substituted with 0-3 R²³, and an electron, provided that when one of R¹⁹ or R²⁰ is an electron, then the other is also an electron;

R²¹ and R²² are each independently selected from the group: a bond to the linking group, a bond to the targeting moiety, -OH, C₁-C₁₀ alkyl substituted with 0-3 R²³, C₁-C₁₀ alkyl substituted with 0-3 R²³, aryl substituted with 0-3 R²³, C₃-₁₀ cycloalkyl substituted with 0-3 R²³, heterocyclo-C₁-₁₀ alkyl substituted with 0-3 R²³, wherein the heterocyclo group is a 5-10 membered heterocyclic ring system containing 1-4 heteroatoms independently selected from N, S, and O, C₆-₁₀ aryl-C₁-₁₀ alkyl substituted with 0-3 R²³, C₁-₁₀ alkyl-C₆-₁₀ aryl- substituted with 0-3 R²³, and a 5-10 membered heterocyclic ring system containing 1-4 heteroatoms independently selected from N, S, and O and substituted with 0-3 R²³;

15

R²³ is independently selected at each occurrence from the group: a bond to the linking group, a bond to the targeting moiety, =O, F, Cl, Br, I, -CF₃, -CN, -CO₂R²⁴, -C(=O)R²⁴, -C(=O)N(R²⁴)₂, -CHO, -CH₂OR²⁴, -OC(=O)R²⁴, -OC(=O)OR^{24a}, -OR²⁴, -OC(=O)N(R²⁴)₂, -NR²⁵C(=O)R²⁴, -NR²⁵C(=O)OR^{24a}, -NR²⁵C(=O)N(R²⁴)₂, -NR²⁵SO₂N(R²⁴)₂, -NR²⁵SO₂R^{24a}, -SO₃H, -SO₂R^{24a}, -SR²⁴, -S(=O)R^{24a}, -SO₂N(R²⁴)₂, -N(R²⁴)₂, -NHC(=S)NHR²⁴, =NOR²⁴, NO₂, -C(=O)NHOR²⁴, -C(=O)NHN(R²⁴)₂, -OCH₂CO₂H, 2-(1-morpholino)ethoxy, C₁-C₅ alkyl, C₂-C₄ alkenyl, C₃-C₆ cycloalkyl, C₃-C₆ cycloalkylmethyl, C₂-C₆ alkoxyalkyl, aryl substituted with 0-2 R²⁴, and a 5-10 membered heterocyclic ring system containing 1-4 heteroatoms independently selected from N, S, and O; and wherein at least one of A¹, A², A³, A⁴, A⁵, A⁶, A⁷, A⁸ or R²³ is a bond to the linking group or targeting moiety;

30

R²⁴, R^{24a}, and R²⁵ are independently selected at each occurrence from the group: a bond to the linking group, a bond to the

targeting moiety, H, C₁-C₆ alkyl, phenyl, benzyl, C₁-C₆ alkoxy, halide, nitro, cyano, and trifluoromethyl; and

R²⁶ is a co-ordinate bond to a metal or a hydrazine protecting group; or

5 a pharmaceutically acceptable salt thereof.

32. A diagnostic agent according to Claim 31, wherein:

h' is 1;

10 w² is NH; and

x' is 1.

33. A diagnostic agent according to Claim 31, wherein:

x is 0;

15 Z is aryl substituted with 0-3 R¹⁶;

k is 1;

g' is 1;

R^{13a}R^{14a} are independently H;

w² is NHC(=O) or -(OCH₂CH₂)₇₆₋₈₄-; and

20 x' is 1.

34. A diagnostic agent according to Claim 31, wherein:

w¹ is C=O;

g is 2;

25 R¹³ and R¹⁴ are independently H;

k is 0;

g' is 0;

h' is 1;

w² is NH; and

30 x' is 1.

35. A diagnostic agent according to Claim 31, wherein:

2-{[5-(3-{2-[(6-Hydroxycarbamoyl-7-isobutyl-8-oxo-2-oxa-9-aza-bicyclo[10.2.2]hexadeca-1(15),12(16),13-triene-10-carbonyl)-

amino)-acetylamino)-propylcarbamoyl)-pyridin-2-yl]-
hydrazonomethyl}-benzenesulfonic acid;

2-{{5-(4-{{(6-Hydroxycarbamoyl-7-isobutyl-8-oxo-2-oxa-9-aza-
5 bicyclo[10.2.2]hexadeca-1(15),12(16),13-triene-10-carbonyl)-
amino)-methyl}-benzylcarbamoyl)-pyridin-2-yl]-hydrazonomethyl)-
benzenesulfonic acid;

2-[7-({N-[3-(2-{{7-(N-hydroxycarbamoyl)(3S,6R,7S)-4-aza-6-(2-
10 methylpropyl)-11-oxa-5-oxobicyclo[10.2.2]hexadeca-
1(15),12(16),13-trien-3-
yl]carbonylamino)acetylamino)propyl]carbamoyl)methyl)-1,4,7,10-
tetraaza-4,10-bis(carboxymethyl)cyclododecyl]acetic acid;

15 2-{7-[(N-{{4-({7-(N-hydroxycarbamoyl)(3S,6R,7S)-4-aza-6-(2-
methylpropyl)-11-oxa-5-oxobicyclo[10.2.2]hexadeca-
1(15),12(16),13-trien-3-yl]-
carbonylamino)methyl)phenyl]methyl]carbamoyl)methyl)-1,4,7,10-
tetraaza-4,10-bis(carboxymethyl)cyclododecyl]acetic acid;

20 2-(7-{{N-(1-{{N-[3-(2-{{7-(N-hydroxycarbamoyl)(3S,6R,7S)-4-aza-6-
(2-methylpropyl)-11-oxa-5-oxobicyclo[10.2.2]hexadeca-
1(15),12(16),13-trien-3-
yl]carbonylamino)acetylamino)propyl]carbamoyl}-2-
25 sulfoethyl)carbamoyl)methyl)-1,4,7,10-tetraaza-4,10-
bis(carboxymethyl)cyclododecyl]acetic acid;


2-[7-({N-[1-(N-{{4-({7-(N-hydroxycarbamoyl)(3S,6R,7S)-4-aza-6-
(2-methylpropyl)-11-oxa-5-oxobicyclo[10.2.2]hexadeca-
30 1(15),12(16),13-trien-3-yl]-
carbonylamino)methyl)phenyl]methyl]carbamoyl)-2-
sulfoethyl]carbamoyl)methyl)-1,4,7,10-tetraaza-4,10-
bis(carboxymethyl)cyclododecyl]acetic acid;

35 2-({2-[(N-[3-(2-{{7-(N-hydroxycarbamoyl)(3S,6R,7S)-4-aza-6-(2-
methylpropyl)-11-oxa-5-oxobicyclo[10.2.2]hexadeca-

- 1 (15), 12 (16), 13-trien-3-yl]carbonylamino}acetylamino)propyl]carbamoyl)methyl} (carboxymethyl) amino}ethyl) {2-[bis(carboxymethyl) amino]ethyl} amino}acetic acid;
- 5 2-[(2-[(N-{[4-([7-(N-hydroxycarbamoyl) (3S, 6R, 7S)-4-aza-6-(2-methylpropyl)-11-oxa-5-oxobicyclo[10.2.2]hexadeca-1 (15), 12 (16), 13-trien-3-yl]-carbonylamino)methyl}phenyl)methyl}carbamoyl)methyl} (carboxymethyl) amino}ethyl) {2-[bis(carboxymethyl) amino]ethyl} amino}acetic acid;
- 10 N-[3-(2-([7-(N-hydroxycarbamoyl) (3S, 6R, 7S)-4-aza-6-(2-methylpropyl)-11-oxa-5-oxobicyclo[10.2.2]hexadeca-1 (15), 12 (16), 13-trien-3-yl]carbonylamino}acetylamino)propyl]-4,5-bis[2-(ethoxyethylthio)acetylamino]pentanamide;
- 15 N-{[4-([7-(N-hydroxycarbamoyl) (3S, 6R, 7S)-4-aza-6-(2-methylpropyl)-11-oxa-5-oxobicyclo[10.2.2]hexadeca-1 (15), 12 (16), 13-trien-3-yl]carbonylamino)methyl}-phenyl)methyl}-4,5-bis[2-(ethoxyethylthio)acetylamino]-pentanamide;
- 20 1-(1,2-Dipalmitoyl-sn-glycero-3-phosphoethanolamino)- α , ω -dicarbonylPEG₃₄₀₀-2-([7-(N-hydroxycarbamoyl) (3S, 6R, 7S)-4-aza-6-(2-methylpropyl)-11-oxa-5-oxobicyclo[10.2.2]hexadeca-1 (15), 12 (16), 13-trien-3-yl]carbonylamino)-N-(3-aminopropyl)acetamide;
- 25 1-(1,2-Dipalmitoyl-sn-glycero-3-phosphoethanolamino)- α , ω -dicarbonylPEG₃₄₀₀-[7-(N-hydroxycarbamoyl) (3S, 6R, 7S)-4-aza-6-(2-methylpropyl)-11-oxa-5-oxobicyclo[10.2.2]hexadeca-1 (15), 12 (16), 13-trien-3-yl]-N-{[4-(aminomethyl)phenyl)methyl}carboxamide conjugate;
- 30 2-[2-([5-[N-(5-(N-hydroxycarbamoyl) (5R)-5-{3-[4-(3,4-dimethoxyphenoxy)phenyl]-3-methyl-2-
- 35

oxopyrrolidinyl}pentyl}carbamoyl}(2-pyridyl)}amino)(1Z)-2-azavinyl}benzenesulfonic acid;

2-(2-{[5-(N-{3-[3-(N-hydroxycarbamoyl)(4S)-4-({4-
5 methylphenyl)methoxy}piperidyl}carbonyl)piperidyl]-3-oxopropyl}carbamoyl)(2-pyridyl)}amino)(1Z)-2-azavinyl}benzenesulfonic acid; and



10

36. A diagnostic agent according to claim 1 wherein the diagnostic metal is selected from the group consisting of: a paramagnetic metal, a ferromagnetic metal, a gamma-emitting radioisotope, or an x-ray absorber.

15

37. A diagnostic agent according to claim 36 wherein the diagnostic metal is radioisotope selected from the group consisting of ^{99m}Tc , ^{95}Tc , ^{111}In , ^{62}Cu , ^{64}Cu , ^{67}Ga , and ^{68}Ga .

20 38. A diagnostic agent according to claim 37 further comprising a first ancillary ligand and a second ancillary ligand capable of stabilizing the radioisotope.

39. A diagnostic agent according to Claim 37, wherein the
25 radioisotope is ^{99m}Tc .

40. A diagnostic agent according to Claim 37, wherein the radioisotope is ^{111}In .

41. A diagnostic agent according to claim 36 wherein the paramagnetic metal ion is selected from the group consisting of Gd(III), Dy(III), Fe(III), and Mn(II).
- 5 42. A diagnostic agent according to claim 36 wherein the x-ray absorber is a metal is selected from the group consisting of: Re, Sm, Ho, Lu, Pm, Y, Bi, Pd, Gd, La, Au, Au, Yb, Dy, Cu, Rh, Ag, and Ir.
- 10 43. A diagnostic composition comprising a compound according to claim 1 or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier.
44. A kit comprising a compound of Claim 1, or a
15 pharmaceutically acceptable salt form thereof and a pharmaceutically acceptable carrier.
45. A kit according to Claim 44, wherein the kit further comprises one or more ancillary ligands and a reducing agent.
- 20 46. A kit according to Claim 45, wherein the ancillary ligands are tricine and TPPTS.
- 47 A kit according to Claim 45, wherein the reducing agent is
25 tin(II).
48. A diagnostic agent comprising an echogenic gas and a compound, wherein the compound comprises:
- iv) 1-10 targeting moieties;
- 30 v) a surfactant (Sf); and
- vi) 0-1 linking groups between the targeting moiety and surfactant;
- wherein the targeting moiety is a matrix metalloproteinase inhibitor; and
- 35 wherein the surfactant is capable of forming an echogenic gas filled lipid sphere or microbubble.

49. A diagnostic agent according to claim 48, wherein the targeting moiety is a matrix metalloproteinase inhibitor having an inhibitory constant K_i of <1000 nM.

5

50. A diagnostic agent according to claim 48, wherein the targeting moiety is a matrix metalloproteinase inhibitor having an inhibitory constant K_i of <100 nM.

10 51. A diagnostic agent according to claim 48, comprising 1-5 targeting moieties.

52. A diagnostic agent according to claim 48, comprising one targeting moiety.

15


53. A diagnostic agent according to claim 48, wherein the targeting moiety is an inhibitor of one or more matrix metalloproteinases selected from the group consisting of MMP-1, MMP-2, MMP-3, MMP-9, and MMP-14.

20

54. A diagnostic agent according to claim 48, wherein the targeting moiety is an inhibitor of one or more matrix metalloproteinases selected from the group consisting of MMP-2, MMP-9, and MMP-14.

25

55. A diagnostic agent according to claim 48, wherein the targeting moiety is of the formulae (Ia) or (Ib):




30

R is independently OH or $-\text{CH}_2\text{SH}$;

R¹ is independently selected at each occurrence from the group:
H, OH, C₁₋₃ alkyl, C₂₋₃ alkenyl, C₂₋₃ alkynyl, and
heterocycle-S-CH₂-;

5 R² is independently C₁₋₂₀ alkyl;

X is independently C=O or SO₂, provided when X is C=O, R³ is



, and when X is SO₂, R³ is independently selected
from the group: aryl substituted with 0-2 R⁶, and
10 heterocycle substituted with 0-2 R⁶;

R⁴ is independently selected at each occurrence from the group:
C₁₋₆ alkyl, phenyl, and benzyl;

15 R⁵ is independently at each occurrence from the group: NH(C₁₋₆
alkyl), NH-phenyl, and NH-heterocycle; wherein said alkyl,
phenyl and heterocycle groups are optionally substituted
with a bond to the linking group or a bond to the
surfactant;

20

R⁶ is independently aryloxy substituted with 0-3 R⁷;

R⁷ is independently halogen or methoxy;

25 or alternatively,

R¹ and R⁴ may be taken together to form a bridging group of the
formula -(CH₂)₃-O-phenyl-CH₂-, optionally substituted with a
bond to the linking group or a bond to the surfactant;

30

or alternatively,

R¹ and R² may be taken together to form a bridging group of the formula $-(CH_2)_3-NH-$, optionally substituted with a bond to the linking group or a bond to the surfactant; or

5 R¹ and R² taken together with the nitrogen and carbon atom through which they are attached form a C₅₋₇ atom saturated ring system substituted with one or more substituents selected from the group consisting of: a bond to Ln, a bond to Sf, and $-C(=O)-NR^{29}R^{30}$;

10

R⁸ is independently at each occurrence OH or phenyl, optionally substituted with a bond to the linking group or a bond to the surfactant, provided that when R⁸ is phenyl, R¹⁰ is $-C(=O)-CR^{12}-NH-CH(CH_3)-COOH$;

15

R⁹ and R^{9'} are independently H, C₁₋₆ alkyl optionally substituted with a bond to the linking group or a bond to the surfactant, or are taken together with the carbon atom to which R⁹ and R^{9'} are attached to form a 5-7 atom saturated, partially unsaturated or aromatic ring system containing 0-3 heteroatoms selected from O, N, SO₂ and S, said ring system substituted with R⁶ and optionally substituted with a bond to the linking group or a bond to the surfactant;

25


R¹⁰ and R¹¹ are independently H, or C₁₋₆ alkyl optionally substituted with a bond to the linking group or a bond to the surfactant, or are taken together with the nitrogen atom to which they are attached to form a 5-7 atom saturated, partially unsaturated or aromatic ring system containing 0-3 heteroatoms selected from O, N, SO₂ and S, said ring system optionally substituted with 0-3 R²⁷, a bond to the linking group or a bond to the surfactant;

30

or alternatively,

- R^9 and R^{10} are taken together with the carbon atom to which they are attached to form a 5-7 atom saturated, partially unsaturated or aromatic ring system containing 0-3 heteroatoms selected from O, N, SO_2 and S, said ring system optionally substituted with a bond to the linking group or a bond to the surfactant; and
- R^{12} is independently C₁-20 alkyl;
- R^{27} is =O, C₁-4 alkyl, or phenyl substituted with R^{28} ;
- R^{28} is a phenoxy group substituted with 0-2 OCH_3 groups;
- R^{29} and R^{30} taken together with the nitrogen atom through which they are attached form a C₅-7 atom saturated ring system substituted with R^{31} ; and
- R^{31} is a benzyloxy group substituted with C₁-4 alkyl.

56. A diagnostic agent according to claim 55 wherein wherein the targeting moiety is a matrix metalloproteinase inhibitor of the formulae (Ia) or (Ib):



- R is OH;

R^1 is independently selected at each occurrence from the group:
H, OH, C₁-3 alkyl, C₂-3 alkenyl, C₂-3 alkynyl, and heterocycle-S-CH₂-;

30

R^2 is independently C₁-6 alkyl;

X is C=O;

R⁴ is independently selected at each occurrence from the group:
C₁₋₆ alkyl, phenyl, and benzyl;

5

R⁵ is independently at each occurrence from the group: NH(C₁₋₆ alkyl), NH-phenyl, and NH-heterocycle; wherein said alkyl, phenyl and heterocycle groups are optionally substituted with a bond to the linking group or a bond to the
10 surfactant;

R⁶ is independently aryloxy substituted with 0-3 R⁷;

R⁷ is independently halogen or methoxy;

15

or alternatively,

R¹ and R⁴ may be taken together to form a bridging group of the
20 formula -(CH₂)₃-O-phenyl-CH₂-, optionally substituted with a
bond to the linking group or a bond to the surfactant;

or alternatively,

R¹ and R² may be taken together to form a bridging group of the
25 formula -(CH₂)₃-NH-, optionally substituted with a bond to
the linking group or a bond to the surfactant; or

R¹ and R² taken together with the nitrogen and carbon atom
through which they are attached form a C₅₋₇ atom saturated
30 ring system substituted with one or more substituents
selected from the group consisting of: a bond to Ln, a bond
to Sf, and -C(=O)-NR²⁹R³⁰;

R⁸ is OH;

R⁹ and R^{9'} are independently H, C₁₋₆ alkyl optionally substituted with a bond to the linking group or a bond to the surfactant, or are taken together with the carbon atom to which R⁹ and R^{9'} are attached to form a 5-7 atom saturated, partially unsaturated or aromatic ring system containing 0-1 heteroatoms selected from O, N, , said ring system optionally substituted with a bond to the linking group or a bond to the surfactant;

10

R¹⁰ and R¹¹ are independently H, or C₁₋₆ alkyl optionally substituted with a bond to the linking group or a bond to the surfactant, or are taken together with the nitrogen atom to which they are attached to form a 5-7 atom saturated, partially unsaturated or aromatic ring system containing 0-1 heteroatoms selected from O, N, , said ring system optionally substituted with 0-3 R²⁷, a bond to the linking group or a bond to the surfactant;

20 or alternatively,

R⁹ and R¹⁰ are taken together with the carbon atom to which they are attached to form a 5-7 atom saturated, partially unsaturated or aromatic ring system containing 0-1 heteroatoms selected from O, N, , said ring system optionally substituted with a bond to the linking group or a bond to the surfactant; and

R¹² is independently C₁₋₆ alkyl;

30 R²⁷ is =O, C₁₋₄ alkyl, or phenyl substituted with R²⁸;

R²⁸ is a phenoxy group substituted with 0-2 OCH₃ groups;

R²⁹ and R³⁰ taken together with the nitrogen atom through which they are attached form a C₅₋₇ atom saturated ring system substituted with R³¹; and

R³¹ is a benzyloxy group substituted with C1-4 alkyl.


57. A diagnostic agent according to claim 55 wherein the
5 targeting moiety is a matrix metalloproteinase inhibitor of the
formulae (Ia) or (Ib):

wherein:

R is -OH;

R² is C₁₋₆ alkyl;

10 X is C=O;



R¹ and R⁴ are taken together to form a bridging group of formula
-(CH₂)₃-O-phenyl-CH₂-;

R⁵ is NH(C1-6alkyl), substituted with a bond to the linking
15 group or a bond to the surfactant.

58. A diagnostic agent according to claim 55 wherein:

R is -OH;

R⁹ is C₁ alkyl substituted with a bond to Ln;

20 R¹⁰ and R¹¹ taken together with the nitrogen atom to which they
are attached form a 5 atom saturated ring system, said right
system is substituted with 0-3 R²⁷;

R²⁷ is =O, C1-4 alkyl, or phenyl substituted with R²⁸; and

R²⁸ is a phenoxy group substituted with 0-2 OCH₃ groups.

25

59. A diagnostic agent according to claim 55 wherein the

R is -OH;

R¹ and R² taken together with the nitrogen and carbon atom
through which they are attached form a C₅₋₇ atom saturated ring

30 system substituted with one or more substituents selected from

the group consisting of: a bond to Ln, a bond to Sf, and -C(=O)-NR²⁹R³⁰;

R²⁹ and R³⁰ taken together with the nitrogen atom through which they are attached form a C5-7 atom saturated ring system

5 substituted with R³¹; and

R³¹ is a benzyloxy group substituted with C1-4 alkyl.

60. A diagnostic agent according to claim 48 wherein the linking group is of the formula:

10

$$((W^1)_h - (CR^{13}R^{14})_g)_x - (Z)_k - ((CR^{13a}R^{14a})_{g'} - (W^2)_{h'})_{x'};$$

W¹ and W² are independently selected at each occurrence from the

group: O, S, NH, NHC(=O), C(=O)NH, NR¹⁵C(=O), C(=O)NR¹⁵,
 15 C(=O), C(=O)O, OC(=O), NHC(=S)NH, NHC(=O)NH, SO₂, SO₂NH, -
 (OCH₂CH₂)₇₆₋₈₄, (OCH₂CH₂)_s, (CH₂CH₂O)_{s'}, (OCH₂CH₂CH₂)_{s''},
 (CH₂CH₂CH₂O)_t, and (aa)_{t'};

aa is independently at each occurrence an amino acid;

20

Z is selected from the group: aryl substituted with 0-3 R¹⁶,
 C₃-10 cycloalkyl substituted with 0-3 R¹⁶, and a 5-10
 membered heterocyclic ring system containing 1-4
 heteroatoms independently selected from N, S, and O and
 25 substituted with 0-3 R¹⁶;

R¹³, R^{13a}, R¹⁴, R^{14a}, and R¹⁵ are independently selected at each
 occurrence from the group: H, =O, COOH, SO₃H, PO₃H, C₁-C₅
 alkyl substituted with 0-3 R¹⁶, aryl substituted with 0-3
 30 R¹⁶, benzyl substituted with 0-3 R¹⁶, and C₁-C₅ alkoxy
 substituted with 0-3 R¹⁶, NHC(=O)R¹⁷, C(=O)NHR¹⁷,
 NHC(=O)NHR¹⁷, NHR¹⁷, R¹⁷, and a bond to the surfactant;

- R^{16} is independently selected at each occurrence from the group:
 a bond to the surfactant, $COOR^{17}$, $C(=O)NHR^{17}$, $NHC(=O)R^{17}$,
 OH, NHR^{17} , SO_3H , PO_3H , $-OPO_3H_2$, $-OSO_3H$, aryl substituted
 with 0-3 R^{17} , C_{1-5} alkyl substituted with 0-1 R^{18} , C_{1-5}
 5 alkoxy substituted with 0-1 R^{18} , and a 5-10 membered
 heterocyclic ring system containing 1-4 heteroatoms
 independently selected from N, S, and O and substituted
 with 0-3 R^{17} ;
- 10 R^{17} is independently selected at each occurrence from the group:
 H, alkyl substituted with 0-1 R^{18} , aryl substituted with
 0-1 R^{18} , a 5-10 membered heterocyclic ring system
 containing 1-4 heteroatoms independently selected from N,
 S, and O and substituted with 0-1 R^{18} , C_{3-10} cycloalkyl
 15 substituted with 0-1 R^{18} , polyalkylene glycol substituted
 with 0-1 R^{18} , carbohydrate substituted with 0-1 R^{18} ,
 cyclodextrin substituted with 0-1 R^{18} , amino acid
 substituted with 0-1 R^{18} , polycarboxyalkyl substituted with
 0-1 R^{18} , polyazaalkyl substituted with 0-1 R^{18} , peptide
 20 substituted with 0-1 R^{18} , wherein the peptide is comprised
 of 2-10 amino acids, 3,6-O-disulfo-B-D-galactopyranosyl,
 bis(phosphonomethyl)glycine, and a bond to the surfactant;

- R^{18} is a bond to the surfactant;
- 25 k is selected from 0, 1, and 2;
 h is selected from 0, 1, and 2;
 h' is selected from 0, 1, and 2;
 g is selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;
 30 g' is selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;
 s is selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;
 s' is selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;
 s'' is selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;

t is selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;
 t' is selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;
 x is selected from 0, 1, 2, 3, 4, and 5; and
 x' is selected from 0, 1, 2, 3, 4, and 5.

5

61. A diagnostic agent according to claim 60 wherein
 W¹ and W² are independently selected at each occurrence from
 the group: O, NH, NHC(=O), C(=O)NH, NR¹⁵C(=O), C(=O)NR¹⁵,
 C(=O), C(=O)O, OC(=O), NHC(=S)NH, NHC(=O)NH, SO₂, -
 10 (CH₂CH₂O)₇₆₋₈₄, (OCH₂CH₂)_s, (CH₂CH₂O)_{s'}, (OCH₂CH₂CH₂)_{s"},
 (CH₂CH₂CH₂O)_t, and (aa)_{t'};

aa is independently at each occurrence an amino acid;

15 Z is selected from the group: aryl substituted with 0-1 R¹⁶,
 C₃₋₁₀ cycloalkyl substituted with 0-1 R¹⁶, and a 5-10
 membered heterocyclic ring system containing 1-4
 heteroatoms independently selected from N, S, and O and
 substituted with 0-1 R¹⁶;

20

R¹³, R^{13a}, R¹⁴, R^{14a}, and R¹⁵ are independently selected at each
 occurrence from the group: H, =O, COOH, SO₃H, C₁-C₅ alkyl
 substituted with 0-1 R¹⁶, aryl substituted with 0-1 R¹⁶,
 benzyl substituted with 0-1 R¹⁶, and C₁-C₅ alkoxy
 25 substituted with 0-1 R¹⁶, NHC(=O)R¹⁷, C(=O)NHR¹⁷,
 NHC(=O)NHR¹⁷, NHR¹⁷, R¹⁷, and a bond to the surfactant;

k is 0 or 1;

s is selected from 0, 1, 2, 3, 4, and 5;

30 s' is selected from 0, 1, 2, 3, 4, and 5;

s" is selected from 0, 1, 2, 3, 4, and 5; and

t is selected from 0, 1, 2, 3, 4, and 5.

62. A diagnostic agent according to claim 60

wherein:

W¹ is C(=O)NR¹⁵;

h is 1;

g is 3;

5 R¹³ and R¹⁴ are independently H;

x is 1;

k is 0;

g' is 0;

h' is 1;

10 W² is NH; and

x' is 1.

63. A diagnostic agent according to claim 60

x is 0;

15 k is 1;

Z is aryl substituted with 0-3 R¹⁶;

g[^] is 1;

W² is NH;

R^{13a} and R^{14a} are independently H;

20 h' is 1; and

x' is 1.

64. A diagnostic agent according to claim 60

W¹ is C(=O)NR¹⁵;

25 h is 1;

g is 2;

R¹³ and R¹⁴ are independently H;

x is 1;

k is 0;

30 g' is 1;

R^{13a} and R^{14a} are independently H; or C1-5 alkyl substituted
with 0-3 R¹⁶;

R¹⁶ is SO₃H;

W² is NHC(=O) or NH;

h' is 1; and
x' is 2.

65. A diagnostic agent according to claim 60
5 W¹ is C(=O)NH;
h is 1;
g is 3;
R¹³ and R¹⁴ are independently H;
k is 0;
10 g' is 0;
x is 1;
W² is -NH(C=O)- or -(OCH₂CH₂)₇₆₋₈₄-;
h' is 2; and
x' is 1.

15 66. A diagnostic agent according to claim 60
x is 0;
k is 0;
g' is 3;
20 h' is 1;
W² is NH; and
x' is 1.

67. A diagnostic agent according to claim 60
25 x is 0;
Z is aryl substituted with 0-3 R¹⁶;
k is 1;
g' is 1;
R^{13a}R^{14a} are independently H;
30 W² is NHC(=O) or -(OCH₂CH₂)₇₆₋₈₄-; and
x' is 1.

68. A diagnostic agent according to claim 60
W¹ is C=O;
35 g is 2;

R¹³ and R¹⁴ are independently H;

k is 0;

g' is 0;

h' is 1;

5 W² is NH; and

x' is 1.

69. A diagnostic agent according to claim 48 wherein the linking group is present.

10

70. A diagnostic agent according to claim 48 wherein

S_f is a surfactant which is a lipid or a compound of the

15 formula:
$$\begin{array}{c} \text{E}^9 - \text{A}^{10} \\ \diagup \\ \text{A}^9 \end{array} ;$$

A⁹ is selected from the group: OH and OR³²;

A¹⁰ is OR³²;

20

R³² is C(=O)C₁₋₂₀ alkyl;

E⁹ is C₁₋₁₀ alkylene substituted with 1-3 R³³;

25 R³³ is independently selected at each occurrence from the group:

R³⁵, -PO₃H-R³⁵, =O, -CO₂R³⁴, -C(=O)R³⁴, -C(=O)N(R³⁴)₂,
-CH₂OR³⁴, -OR³⁴, -N(R³⁴)₂, C₁-C₅ alkyl, and C₂-C₄ alkenyl;


R³⁴ is independently selected at each occurrence from the group:

30 R³⁵, H, C₁-C₆ alkyl, phenyl, benzyl, and trifluoromethyl;

R³⁵ is a bond to L_n;

and a pharmaceutically acceptable salt thereof.

- 5 71. A diagnostic agent according to claim 48 wherein the surfactant is a lipid or a compound of the



10 A^9 is OR^{32} ;

A^{10} is OR^{32} ;

R^{32} is $\text{C}(=\text{O})\text{C}_{1-15}$ alkyl;

15

E^9 is C_{1-4} alkylene substituted with 1-3 R^{33} ;

R^{33} is independently selected at each occurrence from the group:


20 R^{35} , $-\text{PO}_3\text{H}-\text{R}^{35}$, $=\text{O}$, $-\text{CO}_2\text{R}^{34}$, $-\text{C}(=\text{O})\text{R}^{34}$, $-\text{CH}_2\text{OR}^{34}$, $-\text{OR}^{34}$,
and C_1-C_5 alkyl;

R^{34} is independently selected at each occurrence from the group:

R^{35} , H , C_1-C_6 alkyl, phenyl, and benzyl; and

25 R^{35} is a bond to L_n .

72. A diagnostic agent according to claim 48, wherein




wherein:

- A¹ is a bond to Ln;
 E¹ is C₁ alkyl substituted by R²³;
 5 A² is NH;
 E² is C₂ alkyl substituted with 0-1R²³;
 A³ is -O-P(O)(R²¹)-O-;
 E³ is C₁ alkyl;
 A⁴ and A⁵ are each -O-;
 10 E⁴ and E⁶ are each independently C₁₋₁₆ alkyl substituted with 0-1R²³;
 E⁵ is C₁ alkyl;
 A⁵ is -O-;
 R²¹ is -OH; and
 15 R²³ is =O.

73. A diagnostic agent according to claim 48 wherein the compound is of the formula:



wherein, Q is a compound of Formulae (Ia) or (Ib):




25

R is independently OH or -CH₂SH;

R¹ is independently selected at each occurrence from the group:
H, OH, C₁₋₃ alkyl, C₂₋₃ alkenyl, C₂₋₃ alkynyl, and
heterocycle-S-CH₂-;

5 R² is independently C₁₋₂₀ alkyl;

X is independently C=O or SO₂, provided when X is C=O, R³ is



, and when X is SO₂, R³ is independently selected
from the group: aryl substituted with 0-2 R⁶, and
10 heterocycle substituted with 0-2 R⁶;

R⁴ is independently selected at each occurrence from the group:
C₁₋₆ alkyl, phenyl, and benzyl;

15 R⁵ is independently at each occurrence from the group: NH(C₁₋₆
alkyl), NH-phenyl, and NH-heterocycle; wherein said alkyl,
phenyl and heterocycle groups are optionally substituted
with a bond to L_n;

20 R⁶ is independently aryloxy substituted with 0-3 R⁷;

R⁷ is independently halogen or methoxy;

or alternatively,

25

R¹ and R⁴ may be taken together to form a bridging group of the
formula -(CH₂)₃-O-phenyl-CH₂-, optionally substituted with a
bond to L_n;

30 or alternatively,

R¹ and R² may be taken together to form a bridging group of the formula $-(CH_2)_3-NH-$, optionally substituted with a bond to L_n; or

- 5 R¹ and R² taken together with the nitrogen and carbon atom through which they are attached form a C₅₋₇ atom saturated ring system substituted with one or more substituents selected from the group consisting of: a bond to L_n, a bond to Sf, and $-C(=O)-NR^{29}R^{30}$;

10

R⁸ is independently at each occurrence OH or phenyl, optionally substituted with a bond to L_n, provided that when R⁸ is phenyl, R¹⁰ is $-C(=O)-CR^{12}-NH-CH(CH_3)-COOH$;

- 15 R⁹ and R^{9'} are independently H, C₁₋₆ alkyl optionally substituted with a bond to L_n, or are taken together with the carbon atom to which they are attached to form a 5-7 atom saturated, partially unsaturated or aromatic ring system containing 0-3 heteroatoms selected from O, N, SO₂ and S, said ring system substituted with R⁶ and optionally substituted with a bond to L_n;

20

R¹⁰ and R¹¹ are independently H, or C₁₋₆ alkyl optionally substituted with a bond to L_n, or are taken together with the nitrogen atom to which they are attached to form a 5-7 atom saturated, partially unsaturated or aromatic ring system containing 0-3 heteroatoms selected from O, N, SO₂ and S, said ring system optionally substituted with 0-3 R²⁷ or a bond to L_n;

25

30

or alternatively,

R⁹ and R¹⁰ are taken together with the carbon atom to which they are attached to form a 5-7 atom saturated, partially

unsaturated or aromatic ring system containing 0-3 heteroatoms selected from O, N, SO₂ and S, said ring system optionally substituted with a bond to L_n;

5 R¹² is independently C₁₋₂₀ alkyl;

d is selected from 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;

L_n is a linking group having the formula:

10

$$((W^1)_h - (CR^{13}R^{14})_g)_x - (Z)_k - ((CR^{13a}R^{14a})_{g'} - (W^2)_{h'})_{x'};$$

W¹ and W² are independently selected at each occurrence from the

group: O, S, NH, NHC(=O), C(=O)NH, NR¹⁵C(=O), C(=O)NR¹⁵,
 15 C(=O), C(=O)O, OC(=O), NHC(=S)NH, NHC(=O)NH, SO₂, SO₂NH, -
 (OCH₂CH₂)₇₆₋₈₄, (OCH₂CH₂)_s, (CH₂CH₂O)_{s'}, (OCH₂CH₂CH₂)_{s''},
 (CH₂CH₂CH₂O)_t, and (aa)_{t'};

aa is independently at each occurrence an amino acid;

20

Z is selected from the group: aryl substituted with 0-3 R¹⁶,
 C₃₋₁₀ cycloalkyl substituted with 0-3 R¹⁶, and a 5-10
 membered heterocyclic ring system containing 1-4
 heteroatoms independently selected from N, S, and O and
 25 substituted with 0-3 R¹⁶;

R¹³, R^{13a}, R¹⁴, R^{14a}, and R¹⁵ are independently selected at each
 occurrence from the group: H, =O, COOH, SO₃H, PO₃H, C₁₋₅
 alkyl substituted with 0-3 R¹⁶, aryl substituted with 0-3
 30 R¹⁶, benzyl substituted with 0-3 R¹⁶, and C₁₋₅ alkoxy
 substituted with 0-3 R¹⁶, NHC(=O)R¹⁷, C(=O)NHR¹⁷,
 NHC(=O)NHR¹⁷, NHR¹⁷, R¹⁷, and a bond to Sf;

R¹⁶ is independently selected at each occurrence from the group:
a bond to Sf, COOR¹⁷, C(=O)NHR¹⁷, NHC(=O)R¹⁷, OH, NHR¹⁷,
SO₃H, PO₃H, -OPO₃H₂, -OSO₃H, aryl substituted with 0-3 R¹⁷,
C₁₋₅ alkyl substituted with 0-1 R¹⁸, C₁₋₅ alkoxy
5 substituted with 0-1 R¹⁸, and a 5-10 membered heterocyclic
ring system containing 1-4 heteroatoms independently
selected from N, S, and O and substituted with 0-3 R¹⁷;

R¹⁷ is independently selected at each occurrence from the group:
10 H, alkyl substituted with 0-1 R¹⁸, aryl substituted with
0-1 R¹⁸, a 5-10 membered heterocyclic ring system
containing 1-4 heteroatoms independently selected from N,
S, and O and substituted with 0-1 R¹⁸, C₃₋₁₀ cycloalkyl
substituted with 0-1 R¹⁸, polyalkylene glycol substituted
15 with 0-1 R¹⁸, carbohydrate substituted with 0-1 R¹⁸,
cyclodextrin substituted with 0-1 R¹⁸, amino acid
substituted with 0-1 R¹⁸, polycarboxyalkyl substituted with
0-1 R¹⁸, polyazaalkyl substituted with 0-1 R¹⁸, peptide
substituted with 0-1 R¹⁸, wherein the peptide is comprised
20 of 2-10 amino acids, 3,6-O-disulfo-B-D-galactopyranosyl,
bis(phosphonomethyl)glycine, and a bond to Sf;

R¹⁸ is a bond to Sf;


25 k is selected from 0, 1, and 2;
h is selected from 0, 1, and 2;
h' is selected from 0, 1, and 2;
g is selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;
g' is selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;
30 s is selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;
s' is selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;
s" is selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;
t is selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;

t' is selected from 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, and 10;

x is selected from 0, 1, 2, 3, 4, and 5;

x' is selected from 0, 1, 2, 3, 4, and 5;

5 Sf is a surfactant which is a lipid or a compound of the



A⁹ is selected from the group: OH and OR³²;

10

A¹⁰ is OR³²;

R³² is C(=O)C₁₋₂₀ alkyl;

15 E⁹ is C₁₋₁₀ alkylene substituted with 1-3 R³³;

R³³ is independently selected at each occurrence from the group:

R³⁵, -PO₃H-R³⁵, =O, -CO₂R³⁴, -C(=O)R³⁴, -C(=O)N(R³⁴)₂,
-CH₂OR³⁴, -OR³⁴, -N(R³⁴)₂, C₁-C₅ alkyl, and C₂-C₄ alkenyl;

20


R³⁴ is independently selected at each occurrence from the group:

R³⁵, H, C₁-C₆ alkyl, phenyl, benzyl, and trifluoromethyl;

R³⁵ is a bond to L_n; or

25

Sf is of the formula:



wherein:

A¹ is a bond to Ln;

E¹ is C₁ alkyl substituted by R²³;

A² is NH;

5 E² is C₂ alkyl substituted with 0-1R²³;

A³ is -O-P(O)(R²¹)-O;

E³ is C₁ alkyl;

A⁴ and A⁵ are each -O-;

E⁴ and E⁶ are each independently C₁₋₁₆ alkyl substituted with 0-

10 1R²³;

E⁵ is C₁ alkyl;

A⁵ is -O-;

R²¹ is -OH; and

R²³ is =O; or


15 a pharmaceutically acceptable salt thereof.

74. A diagnostic agent according to Claim 73, wherein:

R is -OH;

R² is C₁₋₆ alkyl;

20 X is C=O;



R¹ and R⁴ are taken together to form a bridging group of formula
-(CH₂)₃-O-phenyl-CH₂-;

R⁵ is NH(C₁₋₆alkyl), substituted with a bond to the linking
25 group or a bond to the surfactant.

75. A diagnostic agent according to Claim 73, wherein:

R is -OH;

R⁹ is C₁ alkyl substituted with a bond to Ln;

30 R¹⁰ and R¹¹ taken together with the nitrogen atom to which they
are attached form a 5 atom saturated ring system, said right
system is substituted with 0-3 R²⁷;

R²⁷ is =O, C₁₋₄ alkyl, or phenyl substituted with R²⁸; and

R²⁸ is a phenoxy group substituted with 0-2 OCH₃ groups;

S_f is a surfactant which is a lipid or a compound of the

5 formula:
$$\begin{array}{c} \text{E}^9 - \text{A}^{10} \\ \diagup \\ \text{A}^9 \end{array};$$

A⁹ is OR³²;

10 A¹⁰ is OR³²;

R³² is C(=O)C₁₋₁₅ alkyl;

E⁹ is C₁₋₄ alkylene substituted with 1-3 R³³;

15 R³³ is independently selected at each occurrence from the group:
R³⁵, -PO₃H-R³⁵, =O, -CO₂R³⁴, -C(=O)R³⁴, -CH₂OR³⁴, -OR³⁴,
and C₁₋₅ alkyl;

R³⁴ is independently selected at each occurrence from the group:
20 R³⁵, H, C₁₋₆ alkyl, phenyl, and benzyl; and

R³⁵ is a bond to L_n.


76. A diagnostic agent according to Claim 73, wherein:
25 R is -OH;

R⁹ is C₁ alkyl substituted with a bond to L_n;

R¹⁰ and R¹¹ taken together with the nitrogen atom to which they
are attached form a 5 atom saturated ring system, said right
system is substituted with 0-3 R²⁷;

30 R²⁷ is =O, C₁₋₄ alkyl, or phenyl substituted with R²⁸; and
R²⁸ is a phenoxy group substituted with 0-2 OCH₃ groups;

Sf is a surfactant which is a lipid or a compound of the of the formula:



5 wherein:

A¹ is a bond to Ln;

E¹ is C₁ alkyl substituted by R²³;

A² is NH;

E² is C₂ alkyl substituted with 0-1R²³;

10 A³ is -O-P(O)(R²¹)-O;

E³ is C₁ alkyl;

A⁴ and A⁵ are each -O-;

E⁴ and E⁶ are each independently C₁₋₁₆ alkyl substituted with 0-1R²³;

15 E⁵ is C₁ alkyl;

A⁵ is -O-;

R²¹ is -OH; and

R²³ is =O.

20 77. A diagnostic agent according to Claim 73, wherein:

wherein

R is -OH;

R¹ and R² taken together with the nitrogen and carbon atom through which they are attached form a C₅₋₇ atom saturated ring system substituted with one or more substituents selected from the group consisting of: a bond to Ln, a bond to Sf, and -C(=O)-NR²⁹R³⁰;

25

R²⁹ and R³⁰ taken together with the nitrogen atom through which they are attached form a C₅₋₇ atom saturated ring system

30 substituted with R³¹; and

R³¹ is a benzyloxy group substituted with C1-4 alkyl.

d is selected from 1, 2, 3, 4, and 5;

5 W is independently selected at each occurrence from the group:

O, NH, NHC(=O), C(=O)NH, NR¹⁵C(=O), C(=O)NR¹⁵, C(=O),
C(=O)O, OC(=O), NHC(=S)NH, NHC(=O)NH, SO₂, (OCH₂CH₂)_s,
(CH₂CH₂O)_{s'}, (OCH₂CH₂CH₂)_{s''}, (CH₂CH₂CH₂O)_t, and (aa)_{t'};

10 aa is independently at each occurrence an amino acid;

Z is selected from the group: aryl substituted with 0-1 R¹⁶,
C₃-10 cycloalkyl substituted with 0-1 R¹⁶, and a 5-10
membered heterocyclic ring system containing 1-4
15 heteroatoms independently selected from N, S, and O and
substituted with 0-1 R¹⁶;

R¹³, R^{13a}, R¹⁴, R^{14a}, and R¹⁵ are independently selected at each
occurrence from the group: H, =O, COOH, SO₃H, C₁-C₅ alkyl
20 substituted with 0-1 R¹⁶, aryl substituted with 0-1 R¹⁶,
benzyl substituted with 0-1 R¹⁶, and C₁-C₅ alkoxy
substituted with 0-1 R¹⁶, NHC(=O)R¹⁷, C(=O)NHR¹⁷,
NHC(=O)NHR¹⁷, NHR¹⁷, R¹⁷, and a bond to Sf;

25 k is 0 or 1;

s is selected from 0, 1, 2, 3, 4, and 5;

s' is selected from 0, 1, 2, 3, 4, and 5;

s'' is selected from 0, 1, 2, 3, 4, and 5; and

t is selected from 0, 1, 2, 3, 4, and 5.

30

78. A diagnostic agent according to Claim 73, wherein:

W¹ is C(=O)NR¹⁵;

h is 1;

- g is 3;
R¹³ and R¹⁴ are independently H;
x is 1;
k is 0;
5 g' is 0;
h' is 1;
W² is NH; and
x' is 1.
- 10 79. A diagnostic agent according to Claim 73, wherein:
x is 0;
k is 1;
Z is aryl substituted with 0-3 R¹⁶;
g' is 1;
15 W² is NH;
R^{13a} and R^{14a} are independently H;
h' is 1; and
x' is 1.
- 20 80. A diagnostic agent according to Claim 73, wherein:
W¹ is C(=O)NR¹⁵;
h is 1;
g is 2;
R¹³ and R¹⁴ are independently H;
25 x is 1;
k is 0;
g' is 1;
R^{13a} and R^{14a} are independently H; or C1-5 alkyl substituted
with 0-3 R¹⁶;
30 R¹⁶ is SO₃H;
W² is NHC(=O) or NH;
h' is 1; and
x' is 2.

81. A diagnostic agent according to Claim 73, wherein:

W¹ is C(=O)NH;

h is 1;

g is 3;

5 R¹³ and R¹⁴ are independently H;

k is 0;

g' is 0;

x is 1;

W² is -NH(C=O)- or -(OCH₂CH₂)₇₆₋₈₄-;

10 h' is 2; and

x' is 1.

82. A diagnostic agent according to Claim 73, wherein:

x is 0;

15 k is 0;

g' is 3;

h' is 1;

W² is NH; and

x' is 1.

20

83. A diagnostic agent according to Claim 73, wherein:

x is 0;

Z is aryl substituted with 0-3 R¹⁶;

k is 1;

25 g' is 1;

R^{13a}R^{14a} are independently H;

W² is NHC(=O) or -(OCH₂CH₂)₇₆₋₈₄-; and

x' is 1.

30 84. A diagnostic agent according to Claim 73, wherein:

W¹ is C=O;

g is 2;


R¹³ and R¹⁴ are independently H;

k is 0;

g' is 0;
 h' is 1;
 w² is NH; and
 x' is 1.

5

85. A diagnostic agent according to Claim 1, wherein the compound is selected from the group consisting of:



10

84. A diagnostic agent according to Claim 48, wherein: wherein the echogenic gas is a perfluorocarbon gas or sulfur hexafluoride.

15

87. A diagnostic agent according to claim 86 wherein said perfluorocarbon is selected from the group consisting of perfluoromethane, perfluoroethane, perfluoropropane, perfluorobutane, perfluorocyclobutane, perfluoropentane, and perfluorohexane.

20

88. A diagnostic composition comprising a compound according to claim 48 or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier.

25

89. A diagnostic composition comprising a compound according to claim 48 or a pharmaceutically acceptable salt form

thereof, an echogenic gas and a pharmaceutically acceptable carrier.

- 5 90. A diagnostic composition comprising a compound according to claim 48 further comprising: 1,2-dipalmitoyl-sn-glycero-3-phosphotidic acid, 1,2-dipalmitoyl-sn-glycero-3-phosphatidylcholine, and N-(methoxypolyethylene glycol 5000 carbamoyl)-1,2-dipalmitoyl-sn-glycero-3-phosphatidylethanolamine.
- 10 91. A method of detecting, imaging or monitoring the presence of matrix metalloproteinase in a patient comprising the steps of:
- 15 a) administering to said patient a diagnostic agent of claim 1; and
- b) acquiring an image of a site of concentration of said diagnostic agent in the patient by a diagnostic imaging technique.
- 20 92. A method of detecting, imaging or monitoring the presence of matrix metalloproteinase in a patient comprising the steps of:
- a) administering to said patient a diagnostic agent of claim 48; and
- 25 c) acquiring an image of a site of concentration of said diagnostic agent in the patient by a diagnostic imaging technique.
- 30 93. A method of detecting, imaging or monitoring a pathological disorder associated with matrix metalloproteinase activity in a patient comprising the steps of:
- a) administering to said patient a diagnostic agent of claim 1; and
- 35 b) acquiring an image of a site of concentration of said diagnostic agent in the patient by a diagnostic imaging technique.

94. A method of detecting, imaging or monitoring a pathological disorder associated with matrix metalloproteinase activity in a patient comprising the steps of:
- 5 a) administering to said patient a diagnostic agent according to claim 48; and
- c) acquiring an image of a site of concentration of said diagnostic agent in the patient by a diagnostic imaging technique.
- 10
95. A method of detecting, imaging or monitoring atherosclerosis in a patient comprising the steps of:
- a) administering a diagnostic agent according to claim 1; and
- 15 b) acquiring an image of a site of concentration of said diagnostic agent in the body by a diagnostic imaging technique.
96. A method of detecting, imaging or monitoring atherosclerosis in a patient comprising the steps of:
- 20 c) administering a diagnostic agent according to claim 48; and
- d) acquiring an image of a site of concentration of said diagnostic agent in the body by a diagnostic imaging technique.
- 25
97. A method according to claim 95, wherein the atherosclerosis is coronary atherosclerosis or cerebrovascular atherosclerosis.
- 30 98. A method according to claim 96, wherein the atherosclerosis is coronary atherosclerosis or cerebrovascular atherosclerosis.
99. A method of identifying a patient at high risk for transient ischemic attacks or stroke by determining the degree of active atherosclerosis in a patient comprising
- 35 carrying out the method of claim 96.

100. A method of identifying a patient at high risk for transient ischemic attacks or stroke by determining the degree of active atherosclerosis in a patient comprising carrying out the method of claim 97.
101. A method of identifying a patient at high risk for acute cardiac ischemia, myocardial infarction or cardiac death by determining the degree of active atherosclerosis by imaging the patient by the method of claim 96.
102. A method of identifying a patient at high risk for acute cardiac ischemia, myocardial infarction or cardiac death by determining the degree of active atherosclerosis by imaging the patient by the method of claim 97.
103. A method of simultaneous imaging of cardiac perfusion and extracellular matrix degradation in a patient comprising the steps of:
- a) administering a diagnostic agent according to claim 1, wherein the diagnostic metal is a gamma-emitting radioisotope; and
 - (b) administering a cardiac perfusion compound, wherein the compound is radiolabeled with a gamma-emitting radioisotope which exhibits a gamma emission energy that is spectrally separable from the gamma emission energy of the diagnostic metal conjugated to the targeting moiety in step (a); and
 - (c) acquiring, by a diagnostic imaging technique, simultaneous images of the sites of concentration of the spectrally separable gamma-emission energies of the compounds administered in steps (a) and (b) .

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

☐ BLACK BORDERS

☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES

☐ FADED TEXT OR DRAWING

☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING

☐ SKEWED/SLANTED IMAGES

☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS

☐ GRAY SCALE DOCUMENTS

☐ LINES OR MARKS ON ORIGINAL DOCUMENT

☒ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY

☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.